

Invading, Leading and Navigating Cells in *Caenorhabditis elegans*: Insights into Cell Movement *in Vivo*

David R. Sherwood^{*,1} and Julie Plastino^{†,1}

^{*}Department of Biology, Regeneration Next, Duke University, Durham, North Carolina 27705, [†]Institut Curie, PSL Research University, CNRS, UMR 168, F-75005 Paris, France, ¹Sorbonne Universités, UPMC Univ Paris 06, CNRS, UMR 168, F-75005 Paris, France

ORCID ID: 0000-0002-4448-6917 (D.R.S.)

ABSTRACT Highly regulated cell migration events are crucial during animal tissue formation and the trafficking of cells to sites of infection and injury. Misregulation of cell movement underlies numerous human diseases, including cancer. Although originally studied primarily in two-dimensional *in vitro* assays, most cell migrations *in vivo* occur in complex three-dimensional tissue environments that are difficult to recapitulate in cell culture or *ex vivo*. Further, it is now known that cells can mobilize a diverse repertoire of migration modes and subcellular structures to move through and around tissues. This review provides an overview of three distinct cellular movement events in *Caenorhabditis elegans*—cell invasion through basement membrane, leader cell migration during organ formation, and individual cell migration around tissues—which together illustrate powerful experimental models of diverse modes of movement *in vivo*. We discuss new insights into migration that are emerging from these *in vivo* studies and important future directions toward understanding the remarkable and assorted ways that cells move in animals.

KEYWORDS basement membrane; *C. elegans*; cell invasion; cell migration; cell signaling; F-actin; FGF pathway; integrin; netrin pathway; Wnt pathway; WormBook

TABLE OF CONTENTS

Abstract	53
AC Invasion: Breaching BM Barriers	55
<i>The AC breaches the BM with invadopodia</i>	55
<i>A large invasive protrusion clears an opening in the BM</i>	57
<i>Identification of the transcriptional networks that specify cell invasive behavior</i>	58
<i>The AC and vulval cells collaborate to further widen the BM opening</i>	58
<i>AC invasion: important unanswered questions</i>	59
Distal Tip Cell Migration: a Leader Cell Migration that Shapes a Tissue	61
<i>DTC modification of and interaction with BM during migration</i>	61
<i>Stabilizing the DTC path: the matrix proteins HIM-4 (hemicentin) and DIG-1</i>	63

Continued

Copyright © 2018 by the Genetics Society of America

doi: <https://doi.org/10.1534/genetics.117.300082>

Manuscript received July 29, 2017; accepted for publication October 26, 2017

¹Corresponding authors: Department of Biology, Regeneration Next, Duke University, Box 90338 Science Dr., Durham, NC 27708. E-mail: david.sherwood@duke.edu; and Institut Curie, UMR 168, F-75005 Paris, France. E-mail: julie.plastino@curie.fr

CONTENTS, *continued*

<i>Diffusible cues help orient DTC migration: netrin and Wnts</i>	64
<i>Polarity of the nucleus and cytoskeleton during DTC migration</i>	65
<i>Transcriptional regulation of DTC migration</i>	65
<i>microRNAs confer robustness to DTC migration under temperature stress</i>	66
<i>DTC migration: key unanswered questions</i>	67
Sex Myoblast Migration: Navigating Around Tissues To Position Muscles	68
<i>Attraction and repulsion in SM migration</i>	68
<i>Ras in SM positioning</i>	69
<i>FGF independent mechanisms of SM migration</i>	70
<i>Possible signaling to the actin cytoskeleton for SM migration</i>	70
<i>Specification of migratory SM cells</i>	70
<i>Unanswered questions in SM migration</i>	71
<i>Summary and outlook</i>	71

THE ability of cells to move is crucial for many aspects of normal animal embryogenesis, organ formation, wound healing, tissue regeneration, and immune cell trafficking (Aman and Piotrowski 2010; Nourshargh and Alon 2014; Mayor and Etienne-Manneville 2016). Misregulation of cell movement also underlies human developmental disorders, immune dysfunction, and cancer (Kurosaka and Kashina 2008; Madsen and Sahai 2010; Friedl and Alexander 2011; Paul *et al.* 2017). Thus, understanding the mechanisms by which cells move in diverse cell and tissue environments has important basic and clinical relevance. Because of the challenge of examining dynamic cellular behaviors in native tissue settings, most studies of cell migration have been carried out in cell culture. While these *in vitro* studies have revealed mechanisms underlying key parameters of migration, such as cytoskeletal regulation, cell-cell and cell-extracellular matrix (ECM) adhesion, polarization machinery, and distinct modes of migration (Lammermann and Sixt 2009; Linder *et al.* 2011; Blanchoin *et al.* 2014; Te Boekhorst *et al.* 2016), *in vitro* conditions do not faithfully match the complexity of *in vivo* settings, and, therefore, their physiological significance often remains unclear.

The shortcomings of *in vitro* migration models are highlighted by the fact that cell-substrate adhesions and other cellular structures appear very different in cells plated on two-dimensional (2D) flat, rigid substrates as compared to more native three-dimensional (3D) cell and ECM environments, and often display different dynamics and biochemistry (Fraley *et al.* 2010; Geraldo *et al.* 2012; Petrie *et al.* 2012). Although 3D culture conditions are a step in the right direction, they do not reflect the richness of other physiologically relevant environmental factors that migrating cells encounter. These factors include diverse cell-cell interactions, diffusible cues, fluctuating nutrient conditions, changing oxygen levels, varying fluid dynamics, cell and tissue growth, and

native mechanical properties of cells and extracellular matrices (Even-Ram and Yamada 2005; Friedl *et al.* 2012). Cells also have important intrinsic properties, such as unique transcriptional programs and chromatin states, that are likely not recapitulated in cell culture settings (Feil and Fraga 2012; Chen *et al.* 2013). Thus, *in vivo* models are essential, not only to verify or challenge mechanisms discovered *in vitro*, but also to discover new mechanisms of cell migration that are difficult, if not impossible, to recapitulate *in vitro*.

Studying cell movements in *Caenorhabditis elegans* provides a strong experimental model to examine cell motility in an *in vivo* setting. One of the advantages of studying cell migration in *C. elegans* is the simplicity of the gene families that encode cytoskeleton (Sawa *et al.* 2003; Schonichen and Geyer 2010; Mi-Mi *et al.* 2012; Abella *et al.* 2016; Pizarro-Cerda *et al.* 2017), ECM (Kramer 2005), and signaling proteins (Lai Wing Sun *et al.* 2011; Clevers and Nusse 2012; Sawa and Korswagen 2013) that guide cell migrations. This simplified genetic landscape reduces redundancy and makes gene perturbation studies easier to perform and interpret. Cell migration phenotypes are also straightforward to visualize, as the worm's optical transparency allows for imaging of all cell migrations in real time. In addition, *C. elegans* anatomical simplicity (the adult has <1000 somatic cells) and its highly stereotyped development facilitate detailed analysis of even subtle phenotypes. *C. elegans* is also remarkably easy to manipulate genetically such that genes and proteins can be altered at the organismal and individual cell level using temporally controlled optogenetic, RNAi, CRISPR/Cas-9, and ubiquitin mediated methods (Hagedorn *et al.* 2009; Dickinson *et al.* 2013; Armenti *et al.* 2014; Shen *et al.* 2014; Corsi *et al.* 2015). Finally, the worm's short life cycle and hermaphrodite mode of reproduction coupled with rapid whole-genome RNAi screening facilitate discovery of genes and pathways regulating cell migration that would not be found through

candidate approaches (Jorgensen and Mango 2002; Kamath *et al.* 2003; Corsi *et al.* 2015). Together, these worm attributes permit exceptional experimental access to uncover the molecular and cell biological mechanisms that underlie migration *in vivo*.

C. elegans undergoes numerous cell migrations throughout embryonic and larval development (Hedgecock *et al.* 1987). Much information concerning mechanisms underlying cell migration in *C. elegans* has emerged from the study of a few major motile events. Some of these have recently been reviewed elsewhere, including ventral enclosure (Vuong-Brender *et al.* 2016), Q neuroblast migration (Rella *et al.* 2016) and axon guidance (Chisholm *et al.* 2016). Our review focuses on what has been learned and promising future studies on three distinct cellular movements that are common motility modes in animals: anchor cell (AC) invasion as a model for invasion through basement membrane (BM) barriers; distal tip cell (DTC) migration as a model for how a BM-encased leader cell directs organ formation; and sex myoblast (SM) migration as a model for how cells migrate between tissues.

AC Invasion: Breaching BM Barriers

BMs are thin, dense, highly cross-linked ECM composed of interlinked sheets of laminin and type IV collagen networks that surround and support most tissues (Yurchenco 2011; Jayadev and Sherwood 2017). Despite their barrier properties, BMs are breached and crossed by cells during development, blood vessel formation, and immune functioning (Yang and Weinberg 2008; Kelley *et al.* 2014; Seano *et al.* 2014). Inappropriate invasion also underlies numerous pathologies, most notably cancer cell metastasis (Valastyan and Weinberg 2011). Owing to the complexity of studying dynamic interactions between invasive cells, BMs, and the invaded tissue, cell invasion has been challenging to experimentally examine in native tissue environments (Beerling *et al.* 2011; Hagedorn and Sherwood 2011).

Most *C. elegans* tissues are enwrapped in BM, and the *C. elegans* genome harbors the major BM components laminin and type IV collagen, as well as the BM-associated proteins perlecan, nidogen, fibulin, agrin, hemicentin, SPARC, and collagen XVIII (Kramer 2005). Gene families encoding BM proteins in *C. elegans* have not undergone the extensive expansion observed in vertebrates (Kramer 2005), and many have been tagged with fluorophores allowing analysis of their localization and function (Kelley *et al.* 2014). Notably, *C. elegans* lacks interstitial matrix, and does not harbor genes encoding fibrillar collagens and other interstitial components (Hutter *et al.* 2000). Fibrillar collagens and interstitial matrix are thought to have originated near the time of metazoan emergence, but apparently were lost in the lineage that gave rise to *C. elegans* (Ozbek *et al.* 2010; Fidler *et al.* 2017).

C. elegans AC invasion into the vulval epithelium is an *in vivo* model of invasive behavior that permits single cell and subcellular analysis of invasion through BM (Figure 1). The AC is a specialized uterine cell that invades through BM

separating the uterine and vulval tissue to initiate uterine-vulval attachment—a connection required for mating and laying embryos. AC invasion occurs over a precise 90-min period during the L3 larval stage, and is coordinated with the divisions of the descendants of the P6.p epidermal cell. These cells are the 1° vulval precursor cells, and give rise to the centrally located vulval cells. The stereotyped nature of AC invasion, amenability to forward and reverse genetic screens, and visual accessibility to live-cell imaging have allowed mechanisms regulating invasion to be identified and characterized *in vivo*.

The AC breaches the BM with invadopodia

Studies on AC invasion have revealed that dynamic, ~1.0 μ m-diameter F-actin (filamentous actin) rich structures, termed invadopodia, form along the AC's invasive cell membrane and breach the BM (Figure 1) (Hagedorn *et al.* 2013). Matrix degrading invadopodia were originally observed in transformed fibroblasts, human cancer cell lines, and primary tumor cells from human patients cultured on glass slides covered with simplified ECMs *in vitro* (Chen 1989; Linder *et al.* 2011; Genot and Gligorijevic 2014). Invadopodia have been extensively characterized *in vitro*, and numerous aspects of their composition, regulation, and formation have been elucidated in cell culture settings (Bergman *et al.* 2014). The identification of invadopodia in *C. elegans* has confirmed the *in vivo* existence of these structures, the importance of invadopodia in breaching BM, and regulation of invadopodia in native invasion events (Morrissey *et al.* 2013; Genot and Gligorijevic 2014; Lohmer *et al.* 2014).

AC invadopodia formation is stimulated ~5 hr prior to invasion by an unidentified diffusible cue(s) from the 1° fated vulval precursor cells (Sherwood and Sternberg 2003; Lohmer *et al.* 2016). The vulval signal activates the Rho GTPase CDC-42 (vertebrate Cdc42) within the AC (Figure 1). CDC-42 seeds invadopodia in part through its effector WSP-1 (WASP), which activates the Arp2/3 complex, an actin polymerization nucleator (Shakir *et al.* 2008; Padrick and Rosen 2010; Lohmer *et al.* 2016). Approximately 10 invadopodia are present at any one time in the AC, and turn over rapidly with a median life time of 45 sec (Hagedorn *et al.* 2013). The rapid turnover of AC invadopodia is in stark contrast to invadopodia dynamics characterized *in vitro* in cancer cells, which have average lifetimes of 30 min or longer (Linder *et al.* 2011; Branch *et al.* 2012; Moshfegh *et al.* 2014). The rapid turnover of AC invadopodia might reflect the more physiologically relevant *in vivo* conditions (e.g., BM composition, microenvironmental signals, and intrinsic cell characteristics) or that AC invasion has evolved to be a quick invasion event, whereas cancer cells are less efficient in trans-migrating BMs (Lohmer *et al.* 2014).

Other actin regulatory proteins are also associated with AC invadopodia, and have revealed important aspects of their formation and regulation. The actin filament severing protein UNC-60 (ADF/cofilin) is crucial to invadopodia turnover (Figure 1). In the absence of UNC-60, invadopodia formation




Figure 1 AC invasion, a BM invasion event. Prior to invasion, the AC is positioned over the epidermal P6.p cell, which it induces to a 1° vulval precursor cell fate. The 1° fated P6.p divides three times. Invasion occurs at the time of the division from the P6.p 2-cell stage to P6.p 4-cell stage. Top panel: Prior to breaching the BM, AC invadopodia form along the AC's invasive cell membrane, depress the BM, and then disassemble. Approximately 10 invadopodia are present at any one time. Invadopodia formation is stimulated by a cue from the 1° vulval cells that activates the Rho GTPase CDC-42. CDC-42 activates WSP-1, which presumably stimulates F-actin production through actin polymerization nucleators such as the Arp2/3 complex. UNC-34 (Ena/VASP) may also contribute to F-actin formation. Invadopodia generation is also dependent on an invadopodial membrane rich in PI(4,5)P2 and containing the lipid-anchored Rac GTPases, CED-10 and MIG-2. The invadopodial membrane is recycled through the endolysosome, and its trafficking is dependent on UNC-60 (cofilin) and the Rab GDP dissociation inhibitor (GDI-1). The integrin heterodimer INA-1/PAT-3 is required for trafficking of all known invadopodial components to the plasma membrane. Middle panel: when an invadopodium breaches the BM, the netrin receptor UNC-40 (DCC) traffics to the breach site, and is activated by its ligand UNC-6 (netrin) secreted by the underlying ventral nerve cord. UNC-40 recruits the actin regulators UNC-34 (Ena/VASP) and the Rac GTPases, which shuts down further invadopodia formation. Lower Panel: UNC-40 (DCC) directs the formation of a large invasive protrusion that expands the opening in the BM by degrading and physically displacing BM.

ceases, and instead large aggregates of static F-actin form along the invasive cell membrane (Hagedorn *et al.* 2014). UNC-34 (Ena/VASP), another actin regulator associated with enhanced cell migration, is also localized to AC invadopodia, although its function in the AC has not been characterized (Chesarone and Goode 2009; Hagedorn *et al.* 2013). In addition to F-actin,

invadopodia are constructed from a specialized invadopodial membrane containing the lipid phosphatidyl inositol 4,5 bisphosphate (PI(4,5)P2) and CED-10 and MIG-2, two members of the Rac subfamily of Rho GTPases that have lipid anchors (Lundquist *et al.* 2001; Hagedorn *et al.* 2014; Ridley 2015; Reiner and Lundquist 2016). The invadopodial membrane is

actively recycled through the endolysosome during invadopodia formation and breakdown, and its delivery to the invasive cell membrane is dependent on the Rab GDP dissociation inhibitor **GDI-1** as well as **UNC-60** (cofilin) (Hagedorn *et al.* 2014; Lohmer *et al.* 2016). Cell culture studies with cancer cells have indicated that the membrane tethered matrix metalloproteinase MT1-MMP is delivered to invadopodia through the endolysosome, suggesting that active endolysosome recycling might be a shared feature of invadopodia formation (Castro-Castro *et al.* 2016; Hastie and Sherwood 2016).

Invadopodia formation also requires the ECM integrin receptor heterodimer **INA-1/PAT-3** (Hagedorn *et al.* 2009). Integrin receptors are composed of an α and a β subunit. Unlike vertebrates, however, which encode 18 α and 8 β subunits and construct 24 known $\alpha\beta$ integrin heterodimers, *C. elegans* has only a single β and two α subunits and make only two integrins: α **INA-1**/ β **PAT-3**, most similar to vertebrate laminin binding integrins, and α **PAT-2**/ β **PAT-3**, an RGD containing integrin (Baum and Garriga 1997; Kramer 2005; Campbell and Humphries 2011). **INA-1/PAT-3** is the only integrin expressed in the AC, and mediates the trafficking of F-actin regulators and invadopodial membrane to the invasive cell membrane (Hagedorn *et al.* 2009). **INA-1/PAT-3** might polarize the secretory apparatus of the AC (Wickstrom and Fassler 2011).

Invadopodia breach the BM during a narrow 20-min time period in the mid-L3 larval stage. These observations suggest that invadopodia formed in the 5-hr window prior to breach might not be fully mature and able to penetrate BM. The mechanisms that mediate the maturation/precise timing of the invadopodia breach are not known, but are likely connected to cues generated by the vulval cells, as precocious vulval cell formation accelerates the timing of AC invasion (Sherwood and Sternberg 2003). The AC expresses three matrix metalloproteinases (**zmp-1**, **zmp-3**, and **zmp-6**), a class of proteolytic enzymes that have been implicated in vertebrates in breaking down BMs to promote invasion (Sherwood *et al.* 2005; Hotary *et al.* 2006; Page-McCaw *et al.* 2007; Itoh 2015; Matus *et al.* 2015). Invadopodia also physically deform the BM as they extend, suggesting that a combination of proteolysis and mechanical disruption mediates invadopodial breaching of the BM (Figure 1) (Hagedorn *et al.* 2013).

The similarities of invadopodia regulation in cancer cells and the AC are striking. In numerous cancer cell lines Cdc-42 stimulates invadopodia formation (Di Martino *et al.* 2014; Razidlo *et al.* 2014), cofilin regulates invadopodia turnover (Bravo-Cordero *et al.* 2013; Beaty and Condeelis 2014), and integrins are essential for invadopodia generation (Destaing *et al.* 2011; Murphy and Courtneidge 2011). These observations suggest that invadopodia are conserved subcellular structures that evolved early in animals to allow cells to pass through BM barriers during development and tissue remodeling (Medwig and Matus 2017). A recent sensitized whole genome RNAi screen has identified many additional genes implicated in invadopodia regulation, including genes

involved in G-protein signaling, extracellular matrix remodeling and the Hippo pathway (Lohmer *et al.* 2016). The characterization of these genes will provide new insight into mechanisms mediating invadopodia formation, turnover, and BM breaching *in vivo*.

A large invasive protrusion clears an opening in the BM

Invadopodia formation, turnover, and matrix breaching can be reasonably well recapitulated in cell culture on flat 2D surfaces such as ECM coated glass slides (Even-Ram and Yamada 2005). Capturing later events in BM invasion *in vitro* is challenging, as this requires 3D assays that faithfully mimic native BM and cellular architecture to follow how cells cross BM barriers and enter new tissues (Schoumacher *et al.* 2010, 2011). *C. elegans* AC invasion has been particularly crucial in extending our understanding of BM invasion following invadopodia-mediated BM breaching events.

Live-cell imaging revealed that usually only one or two AC invadopodia breach the BM and only one of these transitions into a protrusion that clears a large opening in the BM and expands between the central vulval cells (Hagedorn *et al.* 2013). The netrin pathway plays a crucial role in directing this dramatic transition (Morrissey *et al.* 2013). The netrin receptor **UNC-40** (vertebrate DCC) clusters at the invadopodial breach site and in response to **UNC-6** (netrin) recruits F-actin regulators (Ena/VASP, Rho GTPases) that direct invasive protrusive formation (Figure 1) (Hagedorn *et al.* 2013). **UNC-40** acts as a molecular sink and depletes F-actin regulators from other invadopodia, shutting them down and focusing F-actin generation and invasion through a single breach site. In the absence of **UNC-40** (DCC), invasive protrusion formation fails and invadopodia continue to form and turn over, creating multiple small breaches in the BM that hinder the ability of the AC to contact the vulval cells. This state likely mimics 2D culture conditions, where numerous holes are generated by tumor cell invadopodia on matrix covered glass slides (Martin *et al.* 2012).

Studies on AC invasion have also uncovered new mechanisms of how **UNC-40** (DCC) receptors polarize toward sources of netrin. Surprisingly, **UNC-40** (DCC) is still active in animals lacking **UNC-6** (netrin): **UNC-40** receptors randomly cluster in the cell membrane, recruit F-actin effectors, and generate F-actin. These transient clusters then break down and reform in a new location in an oscillatory cycle (Wang *et al.* 2014b). **UNC-6** (netrin), which is localized in the BM and below it (secreted by the ventral nerve cord and later by the vulval cells) stabilizes **UNC-40** (DCC) clustering toward the source of **UNC-6** (netrin), thus directing protrusion formation through the BM and between the vulval cells (Wang *et al.* 2014b). This oscillatory behavior is likely a mechanism that allows **UNC-40** (DCC) receptors to rapidly and robustly polarize toward sources of **UNC-6** (netrin) and is probably a universal feature of **UNC-40** (DCC) receptor polarization that is shared with other polarity systems (Bendezu and Martin 2013; Dyer *et al.* 2013; Kulkarni *et al.* 2013; Wang *et al.* 2014b).

Optical highlighting of the BM components laminin and type IV collagen revealed that the AC's invasive protrusion utilizes a combination of proteolysis and physical displacement to both degrade and push aside the BM (Figure 1) (Hagedorn *et al.* 2013). This was an unforeseen finding, as BM invasion was not known to involve physical forces and instead was thought to rely solely on proteases dissolving the BM to clear a path for invasion (Valastyan and Weinberg 2011). How the AC's invasive protrusion generates forces to push aside the BM remains an important area of future investigation (see below).

Identification of the transcriptional networks that specify cell invasive behavior

Study of AC invasion has allowed the transcriptional networks to be identified that endow cells with the specialized ability to breach BM (Figure 2). The AC is first specified during the late L2 larval stage (~8 hr prior to invasion). Specification requires the helix-loop-helix transcription factor protein **HLH-2** (Daughterless/vertebrate E proteins) and the nuclear hormone **NHR-67** (vertebrate TLX) (Karp and Greenwald 2004; Schindler and Sherwood 2011; Verghese *et al.* 2011). These transcription factors are expressed prior to AC specification and throughout its differentiation, and appear to belong to a transcriptional network that regulates distinct transcriptional targets at different stages (Schindler and Sherwood 2011). During early specification, **NHR-67** is required to express the gene encoding the Notch ligand **LAG-2** and **HLH-2** promotes expression of the FAT-like cadherin **CDH-3** (Schindler and Sherwood 2011; Verghese *et al.* 2011), while unknown factors are responsible for upregulation of other invasive genes encoding the β -integrin subunit **PAT-3**, and the Rho GTPase **MIG-2** (Sherwood and Sternberg 2003; Sherwood *et al.* 2005; Ziel *et al.* 2009).

After the AC is specified, **NHR-67** (TLX) transcription factor expression is further upregulated and induces G1 cell-cycle arrest (Matus *et al.* 2015). During development, G1 cell-cycle arrest is thought to allow cells to engage unique transcriptional programs to facilitate differentiation (Gonzales *et al.* 2015; Ruijtenberg and van den Heuvel 2015). This also appears to be the case in the AC as cell-cycle arrest is required for the AC to express genes associated with invasion and adopt the specialized features of an invasive cell. In the absence of G1 arrest, early aspects of AC specification occur normally, including expression of genes encoding **PAT-3** (integrin) and **CDH-3** (cadherin); however, later stages of invasive differentiation fail to occur. For example, the AC does not express genes encoding actin regulators such as the formin **EXC-6** and the Ena/VASP ortholog **UNC-34**, the matrix metalloproteinases **ZMP-1**, -3 and -6, and the matrix component **HIM-4** (hemicentin). In addition, invadopodia do not form (Matus *et al.* 2015). G1 cell cycle arrest might be a common feature of invasive cells, as invasive ability is correlated with G1 arrest in other developmental events involving BM transmigration and with invasive tumor cells (Kohrman and Matus 2017). This may be particularly important in treating metastatic cancers. As most chemotherapy drugs target dividing

cells (Yano *et al.* 2014), such treatments would not be effective against nondividing invasive cells, and could select for more aggressive tumors by leaving invasive cells unscathed and later able to proliferate and seed new metastatic lesions.

G1 arrest is thought to permit cell-cycle dependent alterations in chromatin that allow the expression of differentiation genes (Ma *et al.* 2015). There is evidence in the AC that this might be mediated through the histone deacetylase **HDA-1** and the zinc finger protein **MEP-1**, a component of the nucleosome remodeling NuRD complex. **HDA-1** and **MEP-1** are required for AC invasion and the expression of genes associated with later aspects of AC differentiation (Matus *et al.* 2010, 2015). An emerging transcription factor network is being identified that acts during G1 arrest (Figure 2). This network includes the conserved bZIP transcription factor **FOS-1A** (Fos), **HLH-2** (Daughterless/vertebrate E proteins), and the zinc finger protein **EGL-43B** (vertebrate EVI1). These transcription factors regulate the expression of genes encoding invasion effectors such as matrix metalloproteinases (*e.g.*, **ZMP-1**), extracellular matrix proteins [*e.g.*, **MIG-6** (papilin) and **HIM-4** (hemicentin)], and actin cytoskeleton proteins [*e.g.*, **MIG-10B** (lamellipodin); Sherwood *et al.* 2005; Hwang *et al.* 2007; Rimann and Hajnal 2007; Schindler and Sherwood 2011; L. Wang *et al.* 2014]. Interesting aspects of this network are beginning to emerge, such as an incoherent feed-forward circuit where **FOS-1A** positively controls expression of the genes encoding **MIG-10B** and the transcription factor **EGL-43B**, while the **EGL-43B** protein negatively regulates *mig-10b* gene expression. Such networks likely provide fine control over the expression of key effector targets that promote invasion (L. Wang *et al.* 2014). The vertebrate Fos family of transcription factors are strongly implicated in promoting cell motility and invasion in normal development and multiple tumor types (Milde-Langosch 2005; Ozanne *et al.* 2007; Renaud *et al.* 2014). Further, vertebrate E proteins (**HLH-2** orthologs) promote epithelial-mesenchymal transitions (EMTs) (Lamouille *et al.* 2014), which often involve breaching epithelial BMs. Thus, the transcriptional networks that program invasiveness might be conserved. Additional transcriptional mechanisms controlling AC invasion remain to be discovered, as many pro-invasive genes, such as those encoding **MIG-2** (Rac), **PAT-3** (integrin) and **CDC-37** (Hsp 90 cochaperone), are upregulated in the AC, but are not controlled by any known transcriptional regulators (Sherwood *et al.* 2005; Matus *et al.* 2010; Schindler and Sherwood 2011).

The AC and vulval cells collaborate to further widen the BM opening

Following AC invasion, the BM opening enlarges past the edge of the AC. Widening the BM gap is crucial to allow direct attachment between the uterine and vulval cells that form the mature uterine-vulval connection. Optical highlighting of the BM components laminin and type IV collagen revealed that the BM moves over the growing vulval and uterine tissues to widen the BM gap through a newly described morphogenetic




Figure 2 Transcription factors that specify invasive cell fate. Left: The AC is specified during the late L2 stage by the action of several transcription factors, including NHR-67 (Tailless) and HLH-2 (Daughterless/E proteins) that drive the expression of genes encoding LAG-2 (Notch ligand) and CDH-3 (cadherin), respectively, as well as other unknown transcription factors that promote the expression of genes encoding PAT-3 (β -integrin) and MIG-2 (Rac). Right: During the early L3 stage NHR-67 directs the AC into G1 cell cycle arrest, which allows full invasive fate differentiation. A central transcription factor that operates following G1 arrest is FOS-1A (Fos), which promotes the expression of the transcription factors HLH-2 and EGL-43B (EVI1). These transcription factors regulate the expression of several inva-

sion effector genes that encode MIG-6 (papilin), MIG-10B (lamellipodin), HIM-4 (hemicentin), and ZMP-1 (MMP). Other transcription factors remain to be discovered as genes encoding proteins such as CDC-37 (Hsp90 cochaperone) are expressed in the AC and promote invasion, but are not regulated by the FOS-1A transcriptional network.

mechanism termed BM sliding (Ihara *et al.* 2011). It is thought that the rapid growth of the uterine and vulval tissue (approximately twofold growth during time of sliding) applies forces on the BM that drive its shifting (Ihara *et al.* 2011). In addition, the vulval cells invaginate, and move through the BM opening, which may further shift BM position (Ihara *et al.* 2011; Schindler and Sherwood 2013).

Both vulval cells and uterine cells act to control BM shifting to precisely enlarge the opening in the BM (Figure 3). On the vulval side, the centrally positioned vulE and vulF cells divide, and the rounding of the dividing cells causes these cells to lose contact with the BM, allowing the BM to slide over them to expand the BM gap (Matus *et al.* 2014). The BM position stabilizes on the nondividing vulD cell, where the integrin adhesion receptor INA-1/PAT-3 and VAB-19 (a cytosolic adaptor protein and ortholog of the vertebrate tumor suppressor Kank) localize at high levels to the vulD-BM interface, locking the position of the BM gap boundary (Ding *et al.* 2003; Ihara *et al.* 2011). The vulD cell is the only vulval cell that does not divide during the time of BM sliding in all rhabditid nematodes that have been examined (Kiontke *et al.* 2007), a group of nematodes that last shared a common ancestor 240–430 MYA (Dieterich *et al.* 2008). Examination of over 20 of these rhabditid species revealed that the BM gap always stabilizes over the vulD cell, suggesting that control of vulval cell division is a robust and evolutionary conserved mechanism to control BM gap enlargement.

The invading AC and neighboring uterine cells also have a role in promoting BM gap enlargement. During its invasion, the AC signals with the transmembrane Notch ligand LAG-2 to neighboring uterine π cells through the Notch receptor LIN-12 to upregulate the expression of the gene *ctg-1*, which encodes a Sec14 family phosphatidylinositol-transfer protein (Tripathi *et al.* 2014; McClatchey *et al.* 2016). During LIN-12 (Notch) activation, the intracellular domain of LIN-12 (Notch) is proteolytically cleaved (Notch intracellular domain, NICD) and

enters the nucleus, where it forms a complex with the DNA binding protein LAG-1 (CSL), which promotes LIN-12 (Notch) effector gene expression (Greenwald 2005). The *ctg-1* gene contains 19 putative LAG-1 binding sites, strongly suggesting it is a direct target of LIN-12 (Notch) signaling (Yoo *et al.* 2004). The CTG-1 protein limits the trafficking of the BM-adhesion receptor dystroglycan to the uterine cell-BM interface, allowing the BM to slide over the uterine cells to the position determined by the nondividing vulD on the other side of the BM (Figure 3) (Matus *et al.* 2014; McClatchey *et al.* 2016). The temporally coordinated mechanisms of tissue growth, movement, division, dystroglycan receptor downregulation in uterine cells, and integrin and VAB-19 upregulation in vulval cells at the BM gap edge act to precisely position the BM gap boundary to allow direct uterine-vulval tissue attachment.

The shifting of cell-BM interfaces has been observed in several important morphogenetic events in other organisms, including intestinal epithelial renewal, BM deposition, and branching morphogenesis (Haigo and Bilder 2011; Clevers and Battle 2013; Harunaga *et al.* 2014). Thus, the mechanisms that slide the BM after AC invasion might be used in other important developmental processes to mediate BM remodeling. Further, as dystroglycan loss is a common occurrence in the progression of many epithelia cancers, such as breast and colon (Sgambato *et al.* 2003; Cross *et al.* 2008), it is possible that its loss in cancer allows BM openings to widen further, permitting more extensive tumor cell spread.

AC invasion: important unanswered questions

MMP expression is strongly associated with cell invasion in normal development and cancer (Overall and Kleinfeld 2006; Srivastava *et al.* 2007; Page-McCaw 2008). Further, experimental work with *in vitro* and *ex vivo* models suggest MMPs are essential for BM transmigration (Rowe and Weiss 2008). As a result of their strong association with invasion, MMPs have been targeted in extensive clinical trials, which were




Figure 3 BM sliding following AC invasion widens the breach. Top panel (P6.p 4-cell stage): during invasion, the AC activates LIN-12 (Notch) signaling in neighboring uterine π cells via the ligand LAG-2 (Delta; see upper right box). Notch activation leads to proteolytic release of the LIN-12 (Notch) intracellular domain (NICD), which enters the nucleus, associates with LAG-1 (CSL), and upregulates expression of the gene encoding CTG-1 (Sec14-GOLD protein). Middle panel left (P6.p 6-cell stage): vulval and uterine tissue growth and vulval cell invagination apply forces on the BM that drive its shifting. VulvF cells begin to invaginate and the vulvE cells divide, lose contact with the BM, and allow the BM to slide over these cells, thus widening the BM gap. Middle panel right: CTG-1 activity in the uterine π cells inhibits the trafficking of the BM adhesion receptor dystroglycan to the cell-BM interface, weakening BM adhesion, and allowing the BM to move on the uterine side of the BM. Lower panel left (P6.p 8-cell stage): the vulvF cells divide and further invaginate with the vulvE cells. The BM stops shifting over the nondividing vulvD cell, which sets the width of the opening of the BM gap. Integrin and VAB-19 (KANK) localize to the vulvD-BM interface to stabilize BM adhesion. Lower panel right: dystroglycan levels continue to be reduced at the interface, allowing the BM to slide to a position determined by the underlying vulvD cell.

invade with the vulvE cells. The BM stops shifting over the nondividing vulvD cell, which sets the width of the opening of the BM gap. Integrin and VAB-19 (KANK) localize to the vulvD-BM interface to stabilize BM adhesion. Lower panel right: dystroglycan levels continue to be reduced at the interface, allowing the BM to slide to a position determined by the underlying vulvD cell.

Unfortunately not effective for reasons that remain unclear (Zucker *et al.* 2000; Coussens *et al.* 2002; Dufour and Overall 2013), but could be due to changes in cancer cell invasion strategy (Te Boekhorst and Friedl 2016). There are over 20 encoded MMPs in vertebrate genomes, making genetic assessment of their necessity for invasion unfeasible (Rowe and Weiss 2008). In comparison, only six MMPs are encoded in the *C. elegans* genome (Altincicek *et al.* 2010), and it should be possible to determine if AC invasion can occur in their absence. Given the finding that the AC invades, in part, by displacing BM, it is possible that invasive cells may be capable of invading through solely physical means. An important future direction that should be addressed using the AC invasion model is to determine the necessity of MMPs during BM invasion.

It is also unclear how the invasive protrusion rapidly expands and generates forces to displace the encircling BM and vulval tissue. Actin-binding proteins are known to be important for cellular force production (Blanchoin *et al.* 2014), but it is unclear whether actin polymerization drives the expansion of the invasive protrusion in the AC, or what actin-binding proteins are involved in protrusion stabilization. The regulation of membrane dynamics during cell migration and invasion is also poorly understood (Lecuit and Pilot 2003; Fletcher and Rappoport 2010; Hastie and Sherwood 2016), and it is not known if invasive cells expand or alter their plasma membranes to breach BM barriers.

The highly stereotyped nature of AC invasion should allow for the study of AC membrane addition and how it is regulated. Given the rapid enlargement of the protrusion, it will also be interesting to explore if aquaporins and ion channels play a role by generating hydrostatic forces that displace the BM (Schwab and Stock 2014).

It is unknown how the AC transitions rapidly from AC-BM adhesion prior to invasion to AC-vulval precursor cell (cell-cell) adhesion following invasion. Despite extensive screening, we have not found an adhesion system that mediates this connection (D. R. Sherwood, unpublished data). Interestingly, recent work has shown that when the AC adheres to the vulval precursor cells it stimulates the recruitment of the F-BAR-domain protein TOCA-1 to the vulval precursor cell-AC interface. TOCA-1 concentrates nonmuscle myosin NMY-2 at this cell surface and reorients contractile forces, which constricts the lateral membrane of the vulval precursor cells thus reshaping them (Yang *et al.* 2017). These observations indicate that invasive cells can also induce cell shape changes, which could possibly facilitate tissue invasion.

Finally, single-cell isolation procedures involving cell dissociation and fluorescence activated cell sorting (FACS) have been developed in *C. elegans* to determine the expression profile of individual cells (Spencer *et al.* 2014). Applied to the AC, this approach could reveal the first complete expression signature

of an actively invading cell. By combining AC RNAseq studies with transcription factor mutants, it will be possible to carry out a comprehensive analysis of the transcriptional networks that direct invasive behavior, thus revealing how cells are programmed to invade.

Distal Tip Cell Migration: a Leader Cell Migration that Shapes a Tissue

The distal tip cells (DTCs) are a pair of somatic gonad cells born during the L1 stage that act as leader cells in hermaphrodites, tracing out the symmetrical double lobed gonad through their migration trajectory (Kimble and Hirsh 1979; Kimble and White 1981). The ease of identifying defects in gonad shape with a dissecting microscope has led to the isolation of numerous genes regulating DTC migration through forward genetic (many termed *mig*, for Migration defective) as well as RNAi, screens (Hedgecock *et al.* 1987, 1990; Nishiwaki 1999; Cram *et al.* 2006; Suzuki *et al.* 2006). A similar leader cell function is mediated by the linker cell in the male *C. elegans* gonad, which establishes a single lobed J-shaped gonad (Kimble and Hirsh 1979). The migration of the linker cell has not been studied in as much detail as the DTC, but appears to be regulated by many, but not all, of the same genes controlling DTC movement (Kato and Sternberg 2009).

The DTCs initiate migration during the L2 stage, and continue moving at a variable rate of ~6–10 $\mu\text{m/hr}$ until the early adult stage, a duration of ~25 hr (Lints and Hall 2017). The two DTCs, one in each gonad arm, are located at the anterior and posterior ends of the gonad, and their mirror imaged migration has been classified into three phases (Figure 4 and Figure 5A) (Hedgecock *et al.* 1987; Nishiwaki 1999): (phase 1) During the L2 and early-to-mid L3 stage, the DTCs move ventrally away from each other along the ventral body wall muscle BM toward the anterior (head) and posterior (tail) of the animal; (phase 2) during the late-L3 stage, both DTCs turn 90° and move from the ventral to dorsal surface, migrating along the BM that covers the lateral epidermis; (phase 3) during the L4 stage, the DTCs turn 90° and move back to the midsection along the BM covering the dorsal body wall muscles. Together, these different phases of migration build a gonad with two equivalent U shaped arms (Figure 4 and Figure 5A).

The DTC is shaped as a smooth cap that extends over the most distal three-to-five germ cells during its migration (Hall *et al.* 1999; Lints and Hall 2017). Five pairs of sheath cells encase the remaining germ cells and follow the path of the DTCs (Figure 4) (Killian and Hubbard 2005). The entire growing gonad, including the DTCs and sheath cells, are surrounded by a BM (Hall *et al.* 1999; C. C. Huang *et al.* 2003). The germ cells are thought to follow the DTC through extensive cell divisions, expanding from two cells at the beginning of larval development to ~1000 at the young adult stage. The proliferation of the germ cells might help extend the distal gonad arms (Kimble and White 1981; Killian and

Hubbard 2005). Although not quantified, loss of germline proliferation has not been reported to alter the DTC migration path (and thus the shape of the gonad), but the distance of DTC movement during the third phase of migration appears to be reduced (Austin and Kimble 1987). It is unclear how the sheath cells follow the DTC. They are thought to be either pulled along by the dividing germ cells or to actively migrate over the germ cells (Lints and Hall 2017). Interestingly, the DTC extends a single forward directed protrusion only during the initiation of the dorsal turn that begins the second phase of migration, and, otherwise, does not appear protrusive during its migration (Kim *et al.* 2011). In many respects, DTC migration has similarities to collective cell migration events in vertebrates that build branching tubular organs (Andrew and Ewald 2010). Very little is known about how leader cells in morphogenetic branching events migrate while enwrapped in tough, dense sheets of BM (Friedl and Wolf 2010). Such branching morphogenetic programs are thought to be misregulated in some epithelial tumors, allowing groups of cancer cells to collectively invade neighboring tissues (Friedl and Gilmour 2009; Gray *et al.* 2010). Thus, understanding DTC migration has significance to mechanisms underlying organogenesis and potentially some invasive cancers.

DTC modification of and interaction with BM during migration

The DTC expresses many BM components during its migration, including type IV collagen (*C. elegans* *emb-9*/collagen IV $\alpha 1$ and *let-2*/collagen IV $\alpha 2$), laminin (*epi-1*/laminin αB and *lam-1*/laminin β), nidogen (*nid-1*), and agrin (*agr-1*) (Graham *et al.* 1997; Kim and Wadsworth 2000; C. Huang *et al.* 2003; Hrus *et al.* 2007; Clay and Sherwood 2015), suggesting that the DTC secretes, deposits, and remodels the BM that surrounds it during its migration. Reduction of laminin and type IV collagen severely hampers DTC migration, indicating the important role of BM in DTC movement (C. Huang *et al.* 2003; Kao *et al.* 2006; Kubota *et al.* 2008; Kawano *et al.* 2009; Wong and Schwarzbauer 2012).

Throughout its migration the DTC expresses and secretes **GON-1**, an ortholog to the vertebrate proteases ADAMTS-9 and ADAMTS-20 (A Disintegrin And Metalloprotease with ThromboSpondin repeats protein) (Blelloch and Kimble 1999; Llamazares *et al.* 2003; Somerville *et al.* 2003). **GON-1** is essential to facilitate DTC migration (Figure 5B) (Blelloch *et al.* 1999). In animals harboring null alleles of the *gon-1* gene, or rescue constructs lacking the metalloprotease catalytic domain, the DTC fails to migrate (Figure 6, A and B) (Blelloch *et al.* 1999; Blelloch and Kimble 1999). The activity of **GON-1** may be controlled by **MIG-6L**, a DTC expressed isoform of the conserved extracellular matrix protein papilin (Kawano *et al.* 2009). Genetic reduction of *mig-6(l)* gene activity leads to a similar defect in DTC migration as loss of the ADAMTS *gon-1* (Cram *et al.* 2006; Kawano *et al.* 2009). In *Drosophila*, the ortholog of the **MIG-6** protein, papilin, binds to and regulates the activity of a *Drosophila* ADAMTS collagenase (Kramerova *et al.* 2000). Genetic studies indicate that type IV collagen and




Figure 4 DTC migration, a leading cell that shapes an organ. The pair of DTCs (one on the anterior the other on the posterior arm of the basement membrane enwrapped gonad) initiate migration at the L2 larval stage. During the L2 and L3 larval stages (phase 1 of migration), the DTCs move ventrally away from each other along the BMs of the ventral body wall muscles (data not shown) toward the anterior (head) and posterior (tail) of the animal. In the rest of the figure, the posterior gonad arm (right side) shows the germ cells from the L3 stage onward, while the anterior arm (left side) shows the basement membrane and five pairs of sheath cells (sh1-5) that cover the germ cells. The sheath cells follow the path of the DTCs. The digestive tube is shown in light gray and the anterior gonad arm passes underneath it to the other side of the animal. During the late-L3 stage, both DTCs turn 90° and move from the ventral to dorsal surface (phase 2 of migration), moving along the BM of the lateral epidermis. At the early L4 stage, the DTCs turn 90° and move back to the midsection along the BM of dorsal body wall muscles during the L4 stage (phase 3 of migration). DTC migration ceases in the early adult.

the matrix protein fibulin-1 (*C. elegans* *FBL-1*) oppose *GON-1* function (Hesselson *et al.* 2004; Kubota *et al.* 2012). As fibulin-1 maintains collagen in the BM (Kubota *et al.* 2012), an attractive model is that the ADAMTS *GON-1* cleaves collagen to facilitate DTC movement and BM expansion during DTC migration (Figure 5B).

Another ADAMTS protein, *MIG-17*, also regulates DTC migration (Nishiwaki *et al.* 2000). *MIG-17* is secreted from the ventral and dorsal body wall muscle cells and accumulates in the gonadal BM during the L3 stage, where it regulates the directional migration of the DTC (Figure 5B) (Nishiwaki *et al.* 2000). In the absence of *MIG-17*, DTC migration occurs, but the DTC meanders and the gonad has an abnormal shape (Nishiwaki 1999). BM localization and the catalytic domain of *MIG-17* are required for its function, suggesting that *MIG-17* may cleave BM targets (Ihara and Nishiwaki 2007). *MIG-17* recruits and activates (perhaps by proteolysis) an isoform of fibulin (fibulin-1C), which then recruits the BM protein nidogen to the gonad (Kubota *et al.* 2004, 2008). Based on genetic interactions, *MIG-17* has also been proposed to modify type IV collagen to promote directional migration (Kubota *et al.* 2008). These observations imply that dynamic interactions of matrix proteins and proteases continually remodel the BM covering the DTC in a manner that both allows DTC migration and helps direct its path. ADAMTS proteins have complex functions in vertebrate tissue formation and maintenance, and mutations or misregulation of ADAMTS proteins are associated with numerous diseases, including cancer and arthritis (Cal and Lopez-Otin 2015; Kelwick *et al.* 2015). Understanding the functions of these proteases during DTC migration will likely provide insight into their roles in vertebrate tissue formation, and clues as to how their misregulation contributes to human diseases.

Several transmembrane proteins that act as BM receptors, or are strongly associated with BM, regulate DTC migration. These include the *C. elegans* orthologs of the proteins teneurin (*C. elegans* *TEN-1*), dystroglycan (*DGN-1*), and integrin (Baum and Garriga 1997; Lee *et al.* 2001; Drabikowski *et al.* 2005; Trzebiatowska *et al.* 2008; Topf and Chiquet-Ehrismann 2011). Of these, only the function of integrin has been carefully characterized during DTC migration. Both *C. elegans* integrins, α *INA-1*/ β *PAT-3* and α *PAT-2*/ β *PAT-3*, are expressed in the DTC and regulate its migration (Baum and Garriga 1997; Lee *et al.* 2001; Meighan and Schwarzbauer 2007). The α -integrin *INA-1* is expressed in the DTC prior to, and during, its migration, and promotes DTC motility, but does not appear to play a major role in DTC pathfinding (*i.e.*, the direction of movement) (Baum and Garriga 1997). *INA-1* expression is downregulated when DTC migration ceases and failure to turn off *ina-1* gene expression results in DTCs that continue to migrate in adults (Meighan and Schwarzbauer 2007). Termination of migration is also regulated by the E3 ubiquitin ligases *RNF-121* and *RNF-5*, which can target *PAT-3* and a protein with similarity to the integrin effector paxillin (*UNC-95*) for degradation (Broday *et al.* 2004; Darom *et al.* 2010; Kovacevic *et al.* 2012). However, the specific targets of *RNF-121* and *RNF-5* that promote cessation of DTC migration




Figure 5 DTC migration timing, BM interactions, polarization, and gene regulation. Only the posterior gonad arm is shown. (A) The timing of DTC migration shown is at 20°. (B–D) Details of proteins and interactions that regulate DTC migration are described in the text and outlined here for a global view. Note, the “Wnt –” shown in (C) represents a hypothetical possible function for Wnt in inhibiting polarization along the anterior-posterior axis during phase 2 of DTC migration.

are not yet clear. *CACN-1*, a conserved protein and component of the spliceosome, is also required to halt DTC migration (Figure 5B and Figure 6E). Genetic studies suggest *CACN-1* inhibits the Rac GTPase *MIG-2* (Tannoury *et al.* 2010; Doherty *et al.* 2014), although the precise mechanism of *MIG-2* down-regulation is not known.

The α -integrin *PAT-2* is first expressed in the DTC during phase 2 of migration after the DTC turns dorsally, and is required for correct pathfinding along the dorsal body wall muscle BM surface during phase 3 of DTC migration. Knock-down of key components of the retrograde vesicular trafficking pathway has suggested that the polarized trafficking of integrin to the leading edge of the DTC is important for DTC pathfinding (Shafaq-Zadah *et al.* 2016); however, the effects of these knockdowns on the localization of the pathfinding integrin *PAT-2* have not been reported. The requirement for both *INA-1/PAT-3* and *PAT-2/PAT-3* integrin heterodimers during phase 3 of DTC migration suggests that each simultaneously promotes distinct activities within the DTC, either via separate extracellular ligands or through different intracellular signaling partners (Meighan and Schwarzbauer 2007). Interestingly, there appears to be feedback between the amount of *PAT-3* (β -integrin) in the DTC and the levels of type IV collagen in the BM, as lower concentrations of collagen in the BM decrease the levels of a *PAT-3::GFP* translational reporter (Kubota *et al.* 2012). These observations suggest that dynamic interactions between the composition of the BM and the receptors that bind to it regulate DTC migration.

Many known integrin downstream effectors mediate DTC migration and may function downstream of *INA-1/PAT-3* and *PAT-2/PAT-3*. These include *C. elegans* orthologs of talin, kindlin, ILK, the tyrosine-protein kinase Src, the Rho GTPases Rac (*C. elegans* *MIG-2* and *CED-10*), Cdc42, associated Rho GTPase regulators and effectors such as the nucleoside-diphosphate kinase *NDK-1*, and a GIT/PIX/PAK signaling pathway (Reddien and Horvitz 2000; Lundquist *et al.* 2001; Cram *et al.* 2003; Itoh *et al.* 2005; Xu *et al.* 2006; Meighan and Schwarzbauer 2007; Lucanic and Cheng 2008; Wong and Schwarzbauer 2012; Fancsalszky *et al.* 2014). Loss of most integrin effectors perturbs both DTC motility and pathfinding; however, some effectors predominantly alter motility (e.g., *NDK-1*; Figure 6C), while others pathfinding (e.g., *SRC-1*, Figure 6D) (Itoh *et al.* 2005; Fancsalszky *et al.* 2014). Thus, the *INA-1* and *PAT-2* integrins might achieve at least some of their respective roles in DTC motility and pathfinding through engagement with distinct effectors. Despite the characterization of many effectors, it remains unclear if integrin activity promotes DTC motility and directional migration through known roles in polarization, vesicular trafficking, BM deposition, adhesion strength, or cytoskeletal dynamics (Bokel and Brown 2002; Harburger and Calderwood 2009; Vicente-Manzanares *et al.* 2009; Huttenlocher and Horvitz 2011; Wickstrom and Fassler 2011; Yurchenco 2011) or if *INA-1* and *PAT-2* integrins act through novel mechanisms.

Stabilizing the DTC path: the matrix proteins *HIM-4* (hemocentin) and *DIG-1*

Study of DTC migration and gonad formation has also revealed possible mechanisms that stabilize tissue positioning—a poorly understood aspect of morphogenesis. The DTC and gonad are encased in BM, and migrate along the BMs of the




Figure 6 Examples of DTC migration defects. Only the posterior gonad arm is shown. (A) Wild type migration, (B) a *gon-1* mutant where no DTC migration occurs, (C) an *ndk-1* mutant where DTC migration is incomplete, (D) a *src-1* mutant and an *unc-5* mutant where DTC migration shows pathfinding defects, (E) reduction of *cacn-1* by RNAi leads to extended DTC migration (a cessation of migration defect).

body wall muscle and epidermis, thus forming a BM–BM interface where the gonad must likely adhere to maintain its position within the body cavity. A potential molecule that stabilizes gonad positioning is *HIM-4* (hemicentin), a large (>5000 amino acids) extracellular matrix protein of the immunoglobulin superfamily that is expressed by the DTC from the L1 stage and throughout its migration (Vogel and Hedgecock 2001). *HIM-4* localizes between BMs of neighboring tissues in *C. elegans* and vertebrates (Vogel and Hedgecock 2001; Xu *et al.* 2007), and has recently been shown to connect neighboring BMs (Morrissey *et al.* 2014). Consistent with a possible role in stabilizing the position of the gonad along the neighboring BMs, loss of *HIM-4* (hemicentin) can result in the ventral gonad detaching from the underlying body wall muscle BM (Vogel and Hedgecock 2001). Another secreted matrix associated molecule, *DIG-1*, which is a giant member of the immunoglobulin superfamily (>13,000 amino acids), might also mediate BM–BM interactions (Benard *et al.* 2006; Burket *et al.* 2006). In *dig-1* mutants the entire gonad is often displaced from its normal position (most often relocated anteriorly). Strongly indicative of a possible role in BM–BM adhesion, the gonads of *dig-1* mutants can be shifted within the body cavity by mechanical manipulation (Thomas *et al.* 1990; Burket *et al.* 2006). Reporter constructs indicate that the *dig-1* gene is expressed in many muscles surrounding the gonad, but its expression has not been reported in the DTC (Benard *et al.* 2006).

Diffusible cues help orient DTC migration: netrin and Wnts

Studies on DTC migration have helped reveal a coordinated dorso-ventral (D/V) and anterior-posterior (A/P) positioning

system within the worm involving the diffusible netrin and Wnt cues (Figure 5C). These navigation signals orient numerous migratory and cell outgrowth behaviors in vertebrates and invertebrates (Silhankova and Korswagen 2007; Lai Wing Sun *et al.* 2011; Hikasa and Sokol 2013), suggesting the navigational system revealed in *C. elegans* is conserved. Studies in worms have been simplified by the smaller gene families that encode these pathways. For example, vertebrates harbor five genes for netrin ligands, whereas *C. elegans* has only the *UNC-6* protein (Lai Wing Sun *et al.* 2011). *C. elegans* netrin (*UNC-6*) is secreted from ventral cells and serves as an attractive source for cells expressing the transmembrane receptor *UNC-40* (vertebrate DCC). *UNC-6* (netrin) can also function as a repulsive cue for cells expressing the netrin receptor *UNC-5* either alone or with *UNC-40* (DCC) (Hedgecock *et al.* 1990; Ishii *et al.* 1992; Leung-Hagesteijn *et al.* 1992; Chan *et al.* 1996). *C. elegans* encodes four Wnt frizzled receptors (*CFZ-2*, *LIN-17*, *MIG-1*, and *MOM-5*), and five Wnt ligands (*CWN-1*, *CWN-2*, *EGL-20*, *LIN-44*, and *MOM-5*) (Eisenmann 2005), simplifying analysis as compared to mammalian genomes, which encode 19 Wnt proteins (Clevers and Nusse 2012; Sawa and Korswagen 2013). The *C. elegans* Wnt ligands have distinct graded distributions along the A/P axis, which can polarize cells (Whangbo and Kenyon 1999; Goldstein *et al.* 2006; Hilliard and Bargmann 2006; Pan *et al.* 2006; Levy-Strumpf 2016), thus providing a system for cells to assess positional information along the A/P axis.

Netrin and Wnts act as key directional signals for DTC migration along the body axes. For example, expression of the netrin receptor *UNC-5* is activated in the DTC just prior to phase 2, and its expression helps initiate the 90° DTC dorsal

turn and migration away from ventral **UNC-6** (Figure 5C) (Su *et al.* 2000). In the absence of **UNC-5**, **UNC-6** (netrin), or **UNC-40** (DCC), both the anterior and posterior DTCs often fail to turn dorsally, and instead continue to migrate along the ventral body wall muscle (Figure 6D (Hedgecock *et al.* 1990)). The levels of the glycosaminoglycan chondroitin may regulate netrin signaling, as chondroitin synthase (**SQV-5**) and its cofactor chondroitin polymerizing factor (**MIG-22**) promote **UNC-5** mediated dorsal DTC migration (Suzuki *et al.* 2006). In contrast to loss of netrin, mutations in Wnt pathway components lead to A/P migration defects during the third phase of DTC migration (Cabello *et al.* 2010; Levy-Strumpf and Culotti 2014). Genetic studies suggest that anterior and posterior localized Wnts (anterior **CWN-1** and **LIN-44** and posterior **EGL-20**) oppose each other's activity to provide a precise positioning system to orient the posterior DTC along the A/P axis (Figure 5C). Similarly, the anterior expressed Wnt **CWN-2** and Wnt inhibitor **SFRP-1** and the posteriorly expressed Wnts **LIN-44** and **CWN-1** help direct A/P axis polarization of the anterior DTC during the third phase of migration (Levy-Strumpf and Culotti 2014). Whether Wnts act to polarize phase 1 of migration is unclear. Cross talk between the netrin and Wnt pathways is also important. The Wnt receptor **MOM-5** is upregulated in both DTCs at the end of phase 2, and downregulates the **UNC-5** receptor to ensure proper polarized migration of both the DTCs in phase 3 (Figure 5C) (Levy-Strumpf *et al.* 2015).

Notably, netrin and Wnt pathway mutants show only partially penetrant DTC polarity defects, suggesting that each cue acts with other signal(s) to direct DTC migration. Surprisingly, simultaneous impairment of both the netrin and Wnt pathways reveals that netrin and Wnt act redundantly (*i.e.*, they function together) to polarize DTC migration (Levy-Strumpf and Culotti 2014). Thus, each cue provides polarity information along both axes (Levy-Strumpf and Culotti 2014). How these cues define the axis orthogonal to their graded distribution is not well understood, nor is it clear how they control polarization of both the anterior and the posterior DTCs such that they move in opposite directions during phase 3. There is evidence that Wnt ligands can exclude the polarization machinery along the A/P axis in the HSN neuron during axon outgrowth, thus helping direct polarity along its D/V axis (Kulkarni *et al.* 2013; Levy-Strumpf 2016). Wnt might act similarly during the dorsal migration of the DTC (Figure 5C). Further, opposing activities of Wnts, which are in graded distributions along the A/P axis, may help precisely position both anterior and posterior DTC migration along this axis (Levy-Strumpf and Culotti 2014). As integrin-BM interactions also impact polarity, and integrin, netrin, and Wnts share several downstream effectors, including Rac (**CED-10**) and Src kinase (**SRC-1**) (Gitai *et al.* 2003; Itoh *et al.* 2005; Meighan and Schwarzbauer 2007; Harburger and Calderwood 2009; Cabello *et al.* 2010; Lai Wing Sun *et al.* 2011; Levy-Strumpf and Culotti 2014; Wang *et al.* 2014a; Levy-Strumpf 2016), extensive collaboration and cross talk between Wnt, netrin, and integrin signaling likely exists during DTC polarization and migration.

Polarity of the nucleus and cytoskeleton during DTC migration

The nucleus is positioned at the leading front of the DTC throughout its migration. The **VAB-10B1** protein, the *C. elegans* ortholog of the vertebrate cytoskeleton cross-linker spectraplak, is required to move the nucleus to the leading edge during the dorsal turn of the DTC (phase 2 of migration) (Kim *et al.* 2011; Suozzi *et al.* 2012). The **VAB-10B1** protein contains both F-actin and microtubule binding motifs, and both modules are required for nuclear translocation during the dorsal turn (Kim *et al.* 2011). Examination of microtubules and F-actin revealed that F-actin is present in filamentous structures that are loosely aligned along the axis of DTC migration, while microtubules are enriched at the trailing side of the DTC. Visualization of microtubule growth using a plus-end tracking protein indicated that microtubules grow dynamically toward the nucleus and leading edge during DTC migration, whereas microtubule growth in the rear of the DTC appears random. Notably, **VAB-10B1** is largely dispensable for formation of polarized F-actin filaments, but is required for organization of the microtubule network within the DTC and polarized microtubule growth toward the nucleus and leading edge. Surprisingly, mutant analysis showed that **VAB-10B1** is not required for DTC pathfinding, but is necessary for DTC migration: DTCs in **vab-10** mutants migrate more slowly, and the animals have shortened gonad arms (Kim *et al.* 2011). These findings imply that **VAB-10B1** (spectraplak) and likely polarized microtubule dynamics are components of the engine that drives DTC migration, but they are not involved in the mechanism that orients DTC movement. Instead, the orientation of F-actin fibers along the axis of migration, which is not severely affected by loss of **VAB-10B1**, might be a component of the DTC orientation mechanism. It will be interesting to further explore if **VAB-10B1** (spectraplak) links mechanisms that orient DTC migration (Wnt, netrin, integrin regulation of F-actin polarity) to those that drive migration (microtubule polarity and polarized dynamic growth). Both *C. elegans* septin proteins, **UNC-59** and **UNC-61**, are also required for robust DTC movement and pathfinding, although the specific role(s) this cytoskeletal system plays in DTC migration remains to be explored (Finger *et al.* 2003).

Transcriptional regulation of DTC migration

Transcriptional programming of the DTC plays a crucial role in all aspects of its migration (Figure 5D). As with the AC, an early regulator of the DTC transcriptional program is the basic helix-loop-helix (bHLH) transcription factor **HLH-2** (vertebrate E proteins) (Karp and Greenwald 2004; Chesney *et al.* 2009). **HLH-2** is upregulated in the DTC and controls the expression of key genes that help initiate DTC migration, including genes encoding **GON-1** (ADAMTS), **MIG-6** (papilin), and **INA-1** (α -integrin) (Krause *et al.* 1997; Karp and Greenwald 2004; Cram *et al.* 2006; Tamai and Nishiwaki 2007; Meighan *et al.* 2015). bHLH factors function as heterodimers, and **HLH-2** physically interacts with the Achaete-Scute bHLH transcription factor family member **HLH-12** to control target gene expression in the DTC (Figure

5D). Importantly, *gon-1* and *ina-1* genes are still expressed, albeit weakly, in the DTC after loss of *hlh-2* and *hlh-12* (Tamai and Nishiwaki 2007; Meighan *et al.* 2015), suggesting that other transcription factors are also involved in the initiation of DTC migration. The role of *HLH-2* in migration of the DTC, invasion of the *C. elegans* AC (Schindler and Sherwood 2011), and the role of the vertebrate *HLH-2* ortholog E2A in epithelial-to-mesenchymal transition in mammalian cells (Sobrado *et al.* 2009), suggests a conserved role for *HLH-2/E* proteins in promoting cell migration and invasion.

The dorsal turn of the DTC depends on the *UNC-5* protein, which is upregulated by a conserved circuit of heterochronic genes that controls developmental timing events in the worm (Figure 5D) (Su *et al.* 2000). This gene circuit involves the Blimp-1/PRDI-BF1 zinc finger transcriptional repressor *BLMP-1*, the zinc finger EGR (early growth response) family protein *LIN-29*, the steroid hormone receptor *DAF-12*, and the F-Box protein *DRE-1*, which is the key recognition subunit of the SCF ubiquitin ligase complex (Rougvie and Ambros 1995; Antebi *et al.* 1998, 2000; Su *et al.* 2000; Fielenbach *et al.* 2007; Huang *et al.* 2014). *BLMP-1* is a transcriptional repressor that is expressed in the DTCs during the first phase of DTC migration. *BLMP-1* binds to the upstream regulatory region of the *unc-5* gene and is thought to repress *unc-5* transcription. The repression of *unc-5* is relieved at the beginning of the L3 stage by the global developmental timing cues of dafachronic acid release and lower levels of *LIN-42* (Period protein), which help activate *DAF-12* and *LIN-29*, respectively (Huang *et al.* 2014; Cecchetelli and Cram 2017). The levels of *DRE-1* are also upregulated at the L3 stage (Fielenbach *et al.* 2007). *LIN-29* and *DAF-12* repress the transcription of the *blmp-1* gene, while *DRE-1* targets the *BLMP-1* protein for degradation. Together, these activities remove *BLMP-1* from the DTC and relieve the repression on *unc-5* expression (Figure 5D). Interestingly, prior to its removal from the DTC, *BLMP-1* also represses the expression of *lin-29*, suggesting that a double negative feedback loop between *LIN-29* and *BLMP-1* might act as a robust switch in the decision to turn on the expression of *unc-5* (Huang *et al.* 2014). In addition, *LIN-29* and *DAF-12* act cooperatively to promote *unc-5* expression (Huang *et al.* 2014). A consensus *DAF-12* binding sequence in the *unc-5* promoter suggests that *DAF-12* may directly activate *unc-5* transcription (Huang *et al.* 2014). Expression of *unc-5* in the DTC triggers the dorsal turn, orienting the AC away from ventral sources of *UNC-6* protein (Figure 5C) (Hedgecock *et al.* 1990; Su *et al.* 2000). Precocious expression of the *unc-5* gene through ectopic expression or loss of *blmp-1* function results in early DTCs dorsal turns, indicating the importance of the correct timing of *unc-5* gene expression (Su *et al.* 2000; Huang *et al.* 2014). The vertebrate ortholog of *DRE-1*, Fbxo11, mediates the degradation of the pro-oncogene BLC6, which has sequence homology to *BLMP-1*. Overexpression of BLC6 is implicated in the pathogenesis of human B-cell lymphomas (Ci *et al.* 2008). Fbxo11 is deleted in diffuse large B-cell lymphomas, and, as a consequence, BLC6

expression is upregulated (Duan *et al.* 2012). BLC6 has recently been implicated in other cancers, including leukemia, breast cancer, and nonsmall-cell lung cancer (Cardenas *et al.* 2017). It will be interesting to determine if the *DRE-1/BLMP-1* regulatory circuit that controls DTC migration might be a conserved switch that is misregulated in numerous cancers.

Following the dorsal turn, other transcription factors help guide DTC migration (Figure 5D). During the dorsal migration of the DTC (phase 2), *VAB-3*, a Pax6 transcription factor ortholog, turns on the expression of the α -integrin subunit gene encoding *PAT-2*, which regulates pathfinding during phase 3 of DTC movement (Meighan and Schwarzbauer 2007). In addition, *VAB-3* downregulates the expression of the gene encoding *INA-1* (integrin α -subunit), which is necessary to cease DTC migration in the early adult stage (Meighan and Schwarzbauer 2007). Several other transcriptional regulators also control pathfinding during phase 3. These include a CBP/p300 transcriptional coactivator *CBP-1*, and a CREBH transcription factor *LET-607*, which are both expressed in the DTC, and whose DTC-specific loss results in defects in the second turn of the DTC back toward the mid-section (Wong *et al.* 2014). Functional DTC transcriptional targets of *CBP-1* and *LET-607* associated with integrin activity have been identified, including genes encoding *SRC-1* (SRC kinase), *TLN-1* (talin), *NMY-2* (nonmuscle myosin heavy chain II), and *PAT-2* (α -integrin subunit). Regulation of *pat-2* gene expression is likely indirect, as *CBP-1* and *LET-607* promote *vab-3* expression. Notably, expression of the gene encoding the matrix protein *MIG-6* (papilin), and the bHLH transcription factor *MIG-34*, which promote the initiation of DTC migration, are not regulated by *CBP-1* and *CREBH* (Wong *et al.* 2014). Together, these observations indicate that distinct sets of transcription factors and transcriptional regulators help direct the different steps of the DTC migration program. Thus, a combination of a dynamic transcriptional program within the DTC, and a complex extracellular environment of BM components, proteases, and diffusible signals (Wnt, netrin) directs and guides the specific path of DTC migration.

microRNAs confer robustness to DTC migration under temperature stress

Because of its stereotyped movement and ease of visual analysis, DTC migration can serve as a model to examine how developmental migration programs are buffered against environmental stresses. A pair of conserved microRNAs, *mir-34* and *mir-83* (orthologs of mammalian *mir-34* and *mir-29*, respectively) act to ensure robust DTC migration, and appear to be particularly effective at maintaining the correct execution of DTC migration in the face of temperature stress (Figure 5D) (Burke *et al.* 2015). Loss of both *mir-34* and *mir-83* microRNAs results in a mild DTC migration defect, specifically affecting phase 1 and phase 3 of migration. Evidence suggests that both microRNAs function in the DTC and together directly suppress translation of the mRNA encoding the β -integrin subunit *PAT-3* and one of its downstream

effectors, the Rho GTPase **CDC-42** (Figure 5D) (Burke *et al.* 2015). Temperature oscillations between 15 and 25° within a 2-hr window at the time of DTC birth in the L1 larva dramatically enhance the DTC migration defect of animals harboring mutations in **mir-34** and **mir-83**. This suggests, somewhat perplexingly, that **mir-34** and **mir-83** may act at the birth of the DTC cells in the L1 stage, many hours (~10–30) before DTC migration occurs in the L2, L3, and L4 stages. Alternatively, these microRNAs might function at the time of DTC migration to buffer gene expression changes set in motion by the earlier temperature oscillations. Misregulation of the vertebrate orthologs of **mir-34** and **mir-83** are associated with many cancers (Hermeking 2010; Jiang *et al.* 2014; Yan *et al.* 2015), and they also appear to function together to coregulate gene regulatory networks, such as a p53 network that promotes apoptosis (Burke *et al.* 2015). Thus, understanding how the DTC executes accurate migration in the face of stressful environmental conditions will reveal important mechanisms that maintain the fidelity and flexibility of gene regulatory networks in normal development, and the ways in which these networks go awry in human diseases.

DTC migration: key unanswered questions

An important unanswered question in DTC migration relates to how the DTC actually moves. The DTC migrates encased within a BM, a mode of cell migration that is widespread during branching morphogenesis, but poorly understood (Friedl and Wolf 2010). Germline proliferation may help propel DTC movement; however, the precise contribution of the dividing germ cells has not been determined. As mutants such as **vab-3** lead to perpetual DTC migration in the absence of apparent germline hyperproliferation (Meighan and Schwarzbauer 2007), it strongly suggests that germline proliferation is not an essential driving force. Further, the male linker cell, which is functionally analogous to the DTC, can migrate when detached from the proliferating germline (Kato *et al.* 2014). Thus, it seems likely the DTC can move independently of germline proliferation.

It is not yet clear if the DTC employs the two primary modes of cell migration characterized to date—mesenchymal and bleb-based (Charras and Paluch 2008; Lammermann and Sixt 2009; Petrie and Yamada 2012, 2016; Te Boekhorst *et al.* 2016). During mesenchymal migration, cells extend protrusions through controlled F-actin formation. These protrusions adhere to the cell substrate, while the back end of the cell deadheres and retracts through actomyosin contractility, thus moving the cell or groups of cells forward (Friedl and Gilmour 2009). During bleb-based migration, cells use actomyosin contractility to create rounded membrane protrusions that direct cell migration by wedging through and extending between spaces within the neighboring microenvironment (Friedl and Wolf 2010; Paluch and Raz 2013). Live cell imaging has indicated that the DTC only produces a single large protrusion during the dorsal turn at the initiation of the second migration phase, and this protrusion is not enriched with F-actin (Kim *et al.* 2011). Blebbing has also not

been observed in the DTC during its movement (Kim *et al.* 2011). However, loss of **GEX-3** a component of the WAVE complex that activates the actin nucleator the Arp2/3 complex (Soto *et al.* 2002; Shakir *et al.* 2008), and loss of components of the myosin machinery, cause mild DTC migration defects, consistent with roles in movement (Nishiwaki 1999; Cram *et al.* 2006; Wong *et al.* 2014). Notably, F-actin has only been observed via fusion of GFP to the actin binding domain from Moesin (Kim *et al.* 2011). As different populations of F-actin are bound by distinct F-actin probes (Washington and Knecht 2008), it will be important to examine F-actin using different actin binding probes that might label forms of F-actin that drive movement.

It is possible that the DTC uses a less well-established or novel mechanism to move. Given the importance of nuclear positioning in driving DTC movement, the role of the nucleus, which helps power movement of human fibroblasts and fibrosarcoma cells by acting as a piston to generate forward acting forces (Petrie *et al.* 2014, 2017), could be a contributing factor in DTC migration. In addition, the physically confining environment of the encasing BM might allow polarized water permeation or retrograde flow of actomyosin under the plasma membrane (a migration mode termed “chimneying”) to drive DTC movement, as it does for cancer cells in confining *in vitro* environments (Paluch and Raz 2013; Stroka *et al.* 2014; Bergert *et al.* 2015). The crucial role of integrins, secreted matrix proteases, and BM proteins in DTC migration further suggests the intriguing possibility that secretion or polarized assembly of BM that the DTC generates might help power DTC migration. Indeed, there is evidence that some bacteria power their movement via polysaccharide secretion (Jarrell and McBride 2008; Khayatan *et al.* 2015). It seems likely that the DTC might use multiple ways to propel movement and teasing out these mechanisms are important areas of future study.

Other unanswered questions center around DTC polarization. For example, how A/P localized Wnts oppose each other’s functions to precisely guide A/P migration is unclear. Further, how **UNC-5** directs polarity away from **UNC-6** (netrin) sources is poorly understood. It is also unclear what polarizes migration in the DTC. The polarized orientation of F-actin fibers within the DTC in the direction of migration suggests that these fibers might be associated with DTC polarity, and thus could help orient the mechanism(s) that generate movement. The convergence of polarizing signals from integrin, netrin, and Wnt pathways on small Rho GTPases strongly suggest their involvement in orienting DTC migration; however, their precise roles are uncertain.

Live cell imaging coupled with single cell molecular manipulation studies, which have helped elucidate AC invasion, will be crucial for advancing our understanding of DTC migration (Kelley *et al.* 2017). Dynamic methods to visualize and perturb Rho GTPases, F-actin, BM protein deposition, secretion, and removal, as well as trafficking, activity, and localization of integrins, the Wnt receptor **MOM-5** and the netrin receptors **UNC-40** and **UNC-5** should help clarify how

the DTC polarizes and migrates. Branching morphogenesis in the lung, mammary gland, kidney and salivary gland is driven by BM encased leader cells, which are not protrusive and have a smooth leading front like the DTC (Williams and Daniel 1983; Ewald *et al.* 2008; Andrew and Ewald 2010; Harunaga *et al.* 2014). Thus, a deeper understanding of the mechanisms underlying DTC migration will have significance to vertebrate organ formation. Further, as invasive tumor growth hijacks these morphogenetic mechanisms to spread into adjoining tissues (Gray *et al.* 2010), it is likely that examining DTC migration will reveal mechanisms that are misregulated in metastatic cancers.

Sex Myoblast Migration: Navigating Around Tissues To Position Muscles

Unlike the DTC, which directs a collective cell migration event while encased in a BM, many cells also migrate individually through and around animal tissues unencumbered by a surrounding BM. Examples of this mode of migration include primordial germ cells, leukocytes, hematopoietic stem cells, and tumor cells (Aman and Piotrowski 2010; Friedl and Alexander 2011). The SMs are two bilaterally symmetric cells, born at the end of the first larval stage among the posterior ventral body muscles. Each SM undergoes an individual long-distance migration along the left and right sides of the worm body cavity. In the hermaphrodite, ~2 hr after the SMs are formed and over the course of L2 and early L3 stages of development, the SMs migrate anteriorly to a final position at the exact center of the gonad, where the future uterine-vulval attachment will form (Figure 7). Following migration, the SM cells divide to form 16 vulval and uterine muscle cells (the sex muscles), which make attachments to the uterus, the vulva, and lateral epidermis (Sulston and Horvitz 1977; Lints and Hall 2017). The SMs migrate ~65 μm in 4 hr (Branda and Stern 2000), and are not encased in BM (D. R. Sherwood and E. L. Hastie, unpublished data). SMs in male animals migrate toward the tail, and differentiate into the tail muscles used for mating (Sulston and Horvitz 1977). Little is known about male SM migration, except that it is cell-autonomous: in an otherwise hermaphrodite animal, mosaic loss of *tra-1*, a gene linked with sex determination, creates male SMs that migrate posteriorly and produce male-like sex muscles in the tail (Hunter and Wood 1990). In the following, we discuss SM migration in the hermaphrodite.

Attraction and repulsion in SM migration

SM migration depends on a complex mix of both attractive and repulsive signals originating from somatic gonadal as well as nongonadal tissues that precisely position the SM cells at the uterine-vulval connection [for review see Chen and Stern (1998); Figure 7]. During the L2 and early L3 stages, the central part of the gonad is composed of six somatic (non-germline) cells: three ventral uterine cells, two dorsal uterine cells and the AC. The AC marks the center of the gonad and aligns with the vulval precursor cell P6.p. During the late L2

stage the AC secretes the EGF-like ligand *LIN-3*, which induces the P6.p cell to take on the 1° vulval precursor fate (see *AC invasion* section) (Sternberg 2005). In *dig-1* mutants, where the gonad is displaced anteriorly, and sometimes also dorsally, SMs still target the gonad center, navigating through new territory to precisely center on the displaced AC (Thomas *et al.* 1990). When the gonad is entirely deleted by laser ablation, SMs nevertheless still migrate, but their final positions cover a broad, centrally dispersed region (Thomas *et al.* 1990). SM migration therefore seems to be guided by a combination of gonad-dependent and gonad-independent attractive signals that direct the SM to the site of uterine-vulval connection.

The fibroblast growth factor *EGL-17* (FGF) is a key attractive cue that helps guide SM migration (Burdine *et al.* 1997, 1998; Branda and Stern 2000). Vertebrates encode 23 fibroblast growth factor (FGF) ligands and five FGF receptors, while *C. elegans* has just two FGF ligands (*EGL-17* and *LET-756*) and one receptor (*EGL-15*) (Borland *et al.* 2001). In *egl-17* mutant animals, SMs display severe premature termination of migration, stopping ~50 μm short of the gonad center (Stern and Horvitz 1991). Strikingly, gonad ablation in *egl-17* mutant animals restores the broad positioning around P6.p as observed in wild-type animals with ablated gonads. These experiments suggest that *EGL-17* (FGF) is an attractive cue for SM migration and that the gonad generates an unknown repulsive cue. Further, these results indicate that, in the absence of both the attractive cue (*EGL-17*) and the repulsive cue from the gonad, a separate nongonadal (and non-*EGL-17*) signal acts to help position the SM cells. It is unknown how the presence of *EGL-17* (FGF) masks the repulsive effect of the gonad during normal development, or what the physiological role of repulsion might be.

The *egl-17* gene is expressed in the P6.p vulval precursor cell after it is induced to a 1° vulval fate by the gonadal AC, thus providing a source of *EGL-17* (FGF) protein that guides migration and coordinates SM migration with vulval induction during the early L3 stage (Burdine *et al.* 1998). Analysis of the promoter region of *egl-17* revealed three enhancer regions that control *egl-17* expression both temporally and spatially within the 1° fated P6.p cell (Cui and Han 2003; Sternberg 2005). When *egl-17* is expressed in neighboring vulval precursor cells, the SM targets to the cell expressing *egl-17* (Burdine *et al.* 1998). The vulval precursor cells can be ablated, however, and SMs still position precisely (Burdine *et al.* 1998). *EGL-17* (FGF) is also expressed in the dorsal uterine cells, and there is evidence for its presence in the ventral uterine cells and the AC as well (Branda and Stern 2000). Thus *egl-17* from the uterine cells within the gonad likely also guide SM migration (Branda and Stern 2000). Overall, these observations suggest that *EGL-17* (FGF) from multiple sources near the uterine-vulval connection overcomes the negative SM migration signal. Together these *EGL-17* (FGF) attractive cues have been referred to as the “gonad dependent signal,” as *egl-17* expression in the 1° fated P6.p depends on AC-mediated vulval induction. In addition




Figure 7 SM migration, a cell navigation event. Top panels: two SMs are specified by LIN-12 (Notch) signaling and the action of the zinc finger transcriptional regulator SEM-4 in the tail of the L1 larva and migrate independently of each other through the body of the worm over the course of the L2 and L3 larval stages until reaching the central gonad (shown with darker shading). The transcriptional regulator LIN-42 (Period) prevents the SMs from dividing precociously during their migration. Only one SM is shown for simplicity. Bottom panel: SMs are directed to the proper location by an EGL-17 (FGF) signal emanating from the central gonad region and vulval precursor cells, and by an additional unknown signal originating from nongonadal tissue. In addition to these attractive signals, an unidentified cue, also originating in the gonad, repels the SMs. Both EGL-17 (FGF)-dependent attractive and gonad-dependent repulsive signals are sensed by the FGF receptor EGL-15 in the SM, but different isoforms (5A and 5B) respond to attractive and repulsive cues. Signals are transduced to LET-60 (Ras) via the adaptor molecule SEM-5 (Grb2), which may also communicate to Arp2/3 complex-driven actin assembly. EGL-17 (FGF)-independent attraction is less well-understood, but depends on various molecules with potential roles in cell adhesion and motility such as UNC-53 (Nav2), UNC-71 (ADAM), and UNC-73 (Trio).

to directing SMs toward the gonad, EGL-17 (FGF) acts with the *C. elegans* Robo receptor SAX-3 to maintain SMs ventrally during their migration (Branda and Stern 2000).

The *C. elegans* FGF receptor EGL-15 (FGFR) is expressed in SMs and is necessary for correct targeting of the gonad by the SMs (DeVore *et al.* 1995; Sundaram *et al.* 1996; Branda and Stern 2000; Lo *et al.* 2008). It was originally reported that mutation of EGL-15 phenocopies what is observed with *egl-17* mutant worms, with posteriorly displaced SMs indicating that *egl-17* mediated attraction is turned off, while gonad repulsion is maintained (Stern and Horvitz 1991). However, later it was discovered that alternative splicing generates two EGL-15 (FGFR) isoforms, 5A and 5B, which each have a unique domain in their extracellular region after the first IG motif of EGL-15 (FGFR) (Burdine *et al.* 1997, 1998; Goodman *et al.* 2003; Huang and Stern 2004; Lo *et al.* 2008). The original *egl-15* allele interferes with only the splicing of isoform 5A, leaving 5B intact (Goodman *et al.* 2003). Loss of both EGL-15 (FGFR) isoforms leads to SMs that are loosely centered on the AC—a phenotype observed with gonad ablation (*i.e.*, when both gonad repulsion and attraction are lost) (Lo *et al.* 2008). Site of action studies similarly suggest that 5A is the form of the FGF receptor that mediates attraction to EGL-17 (FGF) sources, while 5B has a role in gonad-dependent repulsion. Expression and genetic evidence suggests that repulsion is mediated through an unidentified ligand that is not the other *C. elegans* FGF, LET-756 (Lo *et al.* 2008).

It is not uncommon for ligands and receptors associated with migration to play both attractive and repulsive roles. This occurs with semaphorins and ephrin signaling systems (Klein 2012; Gurrupu and Tamagnone 2016). As discussed earlier, this is also how the netrin signaling system functions, as the ligand UNC-6 (netrin) attracts cells that express the receptor UNC-40 (DCC) and repels cells expressing the receptor UNC-5 and UNC-40 together or UNC-5 alone (Lai Wing Sun *et al.* 2011). There is evidence that opposing signals might act by biasing self-organizing stochastic intracellular polarity systems (Tang and Wadsworth 2014; Chisholm *et al.* 2016). For example, UNC-6 (netrin) stabilizes where randomly directed self-organizing UNC-40 polarizes during AC invasion (Wang *et al.* 2014b). Whatever the mechanism, the FGF-dependent guidance system for SMs seems to be a conserved: *Ppa-egl-17* phenotypes and ablation experiments indicate that gonad-dependent and independent attraction and gonad-dependent repulsion mechanisms are operational in the evolutionarily distant *Pristionchus pacificus* (Photos *et al.* 2006).

Ras in SM positioning

LET-60 (Ras) is a small GTPase that is part of the core *C. elegans* RTK (Receptor Tyrosine Kinase)/Ras/MAPK (Mitogen-Activated Protein Kinase) signaling pathway, which converts growth factor signals to cellular transformations via LET-60 (Ras)-dependent intracellular phosphorylation events (Sundaram 2013). LET-60 is expressed in the SMs and the

somatic gonad, as well as in the vulval precursor cells, where *EGL-17* (FGF) expression is dependent on induction by *LET-60* (Burdine *et al.* 1998; Dent and Han 1998). The *LET-60* (Ras) signaling cascade begins with *EGL-17* (FGF) binding to *EGL-15* (FGFR) on the cell surface, which triggers dimerization and autophosphorylation of the intracellular domain of *EGL-15* (FGFR), promoting docking of the adaptor protein *SEM-5* (Grb2) ((Sundaram 2013) and Figure 7). In *C. elegans*, *EGL-15* (FGFR) binds directly to *SEM-5* (Grb2), unlike in vertebrates, which require an additional FRS2 molecule to link FGFRs to Grb2 (Borland *et al.* 2001; Lo *et al.* 2010). *SEM-5* associates with the Ras-GEF, *SOS-1* (also known as *LET-341*), which activates *LET-60*, leading to a variety of downstream signaling events including phosphorylations via the MAP kinase cascade and nuclear translocations that change gene expression (Borland *et al.* 2001; Sundaram 2013).

Mutations in the *sem-5* (Grb2) gene lead to posteriorly displaced SMs, but the defects are far weaker than *egl-17* mutants, and SM distributions more closely resemble those of gonad-ablated animals, suggesting that both gonad-dependent repulsion and attraction are eliminated when *SEM-5* is not functional (Clark *et al.* 1992; DeVore *et al.* 1995; Chen *et al.* 1997). Likewise, *let-60* loss-of-function mutations lead to SMs that are loosely centered on the AC, indicating loss of *EGL-15* (FGFR) mediated activity, both instructive and repulsive (Sundaram 2013). When *LET-60* (Ras) is restored mosaically in one SM cell, but not the other, the genetically wild-type SM shows rescue of positioning and the mutant SM does not, indicating that *LET-60* (Ras) is required cell-autonomously (Sundaram 2013). Mutations in Ras regulators, such as *ksr-1*, the *C. elegans* equivalent of the positive Ras regulator KSR, give phenotypes like those observed for *let-60* deletion (Ohmachi *et al.* 2002; Oakes *et al.* 2012); similarly loss of RasGAPs that control Ras activity also affects SM migration (Stetak *et al.* 2008). Overall, *LET-60* and its associated adaptors play a major role in guidance of SM migration by the gonad, via both attraction and repulsion.

FGF independent mechanisms of SM migration

The *EGL-17*/*EGL-15*-independent migration of SMs, revealed in the absence of the gonad, is not well understood, but mutations in the genes *unc-53*, *unc-71*, and *unc-73* (corresponding to mammalian Nav2, ADAM, and Trio, respectively) abrogate this migration mode and SMs are posteriorly displaced in gonad-ablated animals instead of being loosely grouped around the AC (Chen *et al.* 1997). Mutations in these genes have no effect on *EGL-15* (FGFR)-dependent SM migration. The posterior SM displacement of *unc-53* gonad-ablated animals is worsened by mutations in *let-60* and *sem-5*, indicating that *LET-60* (Ras) and *SEM-5* (Grb2) also contribute to gonad-independent SM migration, seemingly in addition to their participation in the FGFR signaling pathway (Chen *et al.* 1997; Chen and Stern 1998). Similarly, in gonad-ablated animals, expression of constitutively active *LET-60* (Ras) produces an almost wild-type targeting of SMs, reducing the broad distribution usually observed in gonad-ablated animals (Sundaram 2013). Taken together, these results suggest that SM migration is controlled by *EGL-15* (FGFR)

signaling to *SEM-5* (Grb2) and *LET-60* (Ras), which also play FGFR-independent roles in SM movement by an undetermined signaling pathway.

Possible signaling to the actin cytoskeleton for SM migration

Due to the different compensatory and competing mechanisms driving SM movement, it is somewhat difficult to link genetic interactions to molecular mechanisms of SM migration, and little is known concerning membrane or cytoskeleton dynamics in the SM downstream of the different signaling pathways described above. However drawing analogies with what is known in mammalian systems, *SEM-5* (Grb2) could directly link *EGL-15* (FGFR) to the actin cytoskeleton, since Grb2 is a known activator of the actin polymerization factor N-WASP, enhancing its ability to activate actin polymerization nucleation via the Arp2/3 complex (Carrier *et al.* 2000, and Figure 7). Additionally the molecules involved in gonad-independent SM migration have connections to the actin cytoskeleton: *UNC-53* (NAV-2) physically interacts with *ABI-1*, part of the WAVE complex, as well as *SEM-5*, thus potentially linking *UNC-53* to WASP activity as well (Stringham *et al.* 2002; Schmidt *et al.* 2009). *UNC-71* (ADAM) is an ADAM protein, lacking protease activity, and involved in cell adhesion during cell motility possibly via integrins (X. Huang *et al.* 2003) and *UNC-73* (Trio) is a GEF that accelerates the activity of the Rac GTPases *CED-10* and *MIG-2*. *MIG-2* is expressed in SMs, and expression of a constitutively active *MIG-2* protein perturbs SM migration (Zipkin *et al.* 1997). *CED-10* and *MIG-2* are the upstream activators of the WAVE complex and WASP, respectively, for actin polymerization via the Arp 2/3 complex (Shakir *et al.* 2008; Walck-Shannon *et al.* 2015). All of this points to a possible scenario in which signals arriving from the gonad, and other, as-yet-unidentified, sources signal to the actin cytoskeleton and adhesion systems of the SM to enable its migration and correct positioning at the center of the gonad.

Specification of migratory SM cells

Specification of the SM fate from surrounding mesodermal precursor cells initiates before movement begins (Figure 7). Genetic analysis has indicated that SM cells require *LIN-12* (Notch) activity in the mid-to-late L1 stage for initial specification (Greenwald *et al.* 1983; Foehr and Liu 2008). In the absence of the *lin-12* gene, the SM precursor cells adopt the coelomocyte fate (another mesoderm progenitor fate). *LIN-12* (Notch) is activated by its transmembrane ligands *LAG-2* and *APX-1*, which are expressed in adjacent epidermal cells in the mid-to-late L1 (Foehr and Liu 2008). A zinc finger-type transcription factor encoded by the gene *sem-4*, appears to function after *LIN-12* (Notch) activity. In the absence of *sem-4*, cells resembling the SMs are generated, but they do not migrate toward the vulva or divide, and instead take on the appearance of body wall muscle cells (Basson and Horvitz 1996). *SEM-4*, which is an ortholog of *Drosophila* SPALT, is conserved in *P. pacificus*, and as in *C. elegans*, disruption of *Ppa-sem-4* causes SM fate specification to fail (Photos *et al.* 2006).

Once SM migration halts, and they are positioned near the vulval precursor cells in the L3 stage, the SM cells divide and

differentiate into four types of sex muscles [for mechanisms regulating postmigratory SM lineage differentiation, see Hale *et al.* (2014)]. Molecular mechanisms regulate the timing of SM division, thereby assuring that SMs arrive at their vulval destination before proceeding with their developmental program and dividing to create the sex muscles. Loss of **LIN-42**, the *C. elegans* ortholog of the Per family of circadian rhythm proteins, causes SMs to prematurely undergo division during the L2 migratory stage (Tennessen *et al.* 2006). **LIN-42** (PER) is believed to interfere with transcriptional activators, thus repressing the expression of target genes. **EGL-15** (FGFR), while necessary for **EGL-17** (FGF)-dependent SM attraction and gonad-dependent repulsion, is a negative regulator of terminal differentiation of SMs into muscle cells. When hyperactivation of **EGL-15** (FGFR) is induced via constitutive dimerization, the vulval muscles descendant from the SMs do not form properly; in particular actin cables and myosin are reduced (Sasson and Stern 2004). This phenotype is strongly dependent on the 5A isoform, and less so on 5B, mirroring the isoform specificities for SM positioning. Additionally, the function of **EGL-15** (FGFR) in sex muscle differentiation is controlled by glycosylation. When a key glycosylation site in **EGL-15** (FGFR) is mutated, an **EGL-15** hyperactivation phenotype is observed, with incorrect formation of the vulval muscles. This result indicates that glycosylation negatively regulates the role of **EGL-15** (FGFR) in SM differentiation to muscle. Glycosylation, however, has no effect on SM migration (Polanska *et al.* 2009), suggesting that glycosylation could be a means of inactivating **EGL-15** (FGFR) specifically during differentiation.

Unanswered questions in SM migration

Many questions remain to be addressed concerning SM migration, including the identity of the gonad-dependent repellant signal, how the presence of **EGL-17** (FGF) masks the repulsive effect, and what physiological function repulsion serves in SM focusing. Additionally, the signals and receptors involved in gonad-independent attraction are unknown. Furthermore, little is known about the molecular details regarding how signals are transduced intracellularly leading to gonad-dependent attraction/repulsion, and whether the motility mechanism is different in gonad-dependent vs. -independent movement. Although actin assembly is surely involved in SM movement, the actin cytoskeleton has not been imaged in moving SMs, nor has the cytoskeleton been examined under conditions that affect SM movement. Real-time imaging has not yet been performed on the migrating SM cells, and we do not know what migration mode—mesenchymal, bleb-based, or novel—they use to move. Finally, since male SMs placed in a hermaphrodite environment still migrate toward their normal posterior location, SM migration could serve as a powerful model to determine how the intrinsic programs of migrating cells control the manner in which they respond to extrinsic guidance cues.

Summary and outlook

Cell migration is crucial during tissue formation and homeostasis in animals, and cells have developed a rich toolkit to drive and control movement in different tissue environments (Aman and

Piotrowski 2010; Mayor and Etienne-Manneville 2016). Our review of three types of cell movement in *C. elegans* has focused on distinct types of movements found in animals: AC invasion, a model for how cells breach BM barriers; DTC migration, an example of how a leader cell encased in a BM directs organ shape; and SM migration, a model for how individual cells migrate between tissues. The importance of studying cell movement *in vivo* is highlighted by many novel findings from these models. These include (1) in the case of the AC, the identification of invadopodia in a native tissue environment, the discovery that the netrin ligand stabilizes intrinsic oscillatory clustering of its receptor **UNC-40** (DCC) to polarize cells, the finding that BM sliding widens BM openings, and that the AC induces cell shape changes when it contacts the underlying vulval precursor cells; (2) in the case of DTCs, the elucidation of a Netrin-Wnt global positioning system that orients migration along the A/V and D/V axes, and a role for conserved microRNAs in buffering cell migration to environmental fluctuations in temperature; and (3) in the case of SMs, the intricate roles played by FGF/FGFR-dependent and independent signaling. These *in vivo* studies have also revealed the key functions that dynamic transcriptional programs play in changing the ways in which migrating cells respond to their environment to alter movement. Finally, work in *C. elegans* has identified significant areas for future study of cell motility, such as the puzzling observation that the DTC is largely nonprotrusive and moves by means that are currently poorly understood.

One important implication from these studies in *C. elegans* is that a deep understanding of cell migration requires the ability to visualize migration, to view the subcellular structures and molecular components (as well as their activity) that regulate migration, and to simultaneously perturb this molecular machinery. The challenge of combining these approaches *in vivo* is a key reason why *in vitro* studies of cell migration have been so predominant in examination of cell movement. However, *in vivo* systems are needed to confirm and extend these *in vitro* findings. With its amenability to live cell imaging, simple tissue architecture, and temporally and spatially controlled methods for gene and protein manipulation (Chai *et al.* 2012; Wei *et al.* 2012; Armenti *et al.* 2014; Corsi *et al.* 2015; Zhang *et al.* 2015), *C. elegans* is emerging as more than just a model to identify genes involved in migration. The worm is developing into a powerful *in vivo* system that should remain at the forefront of discovering key conserved cellular and molecular mechanisms that allow cells to enact exquisite migratory programs to navigate through diverse tissue environments.

Acknowledgments

We thank Agnieszka Kawska of IlluScientia.com for her work on the figures, and thank Naomi Levy-Strumpf, Jane Hubbard, Andy Ewald, Peter Friedl, and David Matus for helpful discussions on the material covered. D.R.S. was supported by National Institute of General Medical Sciences (NIGMS) R01 GM079320, National Institute of Child Health and Human Development (NICHD) R21HD084290 and NIGMS R35 Maximizing Investors' Research Award (MIRA) GM118049. J.P.

was supported by Fondation Association pour la Recherche sur le Cancer (ARC) and the Idex ANR-10-IDEX-0001-02 PSL.

Literature Cited

- Abella, J. V., C. Galloni, J. Pernier, D. J. Barry, S. Kjaer *et al.*, 2016 Isoform diversity in the Arp2/3 complex determines actin filament dynamics. *Nat. Cell Biol.* 18: 76–86.
- Altincicek, B., M. Fischer, K. Luerssen, M. Boll, U. Wenzel *et al.*, 2010 Role of matrix metalloproteinase ZMP-2 in pathogen resistance and development in *Caenorhabditis elegans*. *Dev. Comp. Immunol.* 34: 1160–1169.
- Aman, A., and T. Piotrowski, 2010 Cell migration during morphogenesis. *Dev. Biol.* 341: 20–33.
- Andrew, D. J., and A. J. Ewald, 2010 Morphogenesis of epithelial tubes: insights into tube formation, elongation, and elaboration. *Dev. Biol.* 341: 34–55.
- Antebi, A., J. G. Culotti, and E. M. Hedgecock, 1998 *daf-12* regulates developmental age and the dauer alternative in *Caenorhabditis elegans*. *Development* 125: 1191–1205.
- Antebi, A., W. H. Yeh, D. Tait, E. M. Hedgecock, and D. L. Riddle, 2000 *daf-12* encodes a nuclear receptor that regulates the dauer diapause and developmental age in *C. elegans*. *Genes Dev.* 14: 1512–1527.
- Armenti, S. T., L. L. Lohmer, D. R. Sherwood, and J. Nance, 2014 Repurposing an endogenous degradation system for rapid and targeted depletion of *C. elegans* proteins. *Development* 141: 4640–4647.
- Austin, J., and J. Kimble, 1987 *glp-1* is required in the germ line for regulation of the decision between mitosis and meiosis in *C. elegans*. *Cell* 51: 589–599.
- Basson, M., and H. R. Horvitz, 1996 The *Caenorhabditis elegans* gene *sem-4* controls neuronal and mesodermal cell development and encodes a zinc finger protein. *Genes Dev.* 10: 1953–1965.
- Baum, P. D., and G. Garriga, 1997 Neuronal migrations and axon fasciculation are disrupted in *ina-1* integrin mutants. *Neuron* 19: 51–62.
- Beaty, B. T., and J. Condeelis, 2014 Digging a little deeper: the stages of invadopodium formation and maturation. *Eur. J. Cell Biol.* 93: 438–444.
- Beerling, E., L. Ritsma, N. Vriskoop, P. W. Derksen, and J. van Rhee, 2011 Intravital microscopy: new insights into metastasis of tumors. *J. Cell Sci.* 124: 299–310.
- Benard, C. Y., A. Boyanov, D. H. Hall, and O. Hobert, 2006 *DIG-1*, a novel giant protein, non-autonomously mediates maintenance of nervous system architecture. *Development* 133: 3329–3340.
- Bendezu, F. O., and S. G. Martin, 2013 *Cdc42* explores the cell periphery for mate selection in fission yeast. *Curr. Biol.* 23: 42–47.
- Bergert, M., A. Erzberger, R. A. Desai, I. M. Aspalter, A. C. Oates *et al.*, 2015 Force transmission during adhesion-independent migration. *Nat. Cell Biol.* 17: 524–529.
- Bergman, A., J. S. Condeelis, and B. Gligorijevic, 2014 Invadopodia in context. *Cell Adhes. Migr.* 8: 273–279.
- Blanchoin, L., R. Boujemaa-Paterski, C. Sykes, and J. Plastino, 2014 Actin dynamics, architecture, and mechanics in cell motility. *Physiol. Rev.* 94: 235–263.
- Blelloch, R., and J. Kimble, 1999 Control of organ shape by a secreted metalloprotease in the nematode *Caenorhabditis elegans*. *Nature* 399: 586–590.
- Blelloch, R., S. S. Anna-Arriola, D. Gao, Y. Li, J. Hodgkin *et al.*, 1999 The *gon-1* gene is required for gonadal morphogenesis in *Caenorhabditis elegans*. *Dev. Biol.* 216: 382–393.
- Bokel, C., and N. H. Brown, 2002 Integrins in development: moving on, responding to, and sticking to the extracellular matrix. *Dev. Cell* 3: 311–321.
- Borland, C. Z., J. L. Schutzman, and M. J. Stern, 2001 Fibroblast growth factor signaling in *Caenorhabditis elegans*. *Bioessays* 23: 1120–1130.
- Branch, K. M., D. Hoshino, and A. M. Weaver, 2012 Adhesion rings surround invadopodia and promote maturation. *Biol. Open* 1: 711–722.
- Branda, C. S., and M. J. Stern, 2000 Mechanisms controlling sex myoblast migration in *Caenorhabditis elegans* hermaphrodites. *Dev. Biol.* 226: 137–151.
- Bravo-Cordero, J. J., M. A. Magalhaes, R. J. Eddy, L. Hodgson, and J. Condeelis, 2013 Functions of cofilin in cell locomotion and invasion. *Nat. Rev. Mol. Cell Biol.* 14: 405–415.
- Brodsky, L., I. Kolotuev, C. Didier, A. Bhoumik, B. Podbilewicz *et al.*, 2004 The LIM domain protein UNC-95 is required for the assembly of muscle attachment structures and is regulated by the RING finger protein RNF-5 in *C. elegans*. *J. Cell Biol.* 165: 857–867.
- Burdine, R. D., E. B. Chen, S. F. Kwok, and M. J. Stern, 1997 *egl-17* encodes an invertebrate fibroblast growth factor family member required specifically for sex myoblast migration in *Caenorhabditis elegans*. *Proc. Natl. Acad. Sci. USA* 94: 2433–2437.
- Burdine, R. D., C. S. Branda, and M. J. Stern, 1998 EGL-17(FGF) expression coordinates the attraction of the migrating sex myoblasts with vulval induction in *C. elegans*. *Development* 125: 1083–1093.
- Burke, S. L., M. Hammell, and V. Ambros, 2015 Robust distal tip cell pathfinding in the face of temperature stress is ensured by two conserved microRNAs in *Caenorhabditis elegans*. *Genetics* 200: 1201–1218.
- Burket, C. T., C. E. Higgins, L. C. Hull, P. M. Berninsone, and E. F. Ryder, 2006 The *C. elegans* gene *dig-1* encodes a giant member of the immunoglobulin superfamily that promotes fasciculation of neuronal processes. *Dev. Biol.* 299: 193–205.
- Cabello, J., L. J. Neukomm, U. Gunesdogan, K. Burkart, S. J. Charette *et al.*, 2010 The Wnt pathway controls cell death engulfment, spindle orientation, and migration through CED-10/Rac. *PLoS Biol.* 8: e1000297.
- Cal, S., and C. Lopez-Otin, 2015 ADAMTS proteases and cancer. *Matrix Biol.* 44–46: 77–85.
- Campbell, I. D., and M. J. Humphries, 2011 Integrin structure, activation, and interactions. *Cold Spring Harb. Perspect. Biol.* 3: a004994.
- Cardenas, M. G., E. Oswald, W. Yu, F. Xue, A. D. MacKerell, Jr. *et al.*, 2017 The expanding role of the BCL6 oncoprotein as a cancer therapeutic target. *Clin. Cancer Res.* 23: 885–893.
- Carlier, M.-F., P. Nioche, I. Broutin-ElHermite, R. Boujemaa, C. Le Clainche *et al.*, 2000 GRB2 links signaling to actin assembly by enhancing interaction of neural Wiskott-Aldrich syndrome protein (N-WASP) with actin-related protein (Arp2/3) complex. *J. Biol. Chem.* 275: 21946–21952.
- Castro-Castro, A., V. Marchesin, P. Monteiro, C. Lodillinsky, C. Rosse *et al.*, 2016 Cellular and molecular mechanisms of MT1-MMP-dependent cancer cell invasion. *Annu. Rev. Cell Dev. Biol.* 32: 555–576.
- Cecchetelli, A. D., and E. J. Cram, 2017 Regulating distal tip cell migration in space and time. *Mech. Dev.* pii: S0925–4773(16)30131–9. DOI: 10.1016/j.mod.2017.04.003.
- Chai, Y., W. Li, G. Feng, Y. Yang, X. Wang *et al.*, 2012 Live imaging of cellular dynamics during *Caenorhabditis elegans* postembryonic development. *Nat. Protoc.* 7: 2090–2102.
- Chan, S. S., H. Zheng, M. W. Su, R. Wilk, M. T. Killeen *et al.*, 1996 UNC-40, a *C. elegans* homolog of DCC (deleted in colorectal cancer), is required in motile cells responding to UNC-6 netrin cues. *Cell* 87: 187–195.
- Charras, G., and E. Paluch, 2008 Blebs lead the way: how to migrate without lamellipodia. *Nat. Rev. Mol. Cell Biol.* 9: 730–736.
- Chen, E. B., and M. J. Stern, 1998 Understanding cell migration guidance: lessons from sex myoblast migration in *C. elegans*. *Trends Genet.* 14: 322–327.
- Chen, E. B., C. S. Branda, and M. J. Stern, 1997 Genetic enhancers of *sem-5* define components of the gonad-independent

- guidance mechanism controlling sex myoblast migration in *Caenorhabditis elegans* hermaphrodites. *Dev. Biol.* 182: 88–100.
- Chen, P., J. D. Edelman, and S. A. Gharib, 2013 Comparative evaluation of miRNA expression between in vitro and in vivo airway epithelium demonstrates widespread differences. *Am. J. Pathol.* 183: 1405–1410.
- Chen, W. T., 1989 Proteolytic activity of specialized surface protrusions formed at rosette contact sites of transformed cells. *J. Exp. Zool.* 251: 167–185.
- Chesarone, M. A., and B. L. Goode, 2009 Actin nucleation and elongation factors: mechanisms and interplay. *Curr. Opin. Cell Biol.* 21: 28–37.
- Chesney, M. A., N. Lam, D. E. Morgan, B. T. Phillips, and J. Kimble, 2009 *C. elegans* HLH-2/E/Daughterless controls key regulatory cells during gonadogenesis. *Dev. Biol.* 331: 14–25.
- Chisholm, A. D., H. Hutter, Y. Jin, and W. G. Wadsworth, 2016 The genetics of axon guidance and axon regeneration in *Caenorhabditis elegans*. *Genetics* 204: 849–882.
- Ci, W., J. M. Polo, and A. Melnick, 2008 B-cell lymphoma 6 and the molecular pathogenesis of diffuse large B-cell lymphoma. *Curr. Opin. Hematol.* 15: 381–390.
- Clark, S. G., M. J. Stern, and H. R. Horvitz, 1992 *C. elegans* cell-signalling gene *sem-5* encodes a protein with SH2 and SH3 domains. *Nature* 356: 340–344.
- Clay, M. R., and D. R. Sherwood, 2015 Basement membranes in the worm: a dynamic scaffolding that instructs cellular behaviors and shapes tissues. *Curr. Top. Membr.* 76: 337–371.
- Clevers, H., and E. Batlle, 2013 SnapShot: the intestinal crypt. *Cell* 152: 1198–1198.e2.
- Clevers, H., and R. Nusse, 2012 Wnt/beta-catenin signaling and disease. *Cell* 149: 1192–1205.
- Corsi, A. K., B. Wightman, and M. Chalfie, 2015 A transparent window into biology: a primer on *Caenorhabditis elegans* (June 18, 2015) *WormBook*, ed. The *C. elegans* Research Community, *WormBook*, doi/10.1895/wormbook.1.177.1, <http://www.wormbook.org>.
- Coussens, L. M., B. Fingleton, and L. M. Matrisian, 2002 Matrix metalloproteinase inhibitors and cancer: trials and tribulations. *Science* 295: 2387–2392.
- Cram, E. J., S. G. Clark, and J. E. Schwarzbauer, 2003 Talin loss-of-function uncovers roles in cell contractility and migration in *C. elegans*. *J. Cell Sci.* 116: 3871–3878.
- Cram, E. J., H. Shang, and J. E. Schwarzbauer, 2006 A systematic RNA interference screen reveals a cell migration gene network in *C. elegans*. *J. Cell Sci.* 119: 4811–4818.
- Cross, S. S., J. Lippitt, A. Mitchell, F. Hollingsbury, S. P. Balasubramanian *et al.*, 2008 Expression of beta-dystroglycan is reduced or absent in many human carcinomas. *Histopathology* 53: 561–566.
- Cui, M., and M. Han, 2003 *Cis* regulatory requirements for vulval cell-specific expression of the *Caenorhabditis elegans* fibroblast growth factor gene *egl-17*. *Dev. Biol.* 257: 104–116.
- Darom, A., U. Bening-Abu-Shach, and L. Broday, 2010 RNF-121 is an endoplasmic reticulum-membrane E3 ubiquitin ligase involved in the regulation of beta-integrin. *Mol. Biol. Cell* 21: 1788–1798.
- Dent, J. A., and M. Han, 1998 Post-embryonic expression pattern of *C. elegans* *let-60 ras* reporter constructs. *Mech. Dev.* 72: 179–182.
- Destaing, O., M. R. Block, E. Planus, and C. Albiges-Rizo, 2011 Invadosome regulation by adhesion signaling. *Curr. Opin. Cell Biol.* 23: 597–606.
- DeVore, D. L., H. R. Horvitz, and M. J. Stern, 1995 An FGF receptor signaling pathway is required for the normal cell migrations of the sex myoblasts in *C. elegans* hermaphrodites. *Cell* 83: 611–620.
- Dickinson, D. J., J. D. Ward, D. J. Reiner, and B. Goldstein, 2013 Engineering the *Caenorhabditis elegans* genome using Cas9-triggered homologous recombination. *Nat. Methods* 10: 1028–1034.
- Dieterich, C., S. W. Clifton, L. N. Schuster, A. Chinwalla, K. Delehaunty *et al.*, 2008 The *Pristionchus pacificus* genome provides a unique perspective on nematode lifestyle and parasitism. *Nat. Genet.* 40: 1193–1198.
- Di Martino, J., L. Paysan, C. Gest, V. Lagree, A. Juin *et al.*, 2014 Cdc42 and Tks5: a minimal and universal molecular signature for functional invadosomes. *Cell Adhes. Migr.* 8: 280–292.
- Ding, M., A. Goncharov, Y. Jin, and A. D. Chisholm, 2003 *C. elegans* ankyrin repeat protein VAB-19 is a component of epidermal attachment structures and is essential for epidermal morphogenesis. *Development* 130: 5791–5801.
- Doherty, M. F., G. Adelmant, A. D. Cecchetelli, J. A. Marto, and E. J. Cram, 2014 Proteomic analysis reveals CACN-1 is a component of the spliceosome in *Caenorhabditis elegans*. *G3 Bethesda* 4: 1555–1564.
- Drabikowski, K., A. Trzebiatowska, and R. Chiquet-Ehrismann, 2005 *ten-1*, an essential gene for germ cell development, epidermal morphogenesis, gonad migration, and neuronal path-finding in *Caenorhabditis elegans*. *Dev. Biol.* 282: 27–38.
- Duan, S., L. Cermak, J. K. Pagan, M. Rossi, C. Martinengo *et al.*, 2012 FBXO11 targets BCL6 for degradation and is inactivated in diffuse large B-cell lymphomas. *Nature* 481: 90–93.
- Dufour, A., and C. M. Overall, 2013 Missing the target: matrix metalloproteinase antitargets in inflammation and cancer. *Trends Pharmacol. Sci.* 34: 233–242.
- Dyer, J. M., N. S. Savage, M. Jin, T. R. Zyla, T. C. Elston *et al.*, 2013 Tracking shallow chemical gradients by actin-driven wandering of the polarization site. *Curr. Biol.* 23: 32–41.
- Eisenmann, D. M., 2005 Wnt signaling (June 25, 2005), *WormBook*, ed. The *C. elegans* Research Community, *WormBook*, doi/10.1895/wormbook.1.7.1, <http://www.wormbook.org>.
- Even-Ram, S., and K. M. Yamada, 2005 Cell migration in 3D matrix. *Curr. Opin. Cell Biol.* 17: 524–532.
- Ewald, A. J., A. Brenot, M. Duong, B. S. Chan, and Z. Werb, 2008 Collective epithelial migration and cell rearrangements drive mammary branching morphogenesis. *Dev. Cell* 14: 570–581.
- Fancsalszky, L., E. Monostori, Z. Farkas, E. Pourkarimi, N. Masoudi *et al.*, 2014 NDK-1, the homolog of NM23-H1/H2 regulates cell migration and apoptotic engulfment in *C. elegans*. *PLoS One* 9: e92687.
- Feil, R., and M. F. Fraga, 2012 Epigenetics and the environment: emerging patterns and implications. *Nat. Rev. Genet.* 13: 97–109.
- Fidler, A. L., C. E. Darris, S. V. Chetyrkin, V. K. Pedchenko, S. P. Boudko *et al.*, 2017 Collagen IV and basement membrane at the evolutionary dawn of metazoan tissues. *Elife* 6: e24176.
- Fielenbach, N., D. Guardavaccaro, K. Neubert, T. Chan, D. Li *et al.*, 2007 DRE-1: an evolutionarily conserved F box protein that regulates *C. elegans* developmental age. *Dev. Cell* 12: 443–455.
- Finger, F. P., K. R. Kopish, and J. G. White, 2003 A role for septins in cellular and axonal migration in *C. elegans*. *Dev. Biol.* 261: 220–234.
- Fletcher, S. J., and J. Z. Rappoport, 2010 Moving forward: polarised trafficking in cell migration. *Trends Cell Biol.* 20: 71–78.
- Foehr, M. L., and J. Liu, 2008 Dorsoventral patterning of the *C. elegans* postembryonic mesoderm requires both LIN-12/Notch and TGFbeta signaling. *Dev. Biol.* 313: 256–266.
- Fraley, S. I., Y. Feng, R. Krishnamurthy, D.-H. Kim, A. Celedon *et al.*, 2010 A distinctive role for focal adhesion proteins in three-dimensional cell motility. *Nat. Cell Biol.* 12: 598–604.
- Friedl, P., and S. Alexander, 2011 Cancer invasion and the micro-environment: plasticity and reciprocity. *Cell* 147: 992–1009.
- Friedl, P., and D. Gilmour, 2009 Collective cell migration in morphogenesis, regeneration and cancer. *Nat. Rev. Mol. Cell Biol.* 10: 445–457.

- Friedl, P., and K. Wolf, 2010 Plasticity of cell migration: a multi-scale tuning model. *J. Cell Biol.* 188: 11–19.
- Friedl, P., E. Sahai, S. Weiss, and K. M. Yamada, 2012 New dimensions in cell migration. *Nat. Rev. Mol. Cell Biol.* 13: 743–747.
- Genot, E., and B. Gligorijevic, 2014 Invadosomes in their natural habitat. *Eur. J. Cell Biol.* 93: 367–379.
- Geraldo, S., A. Simon, N. Elkhatib, D. Louvard, L. Fetler *et al.*, 2012 Do cancer cells have distinct adhesions in 3D collagen matrices and in vivo? *Eur. J. Cell Biol.* 91: 930–937.
- Gitai, Z., T. W. Yu, E. A. Lundquist, M. Tessier-Lavigne, and C. I. Bargmann, 2003 The netrin receptor UNC-40/DCC stimulates axon attraction and outgrowth through enabled and, in parallel, Rac and UNC-115/AbLIM. *Neuron* 37: 53–65.
- Goldstein, B., H. Takeshita, K. Mizumoto, and H. Sawa, 2006 Wnt signals can function as positional cues in establishing cell polarity. *Dev. Cell* 10: 391–396.
- Gonzales, K. A., H. Liang, Y. S. Lim, Y. S. Chan, J. C. Yeo *et al.*, 2015 Deterministic restriction on pluripotent state dissolution by cell-cycle pathways. *Cell* 162: 564–579.
- Goodman, S. J., C. S. Branda, M. K. Robinson, R. D. Burdine, and M. J. Stern, 2003 Alternative splicing affecting a novel domain in the *C. elegans* EGL-15 FGF receptor confers functional specificity. *Development* 130: 3757–3766.
- Graham, P. L., J. J. Johnson, S. Wang, M. H. Sibley, M. C. Gupta *et al.*, 1997 Type IV collagen is detectable in most, but not all, basement membranes of *Caenorhabditis elegans* and assembles on tissues that do not express it. *J. Cell Biol.* 137: 1171–1183.
- Gray, R. S., K. J. Cheung, and A. J. Ewald, 2010 Cellular mechanisms regulating epithelial morphogenesis and cancer invasion. *Curr. Opin. Cell Biol.* 22: 640–650.
- Greenwald, I., 2005 LIN-12/Notch signaling in *C. elegans* (August 4, 2005), *WormBook*, ed. The *C. elegans* Research Community, *WormBook*, doi/10.1895/wormbook.1.10.1, <http://www.wormbook.org>.
- Greenwald, I. S., P. W. Sternberg, and H. R. Horvitz, 1983 The lin-12 locus specifies cell fates in *Caenorhabditis elegans*. *Cell* 34: 435–444.
- Gurrapu, S., and L. Tamagnone, 2016 Transmembrane semaphorins: multimodal signaling cues in development and cancer. *Cell Adhes. Migr.* 10: 675–691.
- Hagedorn, E. J., and D. R. Sherwood, 2011 Cell invasion through basement membrane: the anchor cell breaches the barrier. *Curr. Opin. Cell Biol.* 23: 589–596.
- Hagedorn, E. J., H. Yashiro, J. W. Ziel, S. Ihara, Z. Wang *et al.*, 2009 Integrin acts upstream of netrin signaling to regulate formation of the anchor cell's invasive membrane in *C. elegans*. *Dev. Cell* 17: 187–198.
- Hagedorn, E. J., J. W. Ziel, M. A. Morrissey, L. M. Linden, Z. Wang *et al.*, 2013 The netrin receptor DCC focuses invadopodia-driven basement membrane transmigration in vivo. *J. Cell Biol.* 201: 903–913.
- Hagedorn, E. J., L. C. Kelley, K. M. Naegeli, Z. Wang, Q. Chi *et al.*, 2014 ADF/cofilin promotes invadopodial membrane recycling during cell invasion in vivo. *J. Cell Biol.* 204: 1209–1218.
- Haigo, S. L., and D. Bilder, 2011 Global tissue revolutions in a morphogenetic movement controlling elongation. *Science* 331: 1071–1074.
- Hale, J. J., N. M. Amin, C. George, Z. Via, H. Shi *et al.*, 2014 A role of the LIN-12/Notch signaling pathway in diversifying the non-striated egg-laying muscles in *C. elegans*. *Dev. Biol.* 389: 137–148.
- Hall, D. H., V. P. Winfrey, G. Blaeuer, L. H. Hoffman, T. Furuta *et al.*, 1999 Ultrastructural features of the adult hermaphrodite gonad of *Caenorhabditis elegans*: relations between the germ line and soma. *Dev. Biol.* 212: 101–123.
- Harburger, D. S., and D. A. Calderwood, 2009 Integrin signalling at a glance. *J. Cell Sci.* 122: 159–163.
- Harunaga, J. S., A. D. Doyle, and K. M. Yamada, 2014 Local and global dynamics of the basement membrane during branching morphogenesis require protease activity and actomyosin contractility. *Dev. Biol.* 394: 197–205.
- Hastie, E. L., and D. R. Sherwood, 2016 A new front in cell invasion: the invadopodial membrane. *Eur. J. Cell Biol.* 95: 441–448.
- Hedgecock, E. M., J. G. Culotti, D. H. Hall, and B. D. Stern, 1987 Genetics of cell and axon migrations in *Caenorhabditis elegans*. *Development* 100: 365–382.
- Hedgecock, E. M., J. G. Culotti, and D. H. Hall, 1990 The unc-5, unc-6, and unc-40 genes guide circumferential migrations of pioneer axons and mesodermal cells on the epidermis in *C. elegans*. *Neuron* 4: 61–85.
- Hermeking, H., 2010 The miR-34 family in cancer and apoptosis. *Cell Death Differ.* 17: 193–199.
- Hesselson, D., C. Newman, K. W. Kim, and J. Kimble, 2004 GON-1 and fibulin have antagonistic roles in control of organ shape. *Curr. Biol.* 14: 2005–2010.
- Hikasa, H., and S. Y. Sokol, 2013 Wnt signaling in vertebrate axis specification. *Cold Spring Harb. Perspect. Biol.* 5: a007955.
- Hilliard, M. A., and C. I. Bargmann, 2006 Wnt signals and frizzled activity orient anterior-posterior axon outgrowth in *C. elegans*. *Dev. Cell* 10: 379–390.
- Hotary, K., X. Y. Li, E. Allen, S. L. Stevens, and S. J. Weiss, 2006 A cancer cell metalloprotease triad regulates the basement membrane transmigration program. *Genes Dev.* 20: 2673–2686.
- Hrus, A., G. Lau, H. Hutter, S. Schenk, J. Ferralli *et al.*, 2007 *C. elegans* agrin is expressed in pharynx, IL1 neurons and distal tip cells and does not genetically interact with genes involved in synaptogenesis or muscle function. *PLoS One* 2: e731.
- Huang, C. C., D. H. Hall, E. M. Hedgecock, G. Kao, V. Karantza *et al.*, 2003 Laminin alpha subunits and their role in *C. elegans* development. *Development* 130: 3343–3358.
- Huang, P., and M. J. Stern, 2004 FGF signaling functions in the hypodermis to regulate fluid balance in *C. elegans*. *Development* 131: 2595–2604.
- Huang, T. F., C. Y. Cho, Y. T. Cheng, J. W. Huang, Y. Z. Wu *et al.*, 2014 BLMP-1/Blimp-1 regulates the spatiotemporal cell migration pattern in *C. elegans*. *PLoS Genet.* 10: e1004428.
- Huang, X., P. Huang, M. K. Robinson, M. J. Stern, and Y. Jin, 2003 UNC-71, a disintegrin and metalloprotease (ADAM) protein, regulates motor, axon guidance and sex myoblast migration in *C. elegans*. *Development* 130: 3147–3161.
- Hunter, C. P., and W. B. Wood, 1990 The tra-1 gene determines sexual phenotype cell-autonomously in *C. elegans*. *Cell* 63: 1193–1204.
- Huttenlocher, A., and A. R. Horwitz, 2011 Integrins in cell migration. *Cold Spring Harb. Perspect. Biol.* 3: a005074.
- Hutter, H., B. E. Vogel, J. D. Plenefisch, C. R. Norris, R. B. Proenca *et al.*, 2000 Conservation and novelty in the evolution of cell adhesion and extracellular matrix genes. *Science* 287: 989–994.
- Hwang, B. J., A. D. Meruelo, and P. W. Sternberg, 2007 *C. elegans* EVI1 proto-oncogene, EGL-43, is necessary for Notch-mediated cell fate specification and regulates cell invasion. *Development* 134: 669–679.
- Ihara, S., and K. Nishiwaki, 2007 Prodomain-dependent tissue targeting of an ADAMTS protease controls cell migration in *Caenorhabditis elegans*. *EMBO J.* 26: 2607–2620.
- Ihara, S., E. J. Hagedorn, M. A. Morrissey, Q. Chi, F. Motegi *et al.*, 2011 Basement membrane sliding and targeted adhesion remodels tissue boundaries during uterine-vulval attachment in *Caenorhabditis elegans*. *Nat. Cell Biol.* 13: 641–651.
- Ishii, N., W. G. Wadsworth, B. D. Stern, J. G. Culotti, and E. M. Hedgecock, 1992 UNC-6, a laminin-related protein, guides cell and pioneer axon migrations in *C. elegans*. *Neuron* 9: 873–881.
- Itoh, B., T. Hirose, N. Takata, K. Nishiwaki, M. Koga *et al.*, 2005 SRC-1, a non-receptor type of protein tyrosine kinase,

- controls the direction of cell and growth cone migration in *C. elegans*. *Development* 132: 5161–5172.
- Itoh, Y., 2015 Membrane-type matrix metalloproteinases: their functions and regulations. *Matrix Biol.* 44–46: 207–223.
- Jarrell, K. F., and M. J. McBride, 2008 The surprisingly diverse ways that prokaryotes move. *Nat. Rev. Microbiol.* 6: 466–476.
- Jayadev, R., and D. R. Sherwood, 2017 Basement membranes. *Curr. Biol.* 27: R207–R211.
- Jiang, H., G. Zhang, J. H. Wu, and C. P. Jiang, 2014 Diverse roles of miR-29 in cancer (review). *Oncol. Rep.* 31: 1509–1516.
- Jorgensen, E. M., and S. E. Mango, 2002 The art and design of genetic screens: *Caenorhabditis elegans*. *Nat. Rev. Genet.* 3: 356–369.
- Kamath, R. S., A. G. Fraser, Y. Dong, G. Poulin, R. Durbin *et al.*, 2003 Systematic functional analysis of the *Caenorhabditis elegans* genome using RNAi. *Nature* 421: 231–237.
- Kao, G., C. C. Huang, E. M. Hedgecock, D. H. Hall, and W. G. Wadsworth, 2006 The role of the laminin beta subunit in laminin heterotrimer assembly and basement membrane function and development in *C. elegans*. *Dev. Biol.* 290: 211–219.
- Karp, X., and I. Greenwald, 2004 Multiple roles for the E/Daughterless ortholog HLH-2 during *C. elegans* gonadogenesis. *Dev. Biol.* 272: 460–469.
- Kato, M., and P. W. Sternberg, 2009 The *C. elegans* tailless/Tlx homolog nhr-67 regulates a stage-specific program of linker cell migration in male gonadogenesis. *Development* 136: 3907–3915.
- Kato, M., T. F. Chou, C. Z. Yu, J. DeModena, and P. W. Sternberg, 2014 LINKIN, a new transmembrane protein necessary for cell adhesion. *Elife* 3: e04449.
- Kawano, T., H. Zheng, D. C. Merz, Y. Kohara, K. K. Tamai *et al.*, 2009 *C. elegans* mig-6 encodes papilin isoforms that affect distinct aspects of DTC migration, and interacts genetically with mig-17 and collagen IV. *Development* 136: 1433–1442.
- Kelley, L. C., L. L. Lohmer, E. J. Hagedorn, and D. R. Sherwood, 2014 Traversing the basement membrane in vivo: a diversity of strategies. *J. Cell Biol.* 204: 291–302.
- Kelley, L. C., Z. Wang, E. J. Hagedorn, L. Wang, W. Shen *et al.*, 2017 Live-cell confocal microscopy and quantitative 4D image analysis of anchor-cell invasion through the basement membrane in *Caenorhabditis elegans*. *Nat. Protoc.* 12: 2081–2096.
- Kelwick, R., I. Desanlis, G. N. Wheeler, and D. R. Edwards, 2015 The ADAMTS (A disintegrin and metalloproteinase with thrombospondin motifs) family. *Genome Biol.* 16: 113.
- Khayatan, B., J. C. Meeks, and D. D. Risser, 2015 Evidence that a modified type IV pilus-like system powers gliding motility and polysaccharide secretion in filamentous cyanobacteria. *Mol. Microbiol.* 98: 1021–1036.
- Killian, D. J., and E. J. Hubbard, 2005 *Caenorhabditis elegans* germline patterning requires coordinated development of the somatic gonadal sheath and the germ line. *Dev. Biol.* 279: 322–335.
- Kim, H. S., R. Murakami, S. Quintin, M. Mori, K. Ohkura *et al.*, 2011 VAB-10 spectraplakins act in cell and nuclear migration in *Caenorhabditis elegans*. *Development* 138: 4013–4023.
- Kim, S., and W. G. Wadsworth, 2000 Positioning of longitudinal nerves in *C. elegans* by nidogen. *Science* 288: 150–154.
- Kimble, J., and D. Hirsh, 1979 The postembryonic cell lineages of the hermaphrodite and male gonads in *Caenorhabditis elegans*. *Dev. Biol.* 70: 396–417.
- Kimble, J. E., and J. G. White, 1981 On the control of germ cell development in *Caenorhabditis elegans*. *Dev. Biol.* 81: 208–219.
- Kiontke, K., A. Barriere, I. Kolotuev, B. Podbilewicz, R. Sommer *et al.*, 2007 Trends, stasis, and drift in the evolution of nematode vulva development. *Curr. Biol.* 17: 1925–1937.
- Klein, R., 2012 Eph/ephrin signalling during development. *Development* 139: 4105–4109.
- Kohrman, A. Q., and D. Q. Matus, 2017 Divide or conquer: cell cycle regulation of invasive behavior. *Trends Cell Biol.* 27: 12–25.
- Kovacevic, I., R. Ho, and E. J. Cram, 2012 CCDC-55 is required for larval development and distal tip cell migration in *Caenorhabditis elegans*. *Mech. Dev.* 128: 548–559.
- Kramer, J. M., 2005 Basement membranes. (September 1, 2005), *WormBook*, ed. The *C. elegans* Research Community, WormBook, doi/10.1895/wormbook.1.16.1, <http://www.wormbook.org>.
- Kramerova, I. A., N. Kawaguchi, L. I. Fessler, R. E. Nelson, Y. Chen *et al.*, 2000 Papilin in development; a pericellular protein with a homology to the ADAMTS metalloproteinases. *Development* 127: 5475–5485.
- Krause, M., M. Park, J. M. Zhang, J. Yuan, B. Harfe *et al.*, 1997 A *C. elegans* E/Daughterless bHLH protein marks neuronal but not striated muscle development. *Development* 124: 2179–2189.
- Kubota, Y., R. Kuroki, and K. Nishiwaki, 2004 A fibulin-1 homolog interacts with an ADAM protease that controls cell migration in *C. elegans*. *Curr. Biol.* 14: 2011–2018.
- Kubota, Y., K. Ohkura, K. K. Tamai, K. Nagata, and K. Nishiwaki, 2008 MIG-17/ADAMTS controls cell migration by recruiting nidogen to the basement membrane in *C. elegans*. *Proc. Natl. Acad. Sci. USA* 105: 20804–20809.
- Kubota, Y., K. Nagata, A. Sugimoto, and K. Nishiwaki, 2012 Tissue architecture in the *Caenorhabditis elegans* gonad depends on interactions among fibulin-1, type IV collagen and the ADAMTS extracellular protease. *Genetics* 190: 1379–1388.
- Kulkarni, G., Z. Xu, A. M. Mohamed, H. Li, X. Tang *et al.*, 2013 Experimental evidence for UNC-6 (netrin) axon guidance by stochastic fluctuations of intracellular UNC-40 (DCC) outgrowth activity. *Biol. Open* 2: 1300–1312.
- Kurosaka, S., and A. Kashina, 2008 Cell biology of embryonic migration. *Birth Defects Res. C Embryo Today* 84: 102–122.
- Lai Wing Sun, K., J. P. Correia, and T. E. Kennedy, 2011 Netrins: versatile extracellular cues with diverse functions. *Development* 138: 2153–2169.
- Lammermann, T., and M. Sixt, 2009 Mechanical modes of ‘amoeboid’ cell migration. *Curr. Opin. Cell Biol.* 21: 636–644.
- Lamouille, S., J. Xu, and R. Derynck, 2014 Molecular mechanisms of epithelial-mesenchymal transition. *Nat. Rev. Mol. Cell Biol.* 15: 178–196.
- Lecuit, T., and F. Pilot, 2003 Developmental control of cell morphogenesis: a focus on membrane growth. *Nat. Cell Biol.* 5: 103–108.
- Lee, M., E. J. Cram, B. Shen, and J. E. Schwarzbauer, 2001 Roles for beta(pat-3) integrins in development and function of *Caenorhabditis elegans* muscles and gonads. *J. Biol. Chem.* 276: 36404–36410.
- Leung-Hagsteeijn, C., A. M. Spence, B. D. Stern, Y. Zhou, M. W. Su *et al.*, 1992 UNC-5, a transmembrane protein with immunoglobulin and thrombospondin type 1 domains, guides cell and pioneer axon migrations in *C. elegans*. *Cell* 71: 289–299.
- Levy-Strumpf, N., 2016 Orchestrating A/P and D/V guidance—a Wnt/Netrin tale. *Worm* 5: e1146857.
- Levy-Strumpf, N., and J. G. Culotti, 2014 Netrins and Wnts function redundantly to regulate antero-posterior and dorso-ventral guidance in *C. elegans*. *PLoS Genet.* 10: e1004381.
- Levy-Strumpf, N., M. Krizus, H. Zheng, L. Brown, and J. G. Culotti, 2015 The Wnt frizzled receptor MOM-5 regulates the UNC-5 netrin receptor through small GTPase-dependent signaling to determine the polarity of migrating cells. *PLoS Genet.* 11: e1005446.
- Linder, S., C. Wiesner, and M. Himmel, 2011 Degrading devices: invadosomes in proteolytic cell invasion. *Annu. Rev. Cell Dev. Biol.* 27: 185–211.
- Lints, R., and D. H. Hall, 2017 *Reproductive System*. *WormAtlas*. doi:10.3908/wormatlas.1.17.

- Llamazares, M., S. Cal, V. Quesada, and C. Lopez-Otin, 2003 Identification and characterization of ADAMTS-20 defines a novel subfamily of metalloproteinases-disintegrins with multiple thrombospondin-1 repeats and a unique GON domain. *J. Biol. Chem.* 278: 13382–13389.
- Lo, T.-W., C. S. Branda, P. Huang, I. E. Sasson, S. J. Goodman *et al.*, 2008 Different isoforms of the *C. elegans* FGF receptor are required for attraction and repulsion of the migrating sex myoblasts. *Dev. Biol.* 318: 268–275.
- Lo, T. W., D. C. Bennett, S. J. Goodman, and M. J. Stern, 2010 *Caenorhabditis elegans* fibroblast growth factor receptor signaling can occur independently of the multi-substrate adaptor FRS2. *Genetics* 185: 537–547.
- Lohmer, L. L., L. C. Kelley, E. J. Hagedorn, and D. R. Sherwood, 2014 Invadopodia and basement membrane invasion in vivo. *Cell Adh. Migr.* 8: 246–255.
- Lohmer, L. L., M. R. Clay, K. M. Naegeli, Q. Chi, J. W. Ziel *et al.*, 2016 A sensitized screen for genes promoting invadopodia function in vivo: CDC-42 and Rab GDI-1 direct distinct aspects of invadopodia formation. *PLoS Genet.* 12: e1005786.
- Lucanic, M., and H. J. Cheng, 2008 A RAC/CDC-42-independent GIT/PIX/PAK signaling pathway mediates cell migration in *C. elegans*. *PLoS Genet.* 4: e1000269.
- Lundquist, E. A., P. W. Reddien, E. Hartwig, H. R. Horvitz, and C. I. Bargmann, 2001 Three *C. elegans* Rac proteins and several alternative Rac regulators control axon guidance, cell migration and apoptotic cell phagocytosis. *Development* 128: 4475–4488.
- Ma, Y., K. Kanakousaki, and L. Buttitta, 2015 How the cell cycle impacts chromatin architecture and influences cell fate. *Front. Genet.* 6: 19.
- Madsen, C. D., and E. Sahai, 2010 Cancer dissemination—lessons from leukocytes. *Dev. Cell* 19: 13–26.
- Martin, K. H., K. E. Hayes, E. L. Walk, A. G. Ammer, S. M. Markwell *et al.*, 2012 Quantitative measurement of invadopodia-mediated extracellular matrix proteolysis in single and multicellular contexts. *J. Vis. Exp.* 66: e4119.
- Matus, D. Q., X. Y. Li, S. Durbin, D. Agarwal, Q. Chi *et al.*, 2010 In vivo identification of regulators of cell invasion across basement membranes. *Sci. Signal.* 3: ra35.
- Matus, D. Q., E. Chang, S. C. Makohon-Moore, M. A. Hagedorn, Q. Chi *et al.*, 2014 Cell division and targeted cell cycle arrest opens and stabilizes basement membrane gaps. *Nat. Commun.* 5: 4184.
- Matus, D. Q., L. L. Lohmer, L. C. Kelley, A. J. Schindler, A. Q. Kohrman *et al.*, 2015 Invasive cell fate requires G1 cell-cycle arrest and histone deacetylase-mediated changes in gene expression. *Dev. Cell* 35: 162–174.
- Mayor, R., and S. Etienne-Manneville, 2016 The front and rear of collective cell migration. *Nat. Rev. Mol. Cell Biol.* 17: 97–109.
- McClatchey, S. T., Z. Wang, L. M. Linden, E. L. Hastie, L. Wang *et al.*, 2016 Boundary cells restrict dystroglycan trafficking to control basement membrane sliding during tissue remodeling. *Elife* 5: e17218.
- Medwig, T. N., and D. Q. Matus, 2017 Breaking down barriers: the evolution of cell invasion. *Curr. Opin. Genet. Dev.* 47: 33–40.
- Meighan, C. M., and J. E. Schwarzbauer, 2007 Control of *C. elegans* hermaphrodite gonad size and shape by vab-3/Pax6-mediated regulation of integrin receptors. *Genes Dev.* 21: 1615–1620.
- Meighan, C. M., A. P. Kann, and E. R. Egress, 2015 Transcription factor hlh-2/E/Daughterless drives expression of alpha integrin ina-1 during DTC migration in *C. elegans*. *Gene* 568: 220–226.
- Milde-Langosch, K., 2005 The Fos family of transcription factors and their role in tumorigenesis. *Eur. J. Cancer* 41: 2449–2461.
- Mi-Mi, L., S. Votra, K. Kempfues, A. Bretscher, and D. Pruyne, 2012 Z-line formins promote contractile lattice growth and maintenance in striated muscles of *C. elegans*. *J. Cell Biol.* 198: 87–102.
- Morrissey, M. A., E. J. Hagedorn, and D. R. Sherwood, 2013 Cell invasion through basement membrane: the netrin receptor DCC guides the way. *Worm* 2: e26169.
- Morrissey, M. A., D. P. Keeley, E. J. Hagedorn, S. T. McClatchey, Q. Chi *et al.*, 2014 B-LINK: a hemiscentin, plakin, and integrin-dependent adhesion system that links tissues by connecting adjacent basement membranes. *Dev. Cell* 31: 319–331.
- Moshfegh, Y., J. J. Bravo-Cordero, V. Miskolci, J. Condeelis, and L. Hodgson, 2014 A Trio-Rac1-Pak1 signalling axis drives invadopodia disassembly. *Nat. Cell Biol.* 16: 574–586.
- Murphy, D. A., and S. A. Courtneidge, 2011 The ‘ins’ and ‘outs’ of podosomes and invadopodia: characteristics, formation and function. *Nat. Rev. Mol. Cell Biol.* 12: 413–426.
- Nishiwaki, K., 1999 Mutations affecting symmetrical migration of distal tip cells in *Caenorhabditis elegans*. *Genetics* 152: 985–997.
- Nishiwaki, K., N. Hisamoto, and K. Matsumoto, 2000 A metalloprotease disintegrin that controls cell migration in *Caenorhabditis elegans*. *Science* 288: 2205–2208.
- Nourshargh, S., and R. Alon, 2014 Leukocyte migration into inflamed tissues. *Immunity* 41: 694–707.
- Oakes, P. W., Y. Beckham, J. Stricker, and M. L. Gardel, 2012 Tension is required but not sufficient for focal adhesion maturation without a stress fiber template. *J. Cell Biol.* 196: 363–374.
- Ohmachi, M., C. E. Rocheleau, D. Church, E. Lambie, T. Schedl *et al.*, 2002 *C. elegans* *ksr-1* and *ksr-2* have both unique and redundant functions and are required for MPK-1 ERK phosphorylation. *Curr. Biol.* 12: 427–433.
- Overall, C. M., and O. Kleinfeld, 2006 Tumour microenvironment - opinion: validating matrix metalloproteinases as drug targets and anti-targets for cancer therapy. *Nat. Rev. Cancer* 6: 227–239.
- Ozanne, B. W., H. J. Spence, L. C. McGarry, and R. F. Hennigan, 2007 Transcription factors control invasion: AP-1 the first among equals. *Oncogene* 26: 1–10.
- Ozbek, S., P. G. Balasubramanian, R. Chiquet-Ehrismann, R. P. Tucker, and J. C. Adams, 2010 The evolution of extracellular matrix. *Mol. Biol. Cell* 21: 4300–4305.
- Padrick, S. B., and M. K. Rosen, 2010 Physical mechanisms of signal integration by WASP family proteins. *Annu. Rev. Biochem.* 79: 707–735.
- Page-McCaw, A., 2008 Remodeling the model organism: matrix metalloproteinase functions in invertebrates. *Semin. Cell Dev. Biol.* 19: 14–23.
- Page-McCaw, A., A. J. Ewald, and Z. Werb, 2007 Matrix metalloproteinases and the regulation of tissue remodelling. *Nat. Rev. Mol. Cell Biol.* 8: 221–233.
- Paluch, E. K., and E. Raz, 2013 The role and regulation of blebs in cell migration. *Curr. Opin. Cell Biol.* 25: 582–590.
- Pan, C. L., J. E. Howell, S. G. Clark, M. Hilliard, S. Cordes *et al.*, 2006 Multiple Wnts and frizzled receptors regulate anteriorly directed cell and growth cone migrations in *Caenorhabditis elegans*. *Dev. Cell* 10: 367–377.
- Paul, C. D., P. Mistriotis, and K. Konstantopoulos, 2017 Cancer cell motility: lessons from migration in confined spaces. *Nat. Rev. Cancer* 17: 131–140.
- Petrie, R. J., and K. M. Yamada, 2012 At the leading edge of three-dimensional cell migration. *J. Cell Sci.* 125: 5917–5926.
- Petrie, R. J., and K. M. Yamada, 2016 Multiple mechanisms of 3D migration: the origins of plasticity. *Curr. Opin. Cell Biol.* 42: 7–12.
- Petrie, R. J., N. Gavara, R. S. Chadwick, and K. M. Yamada, 2012 Nonpolarized signaling reveals two distinct modes of 3D cell migration. *J. Cell Biol.* 197: 439–455.
- Petrie, R. J., H. Koo, and K. M. Yamada, 2014 Generation of compartmentalized pressure by a nuclear piston governs cell motility in a 3D matrix. *Science* 345: 1062–1065.

- Petrie, R. J., H. M. Harlin, L. I. Korsak, and K. M. Yamada, 2017 Activating the nuclear piston mechanism of 3D migration in tumor cells. *J. Cell Biol.* 216: 93–100.
- Photos, A., A. Gutierrez, and R. J. Sommer, 2006 *sem-4/spalt* and *egl-17/FGF* have a conserved role in sex myoblast specification and migration in *P. pacificus* and *C. elegans*. *Dev. Biol.* 293: 142–153.
- Pizarro-Cerda, J., D. S. Chorev, B. Geiger, and P. Cossart, 2017 The diverse family of Arp2/3 complexes. *Trends Cell Biol.* 27: 93–100.
- Polanska, U. M., L. Duchesne, J. C. Harries, D. G. Fernig, and T. K. Kinnunen, 2009 N-Glycosylation regulates fibroblast growth factor receptor/EGL-15 activity in *Caenorhabditis elegans* in vivo. *J. Biol. Chem.* 284: 33030–33039.
- Razidlo, G. L., B. Schroeder, J. Chen, D. D. Billadeau, and M. A. McNiven, 2014 Vav1 as a central regulator of invadopodia assembly. *Curr. Biol.* 24: 86–93.
- Reddien, P. W., and H. R. Horvitz, 2000 CED-2/CrkII and CED-10/Rac control phagocytosis and cell migration in *Caenorhabditis elegans*. *Nat. Cell Biol.* 2: 131–136.
- Reiner, D. J., and E. A. Lundquist, 2016 Small GTPases (January 17, 2006), *WormBook*, ed. The *C. elegans* Research Community, WormBook, doi/10.1895/wormbook.1.67.1, <http://www.wormbook.org>.
- Rella, L., E. E. Fernandes Pova, and H. C. Korswagen, 2016 The *Caenorhabditis elegans* Q neuroblasts: a powerful system to study cell migration at single-cell resolution in vivo. *Genesis* 54: 198–211.
- Renaud, S. J., K. Kubota, M. A. Rumi, and M. J. Soares, 2014 The FOS transcription factor family differentially controls trophoblast migration and invasion. *J. Biol. Chem.* 289: 5025–5039.
- Ridley, A. J., 2015 Rho GTPase signalling in cell migration. *Curr. Opin. Cell Biol.* 36: 103–112.
- Rimann, I., and A. Hajnal, 2007 Regulation of anchor cell invasion and uterine cell fates by the *egl-43* Evi-1 proto-oncogene in *Caenorhabditis elegans*. *Dev. Biol.* 308: 187–195.
- Rougvie, A. E., and V. Ambros, 1995 The heterochronic gene *lin-29* encodes a zinc finger protein that controls a terminal differentiation event in *Caenorhabditis elegans*. *Development* 121: 2491–2500.
- Rowe, R. G., and S. J. Weiss, 2008 Breaching the basement membrane: who, when and how? *Trends Cell Biol.* 18: 560–574.
- Ruijtenberg, S., and S. van den Heuvel, 2015 G1/S inhibitors and the SWI/SNF complex control cell-cycle exit during muscle differentiation. *Cell* 162: 300–313.
- Sasson, I. E., and M. J. Stern, 2004 FGF and PI3 kinase signaling pathways antagonistically modulate sex muscle differentiation in *C. elegans*. *Development* 131: 5381–5392.
- Sawa, H., and H. C. Korswagen, 2013 Wnt signaling in *C. elegans* (December 9, 2013), *WormBook*, ed. The *C. elegans* Research Community, WormBook, doi/10.1895/wormbook.1.7.2, <http://www.wormbook.org>.
- Sawa, M., S. Suetsugu, A. Sugimoto, H. Miki, M. Yamamoto *et al.*, 2003 Essential role of the *C. elegans* Arp2/3 complex in cell migration during ventral enclosure. *J. Cell Sci.* 116: 1505–1518.
- Schindler, A. J., and D. R. Sherwood, 2011 The transcription factor HLH-2/E/Daughterless regulates anchor cell invasion across basement membrane in *C. elegans*. *Dev. Biol.* 357: 380–391.
- Schindler, A. J., and D. R. Sherwood, 2013 Morphogenesis of the *caenorhabditis elegans* vulva. *Wiley Interdiscip. Rev. Dev. Biol.* 2: 75–95.
- Schmidt, K. L., N. Marcus-Gueret, A. Adeleye, J. Webber, D. Baillie *et al.*, 2009 The cell migration molecule UNC-53/NAV2 is linked to the ARP2/3 complex by ABI-1. *Development* 136: 563–574.
- Schonichen, A., and M. Geyer, 2010 Fifteen formins for an actin filament: a molecular view on the regulation of human formins. *Biochim. Biophys. Acta* 1803: 152–163.
- Schoumacher, M., R. D. Goldman, D. Louvard, and D. M. Vignjevic, 2010 Actin, microtubules, and vimentin intermediate filaments cooperate for elongation of invadopodia. *J. Cell Biol.* 189: 541–556.
- Schoumacher, M., D. Louvard, and D. Vignjevic, 2011 Cytoskeleton networks in basement membrane transmigration. *Eur. J. Cell Biol.* 90: 93–99.
- Schwab, A., and C. Stock, 2014 Ion channels and transporters in tumour cell migration and invasion. *Philos. Trans. R. Soc. Lond. B Biol. Sci.* 369: 20130102.
- Seano, G., G. Chiaverina, P. A. Gagliardi, L. di Blasio, A. Puliafito *et al.*, 2014 Endothelial podosome rosettes regulate vascular branching in tumour angiogenesis. *Nat. Cell Biol.* 16: 931–941, 1–8.
- Sgambato, A., M. Migaldi, M. Montanari, A. Camerini, A. Brancaccio *et al.*, 2003 Dystroglycan expression is frequently reduced in human breast and colon cancers and is associated with tumor progression. *Am. J. Pathol.* 162: 849–860.
- Shafaq-Zadah, M., C. S. Gomes-Santos, S. Bardin, P. Maiuri, M. Maurin *et al.*, 2016 Persistent cell migration and adhesion rely on retrograde transport of beta(1) integrin. *Nat. Cell Biol.* 18: 54–64.
- Shakir, M. A., K. Jiang, E. C. Struckhoff, R. S. Demarco, F. B. Patel *et al.*, 2008 The Arp2/3 activators WAVE and WASP have distinct genetic interactions with Rac GTPases in *Caenorhabditis elegans* axon guidance. *Genetics* 179: 1957–1971.
- Shen, Z., X. Zhang, Y. Chai, Z. Zhu, P. Yi *et al.*, 2014 Conditional knockouts generated by engineered CRISPR-Cas9 endonuclease reveal the roles of coronin in *C. elegans* neural development. *Dev. Cell* 30: 625–636.
- Sherwood, D. R., and P. W. Sternberg, 2003 Anchor cell invasion into the vulval epithelium in *C. elegans*. *Dev. Cell* 5: 21–31.
- Sherwood, D. R., J. A. Butler, J. M. Kramer, and P. W. Sternberg, 2005 FOS-1 promotes basement-membrane removal during anchor-cell invasion in *C. elegans*. *Cell* 121: 951–962.
- Silhankova, M., and H. C. Korswagen, 2007 Migration of neuronal cells along the anterior-posterior body axis of *C. elegans*: Wnts are in control. *Curr. Opin. Genet. Dev.* 17: 320–325.
- Sobrado, V. R., G. Moreno-Bueno, E. Cubillo, L. J. Holt, M. A. Nieto *et al.*, 2009 The class I bHLH factors E2-2A and E2-2B regulate EMT. *J. Cell Sci.* 122: 1014–1024.
- Somerville, R. P., J. M. Longpre, K. A. Jungers, J. M. Engle, M. Ross *et al.*, 2003 Characterization of ADAMTS-9 and ADAMTS-20 as a distinct ADAMTS subfamily related to *Caenorhabditis elegans* GON-1. *J. Biol. Chem.* 278: 9503–9513.
- Soto, M. C., H. Qadota, K. Kasuya, M. Inoue, D. Tsuboi *et al.*, 2002 The GEX-2 and GEX-3 proteins are required for tissue morphogenesis and cell migrations in *C. elegans*. *Genes Dev.* 16: 620–632.
- Spencer, W. C., R. McWhirter, T. Miller, P. Strasbourger, O. Thompson *et al.*, 2014 Isolation of specific neurons from *C. elegans* larvae for gene expression profiling. *PLoS One* 9: e112102.
- Srivastava, A., J. C. Pastor-Pareja, T. Igaki, R. Pagliarini, and T. Xu, 2007 Basement membrane remodeling is essential for *Drosophila* disc eversion and tumor invasion. *Proc. Natl. Acad. Sci. USA* 104: 2721–2726.
- Stern, M. J., and H. R. Horvitz, 1991 A normally attractive cell interaction is repulsive in two *C. elegans* mesodermal cell migration mutants. *Development* 113: 797–803.
- Sternberg, P. W., 2005 Vulval development (June 25, 2005), *WormBook*, ed. The *C. elegans* Research Community, WormBook, doi/10.1895/wormbook.1.6.1, <http://www.wormbook.org>.
- Stetak, A., E. Gutierrez, and A. Hajnal, 2008 Tissue-specific functions of the *Caenorhabditis elegans* p120 Ras GTPase activating protein GAP-3. *Dev. Biol.* 323: 166–176.
- Stringham, E., N. Pujol, J. Vandekerckhove, and T. Bogaert, 2002 *unc-53* controls longitudinal migration in *C. elegans*. *Development* 129: 3367–3379.
- Stroka, K. M., H. Jiang, S. H. Chen, Z. Tong, D. Wirtz *et al.*, 2014 Water permeation drives tumor cell migration in confined microenvironments. *Cell* 157: 611–623.
- Su, M., D. C. Merz, M. T. Killeen, Y. Zhou, H. Zheng *et al.*, 2000 Regulation of the UNC-5 netrin receptor initiates the first

- reorientation of migrating distal tip cells in *Caenorhabditis elegans*. *Development* 127: 585–594.
- Sulston, J. E., and H. R. Horvitz, 1977 Post-embryonic cell lineages of the nematode, *Caenorhabditis elegans*. *Dev. Biol.* 56: 110–156.
- Sundaram, M., J. Yochem, and M. Han, 1996 A Ras-mediated signal transduction pathway is involved in the control of sex myoblast migration in *Caenorhabditis elegans*. *Development* 122: 2823–2833.
- Sundaram, M. V., 2013 Canonical RTK-Ras-ERK signaling and related alternative pathways (July 1, 2013), *WormBook*, ed. The *C. elegans* Research Community, *WormBook*, doi/10.1895/wormbook.1.80.2, //www.wormbook.org.
- Suozi, K. C., X. Wu, and E. Fuchs, 2012 Spectraplakins: master orchestrators of cytoskeletal dynamics. *J. Cell Biol.* 197: 465–475.
- Suzuki, N., H. Toyoda, M. Sano, and K. Nishiwaki, 2006 Chondroitin acts in the guidance of gonadal distal tip cells in *C. elegans*. *Dev. Biol.* 300: 635–646.
- Tamai, K. K., and K. Nishiwaki, 2007 bHLH transcription factors regulate organ morphogenesis via activation of an ADAMTS protease in *C. elegans*. *Dev. Biol.* 308: 562–571.
- Tang, X., and W. G. Wadsworth, 2014 SAX-3 (Robo) and UNC-40 (DCC) regulate a directional bias for axon guidance in response to multiple extracellular cues. *PLoS One* 9: e110031.
- Tannoury, H., V. Rodriguez, I. Kovacevic, M. Ibourk, M. Lee *et al.*, 2010 CACN-1/Cactin interacts genetically with MIG-2 GTPase signaling to control distal tip cell migration in *C. elegans*. *Dev. Biol.* 341: 176–185.
- Te Boekhorst, V., and P. Friedl, 2016 Plasticity of cancer cell invasion-mechanisms and implications for therapy. *Adv. Cancer Res.* 132: 209–264.
- Te Boekhorst, V., L. Preziosi, and P. Friedl, 2016 Plasticity of cell migration in vivo and in silico. *Annu. Rev. Cell Dev. Biol.* 32: 491–526.
- Tennessen, J. M., H. F. Gardner, M. L. Volk, and A. E. Rougvie, 2006 Novel heterochronic functions of the *Caenorhabditis elegans* period-related protein LIN-42. *Dev. Biol.* 289: 30–43.
- Thomas, J. H., M. J. Stern, and H. R. Horvitz, 1990 Cell interactions coordinate the development of the *C. elegans* egg-laying system. *Cell* 62: 1041–1052.
- Topf, U., and R. Chiquet-Ehrismann, 2011 Genetic interaction between *Caenorhabditis elegans* teneurin ten-1 and prolyl 4-hydroxylase phy-1 and their function in collagen IV-mediated basement membrane integrity during late elongation of the embryo. *Mol. Biol. Cell* 22: 3331–3343.
- Tripathi, A., A. H. Nile, and V. A. Bankaitis, 2014 Sec14-like phosphatidylinositol-transfer proteins and diversification of phosphoinositide signalling outcomes. *Biochem. Soc. Trans.* 42: 1383–1388.
- Trzebiatowska, A., U. Topf, U. Sauder, K. Drabikowski, and R. Chiquet-Ehrismann, 2008 *Caenorhabditis elegans* teneurin, ten-1, is required for gonadal and pharyngeal basement membrane integrity and acts redundantly with integrin ina-1 and dystroglycan dgn-1. *Mol. Biol. Cell* 19: 3898–3908.
- Valastyan, S., and R. A. Weinberg, 2011 Tumor metastasis: molecular insights and evolving paradigms. *Cell* 147: 275–292.
- Verghese, E., J. Schocken, S. Jacob, A. M. Wimer, R. Royce *et al.*, 2011 The tailless ortholog nhr-67 functions in the development of the *C. elegans* ventral uterus. *Dev. Biol.* 356: 516–528.
- Vicente-Manzanares, M., X. Ma, R. S. Adelstein, and A. R. Horvitz, 2009 Non-muscle myosin II takes centre stage in cell adhesion and migration. *Nat. Rev. Mol. Cell Biol.* 10: 778–790.
- Vogel, B. E., and E. M. Hedgecock, 2001 Hemicentin, a conserved extracellular member of the immunoglobulin superfamily, organizes epithelial and other cell attachments into oriented line-shaped junctions. *Development* 128: 883–894.
- Vuong-Brender, T. T. K., X. Yang, and M. Labouesse, 2016 *C. elegans* embryonic morphogenesis. *Curr. Top. Dev. Biol.* 116: 597–616.
- Walck-Shannon, E., D. J. Reiner, and J. Hardin, 2015 Polarized Rac-dependent protrusions drive epithelial intercalation in the embryonic epidermis of *C. elegans*. *Development* 142: 3549–3560.
- Wang, L., W. Shen, S. Lei, D. Matus, D. Sherwood *et al.*, 2014 MIG-10 (Lamellipodin) stabilizes invading cell adhesion to basement membrane and is a negative transcriptional target of EGL-43 in *C. elegans*. *Biochem. Biophys. Res. Commun.* 452: 328–333.
- Wang, Z., Q. Chi, and D. R. Sherwood, 2014a MIG-10 (lamellipodin) has netrin-independent functions and is a FOS-1A transcriptional target during anchor cell invasion in *C. elegans*. *Development* 141: 1342–1353.
- Wang, Z., L. M. Linden, K. M. Naegeli, J. W. Ziel, Q. Chi *et al.*, 2014b UNC-6 (netrin) stabilizes oscillatory clustering of the UNC-40 (DCC) receptor to orient polarity. *J. Cell Biol.* 206: 619–633.
- Washington, R. W., and D. A. Knecht, 2008 Actin binding domains direct actin-binding proteins to different cytoskeletal locations. *BMC Cell Biol.* 9: 10.
- Wei, X., C. J. Potter, L. Luo, and K. Shen, 2012 Controlling gene expression with the Q repressible binary expression system in *Caenorhabditis elegans*. *Nat. Methods* 9: 391–395.
- Whangbo, J., and C. Kenyon, 1999 A Wnt signaling system that specifies two patterns of cell migration in *C. elegans*. *Mol. Cell* 4: 851–858.
- Wickstrom, S. A., and R. Fassler, 2011 Regulation of membrane traffic by integrin signaling. *Trends Cell Biol.* 21: 266–273.
- Williams, J. M., and C. W. Daniel, 1983 Mammary ductal elongation: differentiation of myoepithelium and basal lamina during branching morphogenesis. *Dev. Biol.* 97: 274–290.
- Wong, M. C., and J. E. Schwarzbauer, 2012 Gonad morphogenesis and distal tip cell migration in the *Caenorhabditis elegans* hermaphrodite. *Wiley Interdiscip. Rev. Dev. Biol.* 1: 519–531.
- Wong, M. C., W. P. Kennedy, and J. E. Schwarzbauer, 2014 Transcriptionally regulated cell adhesion network dictates distal tip cell directionality. *Dev. Dyn.* 243: 999–1010.
- Xu, X., S. C. Rongali, J. P. Miles, K. D. Lee, and M. Lee, 2006 pat-4/ILK and unc-112/Mig-2 are required for gonad function in *Caenorhabditis elegans*. *Exp. Cell Res.* 312: 1475–1483.
- Xu, X., C. Dong, and B. E. Vogel, 2007 Hemicentins assemble on diverse epithelia in the mouse. *J. Histochem. Cytochem.* 55: 119–126.
- Yan, B., Q. Guo, F. J. Fu, Z. Wang, Z. Yin *et al.*, 2015 The role of miR-29b in cancer: regulation, function, and signaling. *Oncotargets Ther.* 8: 539–548.
- Yang, J., and R. A. Weinberg, 2008 Epithelial-mesenchymal transition: at the crossroads of development and tumor metastasis. *Dev. Cell* 14: 818–829.
- Yang, Q., D. Roiz, L. Mereu, M. Daube, and A. Hajnal, 2017 The invading anchor cell induces lateral membrane constriction during vulval lumen morphogenesis in *C. elegans*. *Dev. Cell* 42: 271–285.e3.
- Yano, S., Y. Zhang, S. Miwa, Y. Tome, Y. Hiroshima *et al.*, 2014 Spatial-temporal Fucci imaging of each cell in a tumor demonstrates locational dependence of cell cycle dynamics and chemoresponsiveness. *Cell Cycle* 13: 2110–2119.
- Yoo, A. S., C. Bais, and I. Greenwald, 2004 Crosstalk between the EGFR and LIN-12/Notch pathways in *C. elegans* vulval development. *Science* 303: 663–666.
- Yurchenco, P. D., 2011 Basement membranes: cell scaffoldings and signaling platforms. *Cold Spring Harb. Perspect. Biol.* 3: a004911.
- Zhang, L., J. D. Ward, Z. Cheng, and A. F. Dernburg, 2015 The auxin-inducible degradation (AID) system enables versatile conditional protein depletion in *C. elegans*. *Development* 142: 4374–4384.
- Ziel, J. W., E. J. Hagedorn, A. Audhya, and D. R. Sherwood, 2009 UNC-6 (netrin) orients the invasive membrane of the anchor cell in *C. elegans*. *Nat. Cell Biol.* 11: 183–189.
- Zipkin, I. D., R. M. Kindt, and C. J. Kenyon, 1997 Role of a new Rho family member in cell migration and axon guidance in *C. elegans*. *Cell* 90: 883–894.
- Zucker, S., J. Cao, and W. T. Chen, 2000 Critical appraisal of the use of matrix metalloproteinase inhibitors in cancer treatment. *Oncogene* 19: 6642–6650.

Communicating editor: M. Labouesse