

Are Behavioral Change Interventions Needed to Make Cash Transfer Programs Work for Children? Experimental Evidence from Myanmar *

Erica Field[†] Elisa M. Maffioli[‡]

September 6, 2023

Abstract

We experimentally evaluate the impact on child malnutrition of a maternal cash transfer program in Myanmar that was supplemented with Social Behavior Change Communication (SBCC) in a subset of villages. The combination of interventions significantly reduced the proportion of children stunted, while cash alone had no impact on stunting. SBCC appears to have worked in conjunction with cash to reduce stunting by encouraging mothers to increase children's diet diversity, in particular proteins consumed. The findings underscore the importance of adding SBCC to cash transfer programs.

Keywords: Nutrition, Cash Transfers, Social and Behavior Change Communication

**Acknowledgments:* We thank the Livelihoods and Food Security Fund and the International Growth Centre for their support. We also thank Save the Children for their guidance, feedback, and faith in the research process. This project would not have been possible without the contributions of the Ministry of Social Welfare, Relief, and Resettlement; the Department of Social Welfare; the Myanmar Nurse and Midwife Association; and the Pact Global Microfinance Fund. Finally, we extend our deepest gratitude to the hardworking staff at Innovations for Poverty Action: Nicholas Tint Zaw, Thein Zaw Oo, and Than Zaw Oo. The trial was registered on AEA RCT registry (AEARCTR-0004189). Ethical approval was obtained by Duke University; Department of Medical Research, Government of the Republic of the Union of Myanmar, and Innovations for Poverty Action.

[†]Department of Economics, Duke University, Durham, NC 27710 (e-mail: field.eric@duke.edu)

[‡]Health Management and Policy, University of Michigan School of Public Health, Ann Arbor, MI 48109 (e-mail: elisamaf@umich.edu)

1 Introduction

Despite widescale improvements in economic wellbeing over the past decades, child malnutrition remains a global health concern, affecting more than 150 million children annually (World Bank, 2017). When families cannot afford to provide children with sufficient nutrient-rich food, malnutrition can lead to irreversible decreases in health and cognitive human capital. In Myanmar, where this study takes place, an estimated 29 percent of the children under age five are stunted (Demographic Health Surveys, 2015; United Nation Standing Committee on Nutrition, 2010).

Medical and social science research contributed to the consensus that the first 1000 days of life constitute the most critical period of development (Almond and Currie, 2011a), and inadequate nutrition produce higher susceptibility to illness, impaired physical and cognitive ability (Victora et al., 2010; Almond and Currie, 2011b; Doyle, 2019), and lower human capital accumulation, productivity and earnings in adult life (Engle et al., 2007; Hoddinott et al., 2013; Richter et al., 2017). This has motivated governments, NGOs, and international agencies to direct cash transfers programs to households with young children, which to date have reached 1.3 billion people globally (Bastagli et al., 2016; Hoddinott et al., 2017; Gentilini, 2022).

There are two principal ways in which cash transfers may directly reduce early life malnutrition. First, households at risk of malnourishment could spend a large fraction of income on child nutrition simply because investing in children during critical windows of physical and cognitive development is one of the highest return investments a family can make. Likewise, cash transfers could also facilitate similarly high-return investments in child health production such as health care expenditures. Second, these programs are frequently directed to mothers (“maternal cash transfer programs”), and increasing women’s control over income is believed to promote investment in children in and of itself.¹

However, it is not obvious that simply giving cash is sufficient to make a dent in

¹There is a substantial body of literature linking female income share to increases in children’s expenditure. Bobonis (2009) finds that climate shocks that contribute to female bargaining power, increased household expenditures on children’s goods (education, health). Similarly, Duflo and Udry (2004) find that positive rainfall shocks to women-controlled crops in Cote D’Ivoire increased shares of education expenditures for children. Lépine and Strobl (2013) find that positive rainfall shocks to women-controlled crops in Ghana increased children’s nutritional status. Duflo (2003) find that providing pensions to South African grandmothers increased height-for-age of grandchildren. There are several theories explaining this bias including gender differences in old age security, preferences, and altruism.

early life malnourishment. In fact, there is no consensus from evaluations of existing programs that cash transfers generate significant reductions in stunting. While cash transfer programs have been associated with a wide range of positive household outcomes (Bastagli et al., 2016), the evidence on child nutrition is limited (Baird et al., 2019; Ahmed et al., 2019). While some program evaluations have shown positive impacts, several others reported mixed or null results (Ritcher, 2010; Soares et al., 2010; Cecchini and Madariaga, 2011; Manley et al., 2013; Bastagli et al., 2016; Tirivayia et al., 2016; Biscaye et al., 2017; Millan et al., 2019; Manley et al., 2020, 2022),.

Maternal cash transfer programs may fail to reduce malnutrition for two reasons. First, households may lack sufficient information on child health production to understand the value of early life nutrition, including the importance of a more diverse diet, reducing the marginal impact of income on child health. Second, households may lack financial resources and/or fail to direct consumption towards child health if those who control income do not fully internalize welfare and hence prefer an alternative consumption bundle.²

To increase the impact on child malnutrition, policymakers increasingly enhance maternal cash transfer programs with complementary features designed to address one or both constraints. In many settings, programs include conditionalities such as mandatory health visits in order to incentivize households who may otherwise prefer to put cash elsewhere to invest in child health. While a number of studies show positive effects of conditional cash transfers (CCTs) on stunting, there is ongoing concern that imposing conditionalities excludes the most vulnerable households from receiving program benefits (Cahyadi et al., 2020; Kandpal et al., 2016). In-kind transfer programs can mitigate both constraints, but are costly to implement and may be difficult to tailor to households' idiosyncratic dietary needs (Jeong and Trako, 2022).

An alternative strategy for promoting child health that is frequently implemented in conjunction with maternal cash transfers are information programs delivered via Social and Behavior Change Communication (SBCC). SBCC sessions attached to maternal cash transfer programs are designed to tackle the information constraint by supplementing cash transfers with information on child health production, and their curricula typically focus on infant and child health and feeding practices. Participation

²Maternal cash transfer programs may also fail to reduce malnutrition if there are important constraints to local food availability, such as occurs during famines or in food deserts. This does not apply in the context of our evaluation, as our sample villages lie within two hours of a major city and no severe economic shocks happened during the study period.

in SBCC programs is voluntary, and hence immune to concerns over systematic exclusion. However, it is unclear whether information frictions impose a binding constraint on the marginal propensity to invest in child health out of cash transfer income, and also whether relevant information on child health production can be effectively delivered at scale.

In empirical work to date, the efficacy of supplementing maternal cash transfers with SBCC in bolstering child health remains uncertain.³ As cash transfers are the starting point for governments, NGOs, and international agencies to fight poverty (Gentilini, 2022), understanding the potential value of supplementing cash transfers with SBCC for child malnutrition is policy-relevant, especially in a context such as in Myanmar where anti-poverty programs are sparse and there is a need to fill the nutrition gap (World Food Program, 2019).

A number of studies have evaluated child health impacts of maternal SBCC programs, but evidence is often inconclusive and limited to behavior change outcomes (Luo et al., 2012; Bhutta et al., 2013; Avula et al., 2013; Hoddinott et al., 2017). An exception is an experimental evaluation of a child nutrition SBCC program in Malawi, which was associated with gains in HAZ (0.27 SD) (Fitzsimons et al., 2016). Yet, cash transfer programs can lead to improved infant and child feeding practices if they are combined with SBCC (Fiszbein et al., 2009), but evidence of the impacts on child health outcomes is scarce. A recent randomized experiment in Nepal, which explored the effect of cash transfers plus SBCC relative to SBCC only, found meaningful effects of supplementing cash transfers with SBCC on health knowledge and behavior, but null effects on child malnutrition (Leveré et al., 2022). A contemporaneous study in Nigeria finds meaningful impacts of cash transfers plus SBCC on child stunting, but is unable to isolate the marginal contribution of SBCC because the research does not evaluate the impact of cash alone (Carneiro et al., 2021). The closest evidence to ours, which corroborates our results, is from a similar RCT that was conducted in Bangladesh at the same time as our study, and finds similar evidence that cash transfers plus SBCC reduces stunting in children relative to both cash alone and control (Ahmed et al., 2019).

We add to the existing literature by investigating the impact of additional interventions (i.e., SBCC) on top of cash transfers relative to cash transfers alone, in the

³There is huge evidence on conditional and unconditional cash transfers aimed at improving health (Fiszbein et al., 2009), but our paper is primarily related and contributes specifically to the literature on unconditional cash transfers for nutrition.

context of Myanmar. Specifically, we evaluate a combination of interventions designed to reduce chronic malnutrition during the first 1000 days of life by providing cash transfers with and without SBCC to women who are pregnant or have children under age two. The program, sponsored by the Government of Myanmar and implemented by Save the Children International (SCI), was run for 30 months in 416 rural villages as a pilot for the government’s national maternal transfer program.⁴ The program targeted all pregnant women in intervention villages, who were provided monthly cash transfers from enrollment until their child reached age two (i.e. for 24-30 months). In a randomly chosen subset of treatment villages, program recipients also received monthly SBCC group sessions for the duration of the program that covered a range of topics relevant to child health and nutrition.

We assess the impact after 30 months of implementation of transfers alone and in combination with SBCC on child height-for-age Z-scores (HAZ scores) and stunting, a well-validated biometric measure of chronic malnutrition in children (Leroy and Frongillo, 2019). We restrict our analysis to women found to be pregnant immediately prior to program launch, which allows us to gauge the impact of receiving the full duration of program transfers while also circumventing concerns over selective fertility or migration into treatment villages.

Our results indicate that the combination of cash transfers and SBCC leads to a 4.6 percentage point (13.5%) statistically significant reduction in the proportion of children who are stunted. The program appears to be effective for children at risk of moderate but not severe stunting, which indicates that more heavy-handed approaches or higher levels of transfers might be required to address malnutrition among the most vulnerable children. Meanwhile, cash alone has no detectable impact on child anthropometrics relative to the control group.

Survey data on health behaviors collected at endline indicate that the cash transfers, when combined with SBCC, reduced stunting through some combination of improvements in total food consumption, dietary diversity, breastfeeding and hand-washing practices, all of which are reported to be significantly higher among those treated with the combined interventions relative to both control and cash only groups. Most notably, relative to the control group, food consumption in the combined treatment group rises by 15%, accompanied by a significant improvement in a standardized index of child dietary diversity. Both amount and type of food like protein can directly reduce chronic

⁴The name of the program was LEGACY, which stands for “Learning, Evidence Generation, and Advocacy for Catalyzing Policy”.

malnutrition by increasing energy availability, while higher rates of breastfeeding and hand-washing lower stunting by reducing nutrient-depleting episodes of diarrheal disease. In this sense, disease environment plays an important role in determining baseline levels of nutrient absorption and therefore stunting.

We evaluate these competing mechanisms by examining survey data on child illness episodes, health care expenditures and **DROP-food diary reports of** specific foods consumed, all of which point towards dietary diversity being the critical behavior change. First, we find no decrease in child illness episodes or health care spending among children in the cash plus SBCC group, which indicates limited roles of hand-washing and breastfeeding behaviors in reducing child stunting. Second, **survey data** reveal that transfer recipients who were also exposed to the SBCC curriculum incorporate significantly more protein-rich foods into children’s diets, including meat, pulses, dairy, and eggs. While fruits and vegetables can improve child nutrition, animal proteins in particular have been shown to have a significant impact on child stunting in multiple settings.⁵ In sum, the weight of evidence indicates that most of the reductions in stunting observed among children whose mothers received cash transfers alongside the SBCC program arise from improvements in dietary diversity.

Meanwhile, cash transfers both with and without SBCC improved reported take-up of prenatal care and lead to higher levels of food consumption. However, the similar rates of stunting between the cash only and control groups indicate that these behavior modifications were insufficient to influence chronic malnutrition. While prenatal care is unlikely to have a significant effect on stunting in any setting, the absence of a stunting effect on the cash only arm is more surprising. However, not only is the increase in food consumption among the cash only group relative to control significantly lower than observed in SBCC villages (7%, $p < 0.01$), but there is no significant change in child dietary diversity, both of which could account for a null result on stunting.

These findings provide novel evidence on the policy importance of combining maternal cash transfers with behavioral change interventions in order to generate meaningful improvements in child nutrition. By targeting both financial and information constraints, this study offers a fundamental lesson for the design and implementation of maternal cash transfer programs in low-income countries. First and foremost, in settings such as rural Myanmar where child malnutrition remains a significant problem,

⁵See [Laplante and Sabatini \(2012\)](#) and [Semba et al. \(2016a,b\)](#) for meat and fish; [Molgaard et al. \(2011\)](#); [Iannotti et al. \(2013\)](#); [Dyer et al. \(2016\)](#) for dairy products; [Semba et al. \(2016c\)](#); [Bekdash \(2016\)](#) for eggs.

policies that increase household income still have large potential to improve child malnutrition as long as mothers are also provided with adequate knowledge to purchase the appropriate quantity and quality of foods. Moreover, given that the improvements in stunting appear to be driven largely by promoting changes in dietary diversity, lessons on infant and child feeding practices should be heavily emphasized in SBCC curriculum and prioritized whenever programs are streamlined for scale-up. While we cannot rule out the possibility that information alone would have had comparable impacts on malnutrition, our results indicate that the SBCC curriculum reduced malnutrition by convincing households to increase food consumption rather than by convincing them to change health practices such as infant feeding and hand-washing, which implies that liquidity constraints potentially bind the behavioral responses to the program. Hence, it is unlikely that SBCC delivered *without* cash transfers would have achieved as large an impact on child stunting.

Our findings also offer a key lesson for the evaluation of maternal health programs. In particular, both cash transfers and SBCC have the potential to generate meaningful changes in parental health behaviors *without* significant reductions in child malnourishment. As a result, tracking child health outcomes and not only behavioral outcomes is critical for comprehensive evaluation of program effects on children.

2 Methods

2.1 Setting

In Myanmar, close to 1 out of 3 children are chronically malnourished. To address concerns over child malnutrition, in 2014 the government of Myanmar committed to rolling out a national maternal cash transfer program, which is projected to reach 2.25 million beneficiaries and 0.32% of GDP by 2024 ([The Republic of The Union of Myanmar, 2014](#)). SCI was chosen to implement a pilot version of the program as a randomized controlled trial (RCT) for 30 months prior to national scale-up in order to test the delivery model, including the inclusion of a maternal behavior change component, and measure impacts on malnutrition.⁶

The pilot was implemented between 2016 and 2019 in 416 villages in three town-

⁶Since 2018 the program has been extended at scale in several states and (in 2020-2021) to further support vulnerable households during the COVID-19 pandemic ([Livelihoods and Food Security Fund, 2020](#)).

ships of Myanmar’s Dry Zone – Pakkoku, Yesagyo, and Mahlaing. All villages within two hours of an urban center were eligible for the study. Overall, study villages have reasonable access to food markets (96% have a food market located in their community), but relatively poor access to medical care: only 18% have a village health facility, and only 22% have a midwife that visits regularly. The majority of households in this area earn income from agriculture (89%) and livestock (27%), and casual labor (77%) (Appendix Table A1 presents these statistics by treatment and control groups).

Malnutrition in this area is representative of the country as a whole. Baseline data collected prior to the intervention reveal that 28.7% of children under 5 in the study villages were stunted, almost identical to the national rate of under 5 stunting of 29% in 2015 (Demographic Health Surveys, 2015) (Appendix Table A3).⁷ In terms of weight-for-height, 18% were wasted and in terms of weight-for age, 31% were underweight.⁸

In terms of parental health behaviors that contribute to child malnourishment and hence are generally included in SBCC curricula, baseline data reveal that households in this setting performed very well on some measures and relatively poorly on others. Breastfeeding is nearly universal, as is early initiation of breastfeeding. Likewise, households do fairly well at baseline in terms of WASH behaviors. Almost every woman reported using soap when washing their hands (99%), and the vast majority reported doing so consistently after going to the toilet (77%), the most critical routine WASH behavior for disease control. In contrast, nutritional intake of children over 6 months is poor in this setting. Only 37% of children 6-23 months have a minimum acceptable diet in terms of food diversity. Moreover, there appears to be room for information interventions to have an impact on complementary feeding practices and to influence health-seeking behavior.

In addition to addressing information constraints on child feeding and health care practices, the cash transfer alone has the potential to lead to improvements in child diet and take-up of health care services by relaxing household budget constraints. However, baseline data from household with stunted children do not seem to be living hand-to-mouth, as they spent only 54% of their budget on food, and reported spending their remaining income on a number of “non-essentials” including an average of 7% on gifts

⁷Note that since baseline data were collected after the program was launched (June-September 2016) due to implementation delays, we do not use baseline data in the main analysis. However, we match our analysis sample with baseline data (55.8%) and use this sub-sample of women and their children to provide some background information on the setting.

⁸The rate at baseline is somewhat higher among children 22-35 months old, the age group used for our endline analysis. In the age group, 30% were stunted, 16% were wasted and 33% under-weight.

and donations.

2.2 Program Design

The program comprised two interventions: 1) monthly cash transfers to mothers beginning in pregnancy until their children turned two; and 2) monthly cash transfers supplemented with monthly SBCC that covered a range of topics relevant to child health and nutrition.⁹

Both interventions were randomized across 102 sub-rural health care center catchment areas (the geographic unit of randomization) located within two hours of an urban center. To minimize differences across experimental arms, prior to random assignment, catchment areas were first grouped into 34 triplets (strata) based on geographic clustering. Within each stratum, individual catchment areas were randomly assigned to one of three experimental arms: (1) Treatment 1 (*Cash + SBCC*), in which cash transfers and SBCC activities were provided jointly (N = 34 catchment areas encompassing 142 villages); (2) Treatment 2 (*CashOnly*), in which only cash transfers were provided (N= 34 catchment areas encompassing 146 villages); and (3) Control, in which neither cash transfers nor SBCC were offered (N = 34 catchment areas encompassing 149 villages) (Appendix Fig A1).

Within both T1 and T2 catchment areas, all pregnant women were assigned to receive monthly cash transfers worth 10,000 MMK (about 6.5 USD) beginning in their second trimester of pregnancy until their child reached age two.¹⁰ As a reference, the legal minimum wage in Myanmar at that time was 3,600 MMK per day, so the cash transfer amount represented about 3-4 days of work at the minimum wage. In addition to monthly cash transfers, beneficiaries in T1 were targeted with SBCC in the form of monthly information sessions on four main topics: infant and young child feeding (IYCF) practices, health-seeking behavior, hygiene practices, and household expenditures.

The program was implemented by SCI in collaboration with the Myanmar Nurse and Midwives Association (MNMA), a national non-governmental organization that

⁹Although this was not a labeled cash transfer, per se, cash and SBCC interventions were likely linked in the sense that they were delivered at the same time by support staff from the same institution (SCI).

¹⁰In October 2017 the implementer (SCI) increased the amount to 15,000 MMK (about 10 USD) to stay in line with similar initiatives in other parts of the country. The value was estimated to cover enough nutritious food for a mother and a child for a month.

provides prevention and community-based care, and Pact Global Microfinance (PGMF), a nonprofit international development organization that delivers microfinance in rural areas. PGMF managed monthly cash disbursements by creating an ad-hoc bank account for each program beneficiary into which transfers were deposited on a monthly basis and delivered through PGMF’s network of rural loan agents. MNMA was responsible for coordinating the sensitization and enrollment of eligible women in each treatment village and organizing SBCC activities in villages assigned to the T1 group.¹¹

SBCC activities were implemented in two stages. First, between May 2016 and January 2017, MNMA delivered basic SBCC programming within each village aimed at mobilizing communities to address poor nutrition. Basic programming included mother-to-mother support groups (including 12-15 pregnant women or mothers of under 5 years old children) in which mothers were brought together monthly to disseminate information and share experiences with feeding practices during pregnancy, lactation, and early childhood; and a handful of participatory community-level sessions (13-15 community members) that explored perceptions and current practices around diet and nutrition, health care, and household and food expenditures.

Based on the information gathered through the basic SBCC activities, SCI then designed a series of intensive SBCC sessions that focused on key behaviors and messaging across four topics: IYCF (including dietary diversity and breastfeeding), health-seeking behavior, hygiene practices, and household expenditures. These sessions were delivered both to the maternal support groups, and also through separate sessions targeted to fathers and elderly household members. The last cohort of mothers was enrolled in May 2018. The last monthly cash transfer and the last SBCC interventions were completed in November 2018 (allowing for at least 6 months of cash transfers for the last cohort enrolled) and May 2019, respectively. Appendix Figure A2 describes the timeline of survey and program activities.¹²

SBCC participation was voluntary. While all mothers in SBCC villages were encouraged to attend the sessions, they still received the full transfer if they were unable or refused to attend. Nevertheless, administrative data indicate high participation in

¹¹One aspect of delivery the government was interested in testing in this pilot program was utilizing a NGO for distribution of cash payments versus government workers. Hence, in 40 villages, payments were delivered by government workers instead of PGMF. A description of the difference in delivery agents and findings from that evaluation are the subject of a companion paper (Field and Maffioli, 2021).

¹²Note that the analysis sample for this evaluation only includes women found to be pregnant at the time of program announcement, i.e., the first enrolled cohort covered by the program for about 30 months.

SBCC sessions: administrative data from SCI found that in *Cash+SBCC* villages, 99% of enrolled mothers attended at least one SBCC session and 81% attended five times or more.¹³ Unsurprisingly, take-up of the cash transfer was also high and relatively “clean” in terms of eligibility criteria: monitoring activities conducted independently by the research team 30 months into implementation in one of three townships revealed low exclusion and inclusion errors to the cash transfer programs (6.8% and 9.8%, respectively), and all inclusion errors were women in treatment villages who received transfers despite not meeting the eligibility criteria rather than non-compliers from outside villages, reducing concern over contamination of the control group.¹⁴

2.3 Analysis Sample

We evaluate the program’s impact on child nutrition among women who were pregnant at the time of the program launch. Restricting the sample to this group mitigates concern over selective fertility and migration into study areas that could confound a comparison between babies that were conceived in treatment versus control groups after program started. Moreover, women who were pregnant at program launch and were first enrolled are the only program beneficiaries to receive the full 30 months of coverage as part of the pilot.¹⁵

¹³Survey data also ask about attendance at SBCC sessions and report lower participation. However, we trust administrative data from SCI more than survey reports, given that self-reported attendance data tend to be underreported due to recall bias whereas administrative data were collected by direct observation.

¹⁴Monitoring data were collected only in one township because of limited funding availability and the implementing partner’s (SCI) preference for the implementation of a related project (Field and Maffioli, 2021).

¹⁵The trial was registered on the AEA RCT registry (AEARCTR-0004189) in 2019, prior to the completion of endline data reconciliation. Although we did not register a separate Pre-Analysis Plan, the design, primary outcomes of interest and analysis sample are described in the trial registration document. More relevant, the analysis plan, primary outcome of interest (stunting) and power calculations are detailed in the 2016 Research Protocol that was delivered to our government partner for study approval prior to the onset of the RCT (https://elisamaffioli.files.wordpress.com/2021/12/160404_legacy_rct_research-protocol_final.pdf). The only change to the protocol that was made in 2019 prior to endline analysis (and described in the trial registry) was the decision to restrict the study to women who were pregnant at the time of program launch (April 2016) in order to mitigate concerns over endogenous selection into program benefits among those who become pregnant after the program was launched. Our experiment was powered on the primary outcome of stunting. Assuming an ICC of 0.03 and standard deviation =1.14, it was determined that 20 women per cluster (34 clusters) were necessary to detect a change in mean HAZ score from -1.42 to -1.12, at 90% power and alpha=0.01. Similarly, assuming an ICC of 0.03, it was determined that a similar number of observations (21 women per cluster) were necessary to detect a reduction of 8 percentage points in the proportion of children stunted, at 80% power and alpha=0.05. Our analysis sample

Women in this group were identified by conducting a full listing of individuals (Appendix Figure A2, listing I) in treatment and control study villages two months prior to the program being launched (February 2016) in which community health workers recorded every woman’s age and pregnancy status. However, several additional women, who were not classified as pregnant at the time of the initial listing, either because they were unaware of, or reluctant to report pregnancy status early on, or because they were not found in the village at the time of the initial listing, were reclassified as eligible midway through the program and received program benefits thereafter. However, significantly more of such women were found in treatment relative to control villages (unsurprisingly, given their greater incentive to reveal themselves in order to receive benefits once they became aware of the program), which could bias our estimates of program effects were we to include them in the analysis. In order to properly identify the true causal impacts of the program on actual beneficiaries, the decision was made to conduct a second listing (June-August 2017) (Appendix Figure A2, listing II) in order to identify all women who were eligible for enrollment before the program launch. We restricted our analysis sample to those who reported to be pregnant in listing II who were also found in listing I (2,338 women). We then confirmed their children’s date of birth with endline data, and select those mothers with an observed child between 22-35 months old at endline - these women were supposed to be 1-9 months gestation age in April 2016. Our final sample is composed by 2,134 women among those 2,338. Given this retrospective selection, we are unable to appropriately measure attrition.

The endline survey gathered data on household and individual characteristics, including weight and height data of all children under age 5, socio-economic status including income and assets, food consumption including dietary diversity, health-seeking behaviors emphasized in the SBCC sessions, credit and saving, decision-making, desired and realized fertility, and program participation. Our analysis estimates the effects of the interventions on the 2,154 children born to the selected 2,134 women for the analysis, i.e, those covered by the LEGACY program for their first 1,000 days of life.

2.4 Empirical Strategy

The random assignment of interventions across villages allows us to identify the causal effect of cash transfers and the relative importance of pairing cash transfers with SBCC

includes approximately 21 women in each of the 102 clusters.

by comparing endline outcomes across study arms. We estimate program effects with the following ordinary least squares (OLS) model:

$$Y_{iv} = \alpha + \beta * [Cash + SBCC]_v + \gamma * [CashOnly]_v + \delta X_{iv} + t + \epsilon_{iv}$$

where Y is the primary health outcome of interest for child or mother i living in village v . To capture nutritional impacts on children, we use child height and age data from endline to construct height-for-age z-scores (HAZ), a well-validated anthropometric measure of chronic malnutrition using the WHO child growth standards ([World Health Organization, 2006](#)). A HAZ value of -1 indicates that, given sex and age, a child’s height is one standard deviation below the median child in her age/sex reference group. In addition, we construct an indicator of stunting that equals one if $HAZ < -2$; an indicator of severe stunting that equals one if $HAZ < -3$; and an indicator of moderate stunting that equals one if $-3 \leq HAZ < -2$.¹⁶

To better understand potential pathways of influence, we examine a number of behavioral outcomes available at endline that capture economic and health determinants of malnutrition, focusing on knowledge and behaviors emphasized in the SBCC curriculum. These include infant feeding practices (dietary diversity and breastfeeding), total expenditures on food and healthcare, illness episodes and visits to skilled health personnel, and hand-washing behavior.

$Cash + SBCC$ is an indicator of whether the respondent’s village was assigned to T1, and $CashOnly$ is an indicator of whether the village was assigned to T2. The excluded group is the control group (CG). The model also controls for a number of predetermined observables, X , which include (i) individual demographic controls, including mother’s age and education, household head’s age and education, and child’s sex and age (child-level analysis only); and (ii) village-level controls, including distance to large and small markets, indicators for main source of livelihood (agriculture, livestock, or casual labor), availability of government provided electricity, and participation in a concurrent water, sanitation and hygiene (WASH) intervention. Village-level controls were collected prior to the launch of the program. In accordance with our randomization design, the model includes fixed effects for geographic strata (one variable for each triplet of sub-rural health catchment areas, t), which control for unobserved

¹⁶As a quality check, Appendix Figure A3 verifies no irregularity in the relationship between HAZ score and age of children in months. Note that, to measure anthropometrics in the field, enumerators were trained to follow [SMART technology](#), the gold standard measurement protocols for nutrition-related anthropometrics ([World Health Organization, 2006](#)).

characteristics that may vary across clusters and influence program delivery. Standard errors are clustered at the village cluster level, the unit of randomization.

To ensure that comparability across study arms was achieved by random assignment, we test for observable differences across experimental arms based on time-invariant individual characteristics (Appendix Table A4) and village-level data collected prior to the program launch (Appendix Table A1). Overall, the randomization was successful with only 3 out of 48 comparisons unbalanced across any of the treatment arms at a 95% confidence interval at the individual level, and 1 out of 51 unbalanced at a 90% confidence interval at the village level. Moreover, the mean differences that are significant are small in magnitude and work against our ability to attribute differences in stunting to treatment: for example, the head of the household’s education is lower in the *Cash + SBCC* arm, so could potentially bias downward a comparison of differences in stature at endline. We control for these unbalanced covariates in the empirical model.

It is important to note that our study design precludes non-random imbalances across treatment arms driven by selection into treatment. Although maternal cash transfers have the potential to incentivize women to become pregnant earlier than they would have otherwise or might encourage migration into treatment villages, our analysis sample is limited to women who were living in the village and pregnant *prior* to learning about the program, so is not subject to concerns over endogenous selection into the sample. However, because we run a handful of regressions on the sample of new mothers – the only group among which we are able to measure detailed infant feeding practices –, we also test directly whether the program led to fertility responses that could bias a comparison of infant feeding outcomes across experimental arms. Relative to the control group, women in *Cash + SBCC* and *CashOnly* are no more likely to be currently pregnant and do not report a greater number of pregnancies since the launch of the program, indicating that there is no increase in fertility in response to the program (Appendix Table A5).

3 Results

Our primary outcome of interest - as specified in the trial registration - is stunting, which provides a cumulative measure of nutritional benefits in the first 1000 days of life to children born to mothers in the treatment groups. For comparison purposes,

we also investigate flow measures of nutritional intake, including whether a child is underweight at endline, in order to gauge whether program beneficiaries continue to experience improved nutrition after program completion.¹⁷

3.1 Program Effects on Chronic Malnutrition

Table 1 reveals that, in control villages, a full 34% of children in this cohort are stunted, including 7% that are severely stunted. Table 1 shows that children born to mothers who received both cash and SBCC (*Cash + SBCC*, T1) from pregnancy until the child reached 24 months are an estimated 4.6 percentage points (a 13.5 percent reduction, $p < 0.05$) less likely to be stunted at 22-35 months of age compared to children living in control group households. Meanwhile, children in *CashOnly* (T2) villages are no less stunted than children in the control group, and we can firmly reject the equality of *Cash + SBCC* and *CashOnly* treatment effects on stunting ($pvalue = 0.02$). This pattern supports the hypothesis that cash transfers – even those directed to mothers – are only able to effectively combat chronic malnutrition in children when paired with an intervention that encourages behavior change.¹⁸ We do not find differential effects by gender of the child: results are statistically significant for both male (6.7 percentage points) and female (5.2 percentage points) children (Appendix Table A6).

Column 4 of Table 1 presents the treatment indicators regressed on a continuous measure of height for age, i.e., *HAZ* score. Although the *Cash + SBCC* intervention arm does not have a statistically significant effect on the continuous measure of height, the point estimate is large (0.074) and close to significance at the 10% level.¹⁹

Moreover, a more nuanced test of the distribution of *HAZ* scores reveals program

¹⁷Note that weight-based measures largely reflect recent nutritional intake and health status, whereas stunting is thought to reflect cumulative early childhood conditions (Hoddinott et al., 2013).

¹⁸We followed Newson and Team 2003 and Newson 2011 and report frequentist q-values (or adjusted p-values) for multiple-test procedures, by defining the discovery set to control either the familywise error rate (FWER) or the false discovery rate (FDR). For each input pvalue, the corresponding q-value is the lowest input uncorrected critical pvalue (FWER or FDR) which would cause the input pvalue to be included in the discovery set, if the specified multiple-test procedure was applied to the full vector of P-values. To be conservative, we use a step-up method where the q-value for each pvalue is equal to the cumulative minimum of all the r-values corresponding to pvalues of rank equal to or greater than that pvalue.


¹⁹As a point of comparison, the point estimate is similar in magnitude to a deworming intervention in Kenya that increased the mean *HAZ* by 0.09 SD, which was significant in a larger sample (Miguel and Kremer, 2004). Our estimates are also in line with the non-experimental estimates of *HAZ* impacts from other conditional or unconditional cash transfer programs (Biscaye et al., 2017; IEG World Bank, 2011) or early stimulation and nutrition interventions (Atanasio et al., 2018).

Table 1: Child stunting

	(1)	(2)	(3)	(4)
	Child stunted	Child moder- ately stunted	Child severely stunted	HAZ score (WHO)
Cash+SBCC	-0.046** (0.021) [0.033]	-0.053*** (0.018) [0.004]	0.007 (0.011) [0.513]	0.074 (0.047) [0.122]
CashOnly	-0.004 (0.021) [0.846]	-0.008 (0.020) [0.687]	0.004 (0.011) [0.728]	-0.017 (0.041) [0.681]
Observations	2151	2151	2151	2151
Mean Control	0.34	0.27	0.07	-1.57
Clusters	102	102	102	102
Cash+SBCC=CashOnly	0.02	0.01	0.75	0.02

Notes: The table presents OLS estimates of the effects of the maternal cash transfer program interventions on measures of stunting for children whose mothers were pregnant at enrollment, following WHO classification. “Cash+SBCC” indicates T1 villages, where cash transfers and SBCC activities were provided jointly; “CashOnly” indicates T2 villages, where only cash transfers were provided; the reference group are villages in the control group where neither cash transfers nor SBCC took place. Outcomes include the proportion of children stunted as children with Height for Age Z score (HAZ) < -2 (1); the proportion of children moderately stunted as children with HAZ < -2 and >= -3 (2); the proportion of children severely stunted as children with HAZ < -3 (3); and, HAZ (4). Controls include (i) individual demographic controls, including child’s sex and age, mother’s age and education, and household head’s age and education; (ii) village-level controls, including distance to large and small markets, main source of livelihood (agriculture, livestock, or casual labor), availability of government provided electricity, and participation in a concurrent WASH intervention. Fixed effects per geographic strata (34) are included. Standard errors are clustered at the village cluster level and reported in parenthesis. Adjusted p-values are calculated to account for multiple hypothesis testing following Newson and Team (2003) and Newson (2011) and reported in squared brackets. P-values from t-test from the difference in means are reported. ***p<0.01, ** p<0.05, * p<0.1.

effects that are consistent with the stunting results.²⁰ In particular, Figure 1 reveals a rightward shift in the distribution of HAZ scores among *Cash + SBCC* beneficiaries compared to the control group. We test whether the distributions are the same across interventions, using a Kolmogorov-Smirnov non-parametric test. We reject the null hypothesis of equal distributions of *HAZ* scores in the *Cash + SBCC* and *CashOnly* arms ($p=0.048$), and in the *Cash + SBCC* arm and the control group ($p=0.098$). We also reject the null hypothesis of equal distributions of *HAZ* scores in the *CashOnly* arm and control group ($p=0.071$).



Notes: This figure describes the distribution of Height for Age Z score (HAZ) for children whose mothers were pregnant at enrollment, by treatment status. “Cash+SBCC” indicates T1 villages, where cash transfers and SBCC activities were provided jointly; “CashOnly” indicates T2 villages, where cash transfers only were provided; “Control” indicates villages in the control group where neither cash transfers nor SBCC took place.

Figure 1: Child HAZ distribution, by treatment

Consistent with these distributional patterns, columns 2 and 3 of Table 1 show that the reduction in malnutrition achieved by the program corresponds to a decrease in the proportion of children moderately stunted (5.3 percentage points, a 19.6 percent reduction, $p<0.01$) but no change in the proportion severely stunted. This implies that, while chronic malnutrition in early childhood fell for many children at risk, the SBCC intervention did not succeed in combating malnutrition among the most vulnerable households.

We also present estimates of program participation on a flow measure of malnutrition, a binary indicator of a child being underweight (Table 2). It is important to

²⁰Note that the distribution of *HAZ* is centered at zero only if the population follows the WHO standard growth curve, which is not the case in our context given the high prevalence of malnutrition.

Table 2: Child underweight

	(1)	(2)	(3)	(4)
	Child un- derweight	Child moder- ately under- weight	Child severely under- weight	WAH score (WHO)
Cash+SBCC	-0.012 (0.025) [0.641]	-0.027 (0.023) [0.227]	0.016 (0.012) [0.197]	-0.004 (0.050) [0.934]
CashOnly	-0.006 (0.024) [0.812]	-0.026 (0.022) [0.237]	0.021 (0.012) [0.103]	-0.036 (0.049) [0.474]
Observations	2145	2145	2145	2145
Mean Control	0.28	0.24	0.04	-1.43
Clusters	102	102	102	102
Cash+SBCC=CashOnly	0.83	0.96	0.72	0.49

Notes: The table presents OLS estimates of the effects of the maternal cash transfer program interventions on measures of stunting for children whose mothers were pregnant at enrollment, following WHO classification. “Cash+SBCC” indicates T1 villages, where cash transfers and SBCC activities were provided jointly; “CashOnly” indicates T2 villages, where only cash transfers were provided; the reference group are villages in the control group where neither cash transfers nor SBCC took place. Outcomes include the proportion of children underweight as children with Weight for Age Z score (WAH) < -2 (1); the proportion of children moderately underweight as children with WAH < -2 and >= -3 (2); the proportion of children severely underweight as children with WAH < -3 (3); and, WAH (4). Controls include (i) individual demographic controls, including child’s sex and age, mother’s age and education, and household head’s age and education; (ii) village-level controls, including distance to large and small markets, main source of livelihood (agriculture, livestock, or casual labor), availability of government provided electricity, and participation in a concurrent WASH intervention. Fixed effects per geographic strata (34) are included. Standard errors are clustered at the village cluster level and reported in parenthesis. Adjusted pvalues are calculated to account for multiple hypothesis testing following Newson and Team (2003) and Newson (2011) and reported in squared brackets. Pvalues from t-test from the difference in means are reported. ***p<0.01, ** p<0.05, * p<0.1.

note that, by the time of endline, the majority of children in our sample were 22-35 months old and thus they had not received cash supplements for several months. For this reason we are less likely to observe differences in underweight at endline relative to what we might have observed during program implementation since flow measures of malnutrition are more likely to change during the program rather than after the program ended. This is in contrast to program effects on stunting, which are likely to be observed indefinitely. Differences in flow measures of nutritional intake at endline would indicate that the program benefits on nutrition continue even when the cash transfer is taken away from mothers, potentially because of habit formation in feeding behavior, knowledge gained directly through program participation, or indirect learning-by-doing about the value of dietary diversity. The evidence in Table 2 suggests no persistence in nutritional benefits (as measured by a child being underweight) of cash transfers once the transfer is terminated.


3.2 Heterogeneity

Although severe stunting is relatively low in this setting (7% in the control group), the absence of a program effect on this tail of the HAZ distribution is somewhat counterintuitive given that severe stunting is likely to be concentrated among the poorest households, and one might anticipate that the same amount of cash makes a bigger difference for households in more dire circumstances. One explanation that we consider is lower SBCC participation rates of extremely poor households. However, self-reported data from endline do not indicate significantly lower participation rates among households below median income relative to those above median income. Alternatively, the SBCC curriculum or mode of delivery may be inappropriately designed to meet the needs of very poor participants. For instance, households at risk of severe stunting may lack sufficient human capital to translate information into behavior change, or might face additional financial barriers to implementing changes such as diet diversification, even with the additional liquidity provided through a cash transfer. Finally, households at risk of severe stunting may be concentrated in villages with poor infrastructure to support the adoption of certain health practices such as access to clean water or food products.

To further evaluate whether the absence of a program effect on severe stunting is related to differences in socio-economic status (SES), we examine patterns of treatment effects across villages according to village-level SES. In the absence of baseline data

on wealth or income, we proxy for village SES with the average number of years of education attained by women in the sample.

It is first worth noting that rates of severe stunting are similar in magnitude across villages with low versus high average levels of parental education. This pattern alone suggests that some fraction of the population may face a poverty trap such as chronic reinfection that keeps them in a state of persistent malnutrition even when village resource levels rise. Interestingly, results from the subsample analysis indicate that the program effects are concentrated in low SES villages. In particular, we observe that the distribution of the HAZ scores is strongly shifted to the right in the *Cash + SBCC* intervention arm compared to the *CashOnly* arm or the control group only in low SES villages (Figure 2), and the difference is statistically significant. Meanwhile, in the above-median villages, *HAZ* score distributions are similar across experimental arms. Appendix Table A7 shows a similar heterogeneous pattern in a regression framework: the results indicate that relatively low-SES villages gain the most in terms of reductions in rates of stunting from maternal cash transfers combined with SBCC.



Notes: This figure describes the distribution of Height-for-age Z-score (HAZ) for children whose mothers were pregnant at enrollment, by treatment status, and by low vs high socio-economic status. As a proxy, we use the average number of years of education attained by women in the sample. “Cash+SBCC” indicates T1 villages, where cash transfers and SBCC activities were provided jointly; “CashOnly” indicates T2 villages, where cash transfers only were provided; “Control” indicates villages in the control group where neither cash transfers nor SBCC took place.

Figure 2: Child HAZ distribution by women village-level average education

Moreover, households below median income *within* the low-SES village benefit disproportionately in terms of reductions in moderate stunting relative to those above the village median (Appendix Table A8). That is, if we divide the sample into four groups according to both village-level SES and median household income at endline, the pattern of results indicates that reductions in moderate stunting due to exposure

to cash plus SBCC are fully concentrated among the quartile of households in the lower half of the income distribution *within* the lowest SES half of villages. Still, even among this subsample, severe stunting does not improve with either version of the program, consistent with non-convexities at extremely low levels of income.

Figure 2 also reveals that, within low-SES villages, the *CashOnly* and the *Cash + SBCC* treatments appear to operate on the same distribution of children who are on the left-hand side of the stunting distribution, but the *Cash + SBCC* treatment appears to push them relatively further rightward in terms of HAZ scores relative to the *CashOnly* treatment. That is, the patterns of HAZ score distributions indicates that the *CashOnly* treatment is effective for the same number of marginal responders as the *Cash + SBCC* treatment just to a lesser extent, rather than being equally effective for fewer kids.

3.3 Mechanisms

To better understand the channels through which a combination of *Cash + SBCC* generates positive effects on child health, in this section we explore program effects on maternal health behaviors, including the amount and type of food consumed, as measured by total food consumption and child food diversity, and health-care utilization as measured by total health expenditures. We also investigate whether treatment is associated with increases in maternal knowledge about child health production. Finally, to disentangle whether stunting impacts are driven by reductions in nutrition-depleting illnesses versus increases in the intake of nutritious food, we examine whether treatment is associated with reductions in reported episodes of child illness.

3.3.1 Program Effects on Maternal Health Behaviors

Table 3 describe the program impacts on behaviors that were emphasized in the SBCC curriculum. Specifically, we focus on the following key topics covered by the education sessions: food consumption, dietary diversity, breastfeeding, hand washing practices, and health-seeking behavior. To capture dietary diversity, we take the standard approach in the literature (based on WHO guidelines) of constructing a dietary diversity score (DDS) measured as the number of food groups consumed by the child in the previous day out of the following seven: (1) cereals, roots and tubers; (2) legumes and nuts; (3) milk and milk derivatives; (4) meat products (meat, poultry, offal, and fish); (5) eggs; (6) vitamin A-rich fruits and vegetables (leafy green vegetables, yellow fruits

and vegetables); and (7) other fruits and vegetables. A DDS of four is considered the minimum DDS for a healthy diet. As children in our sample are at least 22 months old, the DDS is measured excluding milk, following WHO guidelines.

Because stunting is associated with low levels of protein-rich foods in particular, we also look specifically at how treatment assignment influences food consumption in categories 2-5 aggregated. Not only are higher levels of protein-rich food consumption most likely to translate into reductions in child stunting, but – in addition to emphasizing the general importance of food diversity – SBCC health messaging focused specifically on the importance of feeding young children with protein-rich food groups.

Hand-washing practices are measured as a cumulative score of regularly adopted practices, where each practice is counted as 1 when the respondent reports washing hands with soap in that specific situation and 0 otherwise: after cleaning a baby’s bottom, after using the toilet, before preparing and eating food, before feeding children, after disposing of baby feces, before and after handling children, and on other occasions. Total food consumption is measured as recalled household consumption in the past 7 days and is winsorized at the 99th percentile level.²¹

Consistent with the stunting results, women assigned to the *Cash + SBCC* intervention spend significantly more money on food relative to the control group (increase of 2.2 USD, column 1). Women assigned to the *CashOnly* intervention also exhibit a positive change, but the increase in spending is significantly less stark: weekly food consumption in the *CashOnly* arm is 7.65% higher than the control group compared with a 15.13% difference among the *Cash + SBCC* arm (column 1). The changes in the weekly food consumption represents about 17% (for *CashOnly*) and 33% (for *Cash + SBCC*) of the cash amount (10,000MMK), suggesting a meaningful change in household food consumption that could account for large changes in stunting.

We also find positive changes in behaviors related to child food diversity and breastfeeding practices (Table 3), and these results are all significantly larger for the *Cash + SBCC* arm. In particular, we find a 0.655 unit increase in the child food diversity score (column 2) in the *Cash + SBCC* arm that is significantly different from the *CashOnly* and the control group. We also find a change in the proportion of children ever breastfed (0.7 percentage points, column 3) and in the proportion of children who received colostrum (2.1 percentage points, column 4) in the *Cash + SBCC* intervention arm only. Although these treatment effects are statistically significant, it is worth

²¹Results are robust to using raw consumption data.

Table 3: Maternal health behaviors

	(1)	(2)	(3)	(4)	(5)	(6)
	Tot. food consump- tion (USD)	Child dietary diversity (non-milk, 24 hrs recall)	Child ever breastfed	Child received colostrum	Index of hand- washing behavior (0-9)	Mother with at least 4 ANC visits to skilled health personnel
Cash+SBCC	2.168*** (0.373)	0.655*** (0.063)	0.007* (0.003)	0.021** (0.008)	0.651*** (0.128)	0.161*** (0.024)
CashOnly	1.097*** (0.365)	0.096 (0.070)	0.003 (0.004)	0.001 (0.010)	0.151 (0.118)	0.117*** (0.024)
Observations	2134	2154	2154	2151	2134	2134
Mean Control	14.33	3.39	0.99	0.96	2.60	0.67
Clusters	102	102	102	102	102	102
Cash+SBCC=CashOnly	0.00	0.00	0.26	0.06	0.00	0.05

Notes: The table presents OLS estimates of the effects of the maternal cash transfer program interventions on measures of behavior related to four topics covered by the education sessions in SBCC activities: Total food consumption (column 1), IYCF -including diet diversity (column 2), breastfeeding (columns 3-4), hand-washing practices (column 5), health-seeking behavior (column 6). “Cash+SBCC” indicates T1 villages, where cash transfers and SBCC activities were provided jointly; “CashOnly” indicates T2 villages, where cash transfers only were provided; the reference group are villages in the control group where neither cash transfers nor SBCC took place. Outcomes include: total food consumption, winsorized at the 99th percentile level (in last 7 days, in USD, exchange rate at 31 December 2018, 1); child diversity score constructed following WHO standards, for children at least 22 months old (2); the proportion of children ever breastfed (3); the proportion of children who received colostrum (4); an index of hand washing practices combining whether mothers report always washing hands after cleaning a baby’s bottom, after using the toilet, before preparing and eating food, before feeding children, after disposing of baby feces, before and after handling children, and on other occasions (5); the proportion of mothers receiving at least 4 Antenatal Care visits with skilled health personnel, as defined by WHO standards (6); Controls include (i) individual demographic controls, including child’s sex and age, mother’s age and education, and household head’s age and education for child-level analysis; mother’s age and education, and household head’s age and education for mother-level analysis; (ii) village-level controls, including distance to large and small markets, main source of livelihood (agriculture, livestock, or casual labor), availability of government provided electricity, and participation in a concurrent WASH intervention. Fixed effects per geographic strata (34) are included. Standard errors are clustered at the village cluster level and reported in parenthesis. ***p<0.01, ** p<0.05, * p<0.1.

noting that they are small in magnitude on account of the near universality of these practices prior to the intervention, as exhibited also by the high control group means. Essentially, SBCC participation shifts the fraction of children receiving colostrum from 96% to 98%, relative to both *CashOnly* and the control group. Finally, we find a 0.651 unit increase in the index of hand-washing behavior (column 5) in the *Cash + SBCC* group relative to both *CashOnly* and the control group, as well as a significant increase in the proportion of mothers attending at least 4 antenatal care visits (column 6) that is observed in both treatment arms but is significantly higher for the *Cash + SBCC* arm (16.1 percentage points for *Cash + SBCC* and 11.7 percentage points for *CashOnly*).

The absence of a program effect from cash alone on WASH and breastfeeding behaviors is unsurprising given that income alone should not be expected to increase rates of early initiation, so this can be readily interpreted as an impact of information on maternal health behavior. In contrast, ANC visits among the *Cash + SBCC* arm have the potential to be influenced by both an income effect of receiving cash transfers as well as an information effect of SBCC participation. However, the *difference* in health-seeking behavior between *CashOnly* and *Cash + SBCC* can be interpreted as the impact of information on health care utilization whereas the treatment effect women in the *CashOnly* group picks up the income effect on health-seeking behavior.

The results in Table 3 column 1 highlight how cash transfers with and without SBCC increase total household food consumption. Table 4 additionally explores the shares of the household food budget (in the last 7 days) spent on specific categories of foods: (animal or vegan) protein-rich foods, fruits and vegetables, staple carbohydrates including rice, wheat, maize, soybeans, potatoes, and other food, including oil and condiments. We find that cash transfers with or without SBCC increased household consumption of animal proteins (meat, fish, eggs and dairy), while decreasing consumption of less nutritious food such as staples and other food. However, the increase is statistically significantly higher for *Cash + SBCC* relative to *CashOnly* (5.8 percentage points vs 3.8 percentage points, respectively, column 1). In addition, those households exposed to the SBCC curriculum spent more of their budget on vegan proteins (pulses, 1.5 percentage points) and slightly more on vegetables and fruits (0.5 percentage points).²²

²²As shown in related research (Carneiro et al., 2021), if the program changed behavior and the effects were sustained, it is unsurprising that we find impacts on budget shares of food consumption in the previous seven days, even if the program have stopped some months before endline for the older children in the sample.

Table 4: Household budget shares of food consumption

	(1)	(2)	(3)	(4)	(5)
	Animal proteins	Vegan proteins	Vegetables and fruits	Staples	Other
Cash+SBCC	0.058*** (0.006)	0.015*** (0.002)	0.008** (0.004)	-0.040*** (0.007)	-0.039*** (0.003)
CashOnly	0.038*** (0.007)	0.005* (0.002)	-0.001 (0.004)	-0.015** (0.007)	-0.027*** (0.004)
Observations	2134	2134	2134	2134	2134
Mean Control	0.28	0.05	0.15	0.34	0.19
Clusters	102	102	102	102	102
Cash+SBCC=CashOnly	0.00	0.00	0.00	0.00	0.00

Notes: The table presents OLS estimates of the effects of the maternal cash transfer program interventions on household budget shares of food consumption. Outcomes include the share of total household food consumption spent on animal proteins (dairy, meat and fish, eggs, 1); vegan proteins (pulses and nuts, 2); vegetables and fruits (3); staples (4) and other food, including oil and other condiments (5). “Cash+SBCC” indicates T1 villages, where cash transfers and SBCC activities were provided jointly; “CashOnly” indicates T2 villages, where cash transfers only were provided; the reference group are villages in the control group where neither cash transfers nor SBCC took place. Controls include (i) individual demographic controls, including child’s sex and age, mother’s age and education, and household head’s age and education for child-level analysis; mother’s age and education, and household head’s age and education for mother-level analysis; (ii) village-level controls, including distance to large and small markets, main source of livelihood (agriculture, livestock, or casual labor), availability of government provided electricity, and participation in a concurrent WASH intervention. Fixed effects per geographic strata (34) are included. Standard errors are clustered at the village cluster level and reported in parenthesis. Pvalues from t-test from the difference in means are reported. *** $p < 0.01$, ** $p < 0.05$, * $p < 0.1$.

Reassuringly, these patterns are also reflected in measures of child dietary intake **as reported by the respondent for the previous 24 hours, including day and night feedings**. In particular, Table 5 investigates food intake from various food groups among the sample of children in the endline analysis sample. Data on child diets isolate changes in child feeding practices among mothers exposed to the SBCC curriculum rather than just household-level changes in food consumption, which may not be directed towards children. Table 5 shows clearly that children’s (reported) food intake improved systematically in the *Cash+SBCC* arm. Mothers in villages where cash transfers were supplemented with SBCC report that their children were 13.6 percentage points more likely to consume animal proteins than those in the control group. Households in the *CashOnly* arm are also 7.7 percentage points more likely to eat animal proteins than households in the control group, but the difference is significantly lower than observed among children in the *Cash + SBCC* arm (pvalue=0). In addition, children in households exposed to SBCC are also 21.7 percentage points and 9.3 percentage points more likely to consume vegan proteins and vegetables and fruits, respectively, compared to children in the control group. Instead, we do not find any statistically significant changes in children’s diets for the *CashOnly* arm, and more generally we do not observe shifts towards less nutritious food (columns 4 and 5).

These patterns suggest that one source of explanation for why cash transfers alone increase food consumption but do not reduce stunting may be that the additional food is not going towards children’s diets. This implies that one key mechanism through which SBCC might enhance the malnutrition impact of cash transfers is by convincing parents to direct increases in food consumption towards young children. For instance, they may be feeding school-going children without realizing the importance of critical windows of growth.²³

This pattern of results on program effects on specific food groups is consistent with the observed findings on child stunting, as protein-rich foods are generally needed to generate medium-run changes in chronic malnutrition through diet alone. In particular, a large literature in medicine and nutrition posits that nutrients from proteins primarily support child growth. Animal proteins from meat and fish are a source of amino acids (Laplante and Sabatini, 2012; Semba et al., 2016a,b), dairy products – and specifically cow milk – are an important source of amino acids and micronutrients (calcium, vitamin

²³Unfortunately, our data do not allow us to test this directly by looking at program effects on **weight-for-age** among older children and adults in the sample as biometric data were only collected for young children.

Table 5: Inclusion of protein-rich food groups in children’s diet

	(1)	(2)	(3)	(4)	(5)
	Animal proteins	Vegan proteins	Vegetables and fruits	Staples	Other
Cash+SBCC	0.136*** (0.015)	0.217*** (0.025)	0.093*** (0.022)	0.005 (0.006)	0.005 (0.006)
CashOnly	0.077*** (0.015)	0.025 (0.024)	-0.012 (0.023)	-0.001 (0.007)	0.001 (0.006)
Observations	2154	2154	2154	2154	2154
Mean Control	0.80	0.31	0.76	0.99	0.99
Clusters	102	102	102	102	102
Cash+SBCC=CashOnly	0.00	0.00	0.00	0.26	0.34

Notes: The table presents OLS estimates of the effects of the maternal cash transfer program interventions on children’s diet. Outcomes include the proportion of children eating: animal proteins (dairy, meat and fish, eggs, 1); vegan proteins (pulses and nuts, 2); vitamin-rich vegetables and fruits (3); staples (4); other, including oil and other condiments (5). “Cash+SBCC” indicates T1 villages, where cash transfers and SBCC activities were provided jointly; “CashOnly” indicates T2 villages, where cash transfers only were provided; the reference group are villages in the control group where neither cash transfers nor SBCC took place. Controls include (i) individual demographic controls, including child’s sex and age, mother’s age and education, and household head’s age and education for child-level analysis; mother’s age and education, and household head’s age and education for mother-level analysis; (ii) village-level controls, including distance to large and small markets, main source of livelihood (agriculture, livestock, or casual labor), availability of government provided electricity, and participation in a concurrent WASH intervention. Fixed effects per geographic strata (34) are included. Standard errors are clustered at the village cluster level and reported in parenthesis. Pvalues from t-test from the difference in means are reported. *** p<0.01, ** p<0.05, * p<0.1.

A, zinc) (Molgaard et al., 2011; Iannotti et al., 2013; Dyer et al., 2016), and eggs are an excellent source of choline (Semba et al., 2016c; Bekdash, 2016). Similarly, vegan proteins contain essential amino-acids, although in smaller doses. Meanwhile, the key nutritional value of vegetables and fruits is the vitamins (including vitamin A) and minerals (Gilbert, 2013) they contain. While vitamin deficiencies can increase risk of infections, their impact on stunting is likely to be of second-order importance compared to the role of protein-rich foods.

Compared to animal proteins, households are less likely to increase consumption of non-animal proteins and vitamin-rich foods without outside information and encouragement since they are less likely to be informed about the nutritional value of these food groups. The fact that SBCC is successful in promoting child consumption of vegan proteins in addition to greater consumption of animal proteins is particularly valuable given that vegan proteins are likely to be significantly more cost-effective means of increasing child protein intake.

Consistent with the changes in behavior, we find treatment effects on maternal knowledge (Appendix Table A9). Overall, we observe that maternal knowledge is quite high, with more than 90% of mothers in the control group reporting knowing that child food diversity is important (91%), the correct meaning of exclusive breastfeeding (94%), the best time to initiate breastfeeding (98%), and the best time to introduce complementary feeding (81%). While these high means suggest little scope for the SBCC intervention to improve maternal health knowledge, we find a small but statistically significant difference between *Cash + SBCC* (0.143 SD) and *CashOnly* (0.063 SD) when knowledge is indexed across all survey measures (pvalue=0.03). The absence of stronger treatment effects on individual indicators of maternal knowledge may reflect the fact that the survey instrument poorly captured key changes in knowledge gleaned through the program. Alternatively, SBCC may have operated not by changing specific knowledge of child health production, per se, but rather by way of cultural or social capital created during a variety of SBCC activities among mothers and other members of the community.

3.3.2 Program Effects on Child Illness

The evidence on maternal behavior indicates that reductions in child stunting were achieved in the *Cash + SBCC* treatment by encouraging mothers to increase total food consumption and shift children’s diets towards a broader array of protein-rich

foods. In addition, combining cash transfers with SBCC appears to have encouraged a higher rate of early initiation of breastfeeding, better hand-washing practices, and greater use of prenatal care. While these specific behaviors are unlikely to account for the large reduction in stunting observed in Table 1, it is possible that they correlate with a broader range of changes in infant feeding practices (e.g. longer duration of exclusive breastfeeding) and health care utilization (e.g. use of Oral Rehydration Therapy to curtail episodes of diarrheal disease) and disease control measures that could have more direct effects on stunting but are unobserved in our data.²⁴ That is, some of the impact on stunting may have resulted from changes in health behaviors other than food intake if they led to significant reductions in diarrheal disease, which in and of itself can produce chronic undernourishment in children.

To ascertain whether this is a source of stunting impacts, it is useful to note that hand-washing, breastfeeding exclusivity and health-seeking behavior can only impact child stunting via a reduction in nutrition-depleting illness. Hence, if these behavior changes contributed to reductions in child stunting, we should see corresponding reductions in diarrheal disease in the *Cash + SBCC* arm. Likewise, although antenatal care is unlikely to impact stunting itself, it could be correlated with an increase in expenditures on other health care services like oral rehydration therapy (ORT) that could have directly reduced nutrition-depleting illnesses.

Hence, to provide further evidence on the mechanisms through which the program reduces stunting, we examine survey data on child illness episodes and health care expenditures as a proxy for severity of illness. **However, please note that our data focus on episodes of illness in the last two weeks reported at endline when children covered by the program were 22-35 months old. While we would be interested in looking at changes in illness episodes when children were younger as those would be more informative of the effect on nutrition outcomes measured at endline, we argue that contemporaneous effects on illness still suggest that children in the *Cash + SBCC* arm appear to be stronger at endline, as they are affected by less nutrition-depleting illnesses.**

As shown in Table 6, we do not find any evidence that the interventions led to

²⁴Unfortunately, we did not collect data on the duration of breastfeeding at endline for those who stopped breastfeeding. Our endline survey asked mothers whether their children were ever breastfed and whether the child was still breastfed at the time of the data collection. At endline, the sample of children were 22-35 months old. About 71% were still breastfed. For those who were still breastfed, we asked when they would plan to interrupt. 59% mentioned a child age (mean 32 months), while 41% mentioned until the child won't breastfeed anymore.

changes in whether the child was brought in for treatment (column 4-5), or on total annual health expenditures on children under five (column 6). We find weak evidence that children were less likely to experience diarrhea in the last two weeks, but the effects are not statistically significantly different between the *CashOnly* and *Cash + SBCC* arms (pvalue=0.39), suggesting that SBCC, along with cash, did not contribute more to reducing the risk of infections. There are also no significant effects on the likelihood that children had pneumonia or fever in the past two weeks.²⁵

Table 6: Child Illness, seeking behavior and health expenditures

	(1)	(2)	(3)	(4)	(5)	(6)
	Diarrhea	Pneumonia	Fever	Seek treatment	Pay treatment	Health expenditures (children U5)
Cash+SBCC	-0.014* (0.008)	0.003 (0.027)	0.008 (0.026)	-0.021 (0.020)	-0.022 (0.021)	0.417 (2.516)
CashOnly	-0.020** (0.008)	0.002 (0.026)	-0.017 (0.025)	-0.008 (0.023)	-0.005 (0.025)	-0.033 (2.143)
Observations	2154	2154	2154	2154	2153	2134
Mean Control	0.03	0.20	0.72	0.89	0.86	27.75
Clusters	102	102	102	102	102	102
Cash+SBCC=CashOnly	0.39	0.93	0.31	0.56	0.46	0.85

Notes: The table presents OLS estimates of the effects of the maternal cash transfer program interventions on children outcomes. Outcomes include: the proportion of children with diarrhea in the past two weeks (1); the proportion of children with pneumonia in the past two weeks (2); the proportion of children with fever in the past two weeks (3); the proportion of children who sought treatment for that illness (4); the proportion of children who paid for the treatment (5); total health expenditures for children under 5 years old in the last year (in USD, 6). “Cash+SBCC” indicates T1 villages, where cash transfers and SBCC activities were provided jointly; “CashOnly” indicates T2 villages, where cash transfers only were provided; the reference group are villages in the control group where neither cash transfers nor SBCC took place. Controls include (i) individual demographic controls, including child’s sex and age, mother’s age and education, and household head’s age and education for child-level analysis; mother’s age and education, and household head’s age and education for mother-level analysis; (ii) village-level controls, including distance to large and small markets, main source of livelihood (agriculture, livestock, or casual labor), availability of government provided electricity, and participation in a concurrent WASH intervention. Fixed effects per geographic strata (34) are included. Standard errors are clustered at the village cluster level and reported in parenthesis. Pvalues from t-test from the difference in means are reported. *** p<0.01, ** p<0.05, * p<0.1.

²⁵We observe a reduction in diarrhea, but no effects on health seeking behavior. On one hand, it is possible that the interventions reduced illnesses, thus reducing demand for health care. On the other hand, cash transfers could increase demand for health care, alleviating financial constraints. This would potentially result in a net null effect. Yet, conditioning on experiencing an illness, we do not find impacts on health seeking behavior (not shown), suggesting that this is not the case.

3.3.3 Program Effects on Women’s Decision Making

We also confirm that the effects on child health are not driven by disproportionate changes in women’s decision-making power. We test whether female financial empowerment increased due to the amount of cash received in both interventions. In addition, it is theoretically possible that female decision-making power improved disproportionately in the *Cash + SBCC* arm due to an interaction effect on female agency of cash provision in conjunction with increased knowledge from participation in SBCC sessions. For example, fathers and elderly family members were also invited to participate in some of the SBCC sessions and their knowledge could shift decision making dynamics within the household, and thus the woman’s capacity to direct resources to child health. To explore this mechanisms, we use endline data on spousal decision-making over various spending categories. Appendix Table A10 shows no differential effects on female decision-making of either *CashOnly* or *Cash + SBCC*. Endline data indicate that mothers in both *CashOnly* and *Cash + SBCC* are no more likely to decide on expenditures from their own or spouse’s earnings, health care, major household purchases, or visiting relatives, than those in the control group.

3.3.4 Other Mechanisms

There a few other mechanisms that could have contributed to the additional benefits of *Cash + SBCC* compared to *CashOnly* on child health. First, SBCC is the strategic use of communication approaches to promote changes in knowledge, attitudes, norms, beliefs and behaviors, and this entails four levels of influence that interact to affect behavior: individual, family and peer networks, community and social/structural change (HC3, n.d.). Mother-to-mother support groups or community-level meetings could have contributed in changing social norms around the best young and child feeding practices. While we expect social norms to take time to change, we cannot exclude this played a role in our context. Since there is no variation in the SBCC activities conducted in *Cash + SBCC*, we are unable to explicitly test for this mechanism. Second, it is also possible that SBCC activities increased the salience and attentions on the targeted behaviors in the communities, as suggested by the high level of maternal health knowledge (Appendix Table A9). Finally, and related to the point above, experimenter demand effects could exist, since SBCC activities could have increased the attention to specific behaviors that women were asked about in the survey. Unfortunately, the data gathered in our survey tools do not allow us to test for these mechanisms. We

acknowledge their potential role in partly contributing to explaining the benefits on chronic malnutrition.

3.4 Robustness Checks

Although administrative program data show no documented cases of cash being delivered through the program to individuals residing in control villages, two forms of contamination in the SBCC intervention may compromise the validity of some of our estimates. First, according to SCI’s administrative data on the SBCC rollout, 18 villages assigned to the *CashOnly* treatment received SBCC activities for 20 rather than 30 months because of an error in program implementation. Second, SBCC activities were expanded to all *CashOnly* villages beginning on January 1, 2019, although part of the endline data were collected after December 31, 2018 at which point those respondents had already received at least one month of SBCC activities.

To address these two issues, we re-run the analysis excluding those 18 *CashOnly* villages and all mothers interviewed after December 2018, for a total of 138 mothers or 6.5 percent of the 2,134 women in the analysis sample. As reported in Appendix Table A11, the results are robust to these exclusions.

In addition, the main results are robust to considering the full endline sample (Appendix Tables A12) identified in the 2017 listing of pregnant mothers, which includes women who may have migrated into the village or become pregnant after the launch of the program. Results are also robust to the clustering of standard errors at the level of the program delivery (village) rather than the unit of randomization (health center catchment area), as shown in Appendix Table A13. Finally, the main results on moderate stunting are robust in a specification that includes only unbalanced covariates or excludes control variables (Appendix Table A14).

4 Conclusion

Our findings provide novel evidence from anthropometric data that (unconditional) maternal cash transfer programs delivered for the first 1000 days of life lead to statistically significant reductions in the proportion of children (moderately) stunted, but only when they are combined with intensive Social Behavior Change Communication (SBCC). The significant effects on stunting are concentrated among below-median-income households in low-SES villages, consistent with the notion that nutrition pro-

grams matter most where vulnerability to malnutrition is highest, yet the combination of interventions improved outcomes only among those at risk of moderate but not severe stunting. These patterns provide insight into which sub populations are most easily reached by such interventions, and indicate that greater efforts are needed to combat severe malnutrition. Tailoring SBCC programming to households with low levels of resources or facing chronic infections may be needed to address the most severe cases of malnutrition.

The program was successful in changing a number of maternal health behaviors, including total food consumption, dietary diversity, breastfeeding, hand-washing practices, and utilization of prenatal care. However, given that we do not find evidence that the combined interventions reduced child illness episodes, our pattern of results suggests that SBCC succeeded in reducing child malnutrition primarily through improvements in children’s diet, including a more diverse diet, and in particular protein-rich foods. In contrast, while cash alone increased child food intake and consumption of animal proteins, the changes were significantly smaller than those observed in the *Cash + SBCC* arm, and there was no increase in consumption of the more affordable vegan proteins emphasized in the SBCC curriculum. Moreover, the changes in child diet achieved through cash alone were insufficient to improve anthropometric indicators of malnutrition in young children who are in a critical window of growth.

Together, these findings underscore the importance of adding information components to social safety net programs involving cash disbursement in order to successfully change investment in human capital and thereby disrupt the inter-generational cycle of poverty. In our setting, the interventions show that (unconditional) cash transfers alone are insufficient to reduce chronic malnutrition. Instead, providing mothers with knowledge on how to use the additional disposable income can be transformative in reducing stunting. More generally, our analysis reveals that the combination of cash and SBCC tackled both the financial and information constraints that households faced in our context, and SBCC was fundamental in changing mothers’ feeding practices, thus reducing chronic malnutrition.

Further research is needed to better understand which particular curricular components are key to maximizing the child health gains of maternal cash transfer programs. In addition, more research is needed to establish whether information alone would be similarly effective in improving child health outcomes.

References

- Ahmed, Akhter, John Hoddinott, and Shalini Roy**, “Food transfers, cash transfers, behavior change communication and child nutrition: Evidence from Bangladesh,” 2019.
- Almond, Douglas and Janet Currie**, *Human Capital Development Before Age Five*. in *O. Ashenfelter and D. Card (eds.) Handbook of Labor Economics. Vol. 4*, Elsevier, 2011.
- and –, “Killing Me Softly: The Fetal Origins Hypothesis,” *Journal of Economic Perspectives*, 2011, 25 (3), 153–172.
- Attanasio, Orazio, Helen Baker-Henningham, Raquel Bernal, Costas Meghir, Diana Pineda, and Marta Rubio-Codina**, “Early Stimulation and Nutrition: The Impacts of a Scalable Intervention,” *Cowles Foundation Discussion Paper No. 2145*, 2018.
- Avula, Rasmi, Purnima Menon, Kuntal K Saha, Mahbubul Bhuiyan, Saiqa Siraj Anita S Chowdhury, and Edward E. Frongillo**, “A program impact pathway analysis identifies critical steps in the implementation and utilization of a behavior change communication intervention promoting infant and child feeding practices in Bangladesh,” *The Journal of Nutrition*, 2013, 143 (12), 2029–2037.
- Baird, Sarah, Craig McIntosh, and Berk Ozler**, “When the money runs out: Do cash transfers have sustained effects on human capital accumulation?,” *Journal of Development Economics*, 2019, 140, 169–185.
- Bastagli, Francesca, Jessica Hagen-Zanker, Luke Harman, Valentina Barca, Georgina Sturge, Tanja Schmidt, and Luca Pellerano**, “Cash transfers: what does the evidence say? A rigorous review of programme impact and of the role of design and implementation features,” 2016.
- Bekdash, Rola Aldana**, “Choline and the Brain: An Epigenetic Perspective,” *Advances in Neurobiology*, 2016, 12, 381–399.
- Bhutta, Zulfiqar A, Jai K Das, Arjumand Rizvi, Michelle Gaffey, Neff Walker, Susan Horton, Patrick Webb, Anna Lartey, and Robert Black**, “Evidence-based interventions for improvement of maternal and child nutrition: What can be done and what cost?,” *The Lancet*, 2013, 382 (9890), 452–477.
- Biscaye, Pierre, David Coomes, Elan Ebeling, Annie Rose Favreau, and Trygve Madsen**, “Review of Evidence on Long-term Impacts of Cash Transfer Programs,” 2017.

- Bobonis, Gustavo J**, “Is the allocation of resources within the household efficient? New evidence from a randomized experiment,” *Journal of political Economy*, 2009, 117 (3), 453–503.
- Cahyadi, Nur, Rema Hanna, Benjamin A. Olkena, Rizal Adi Prima, Elan Satriawan, and Ekki Syamsulhakim**, “Cumulative Impacts of Conditional Cash Transfer Programs: Experimental Evidence from Indonesia,” *American Economic Journal: Economic Policy*, 2020.
- Carneiro, Pedro, Lucy Kraftman, Giacomo Mason, Lucie Moore, Imran Rasul, and Molly Scott**, “The impacts of a multifaceted prenatal intervention on human capital accumulation in early life,” *American Economic Review*, 2021, 111 (8), 2506–49.
- Cecchini, Simone and Aldo Madariaga**, “Conditional cash transfer programmes: the recent experience in Latin America and the Caribbean,” 2011.
- Demographic Health Surveys**, 2015.
- Doyle, Orla**, “The First 2000 Days and Children’s Skills,” *Journal of Political Economy*, 2019.
- Duflo, Esther**, “Grandmothers and granddaughters: old-age pensions and intra household allocation in South Africa,” *The World Bank economic review*, 2003, 17 (1), 1–24.
- **and Christopher Udry**, “Intrahousehold resource allocation in Cote d’Ivoire: Social norms, separate accounts and consumption choices,” Technical Report, National Bureau of Economic Research 2004.
- Dyer, Adam, Cyrus Vahdatpour, Albert Sanfeliu, and Daniela Tropea**, “The Role of Insulin Like Growth Factor 1 (IGF-1) in Brain Development, Maturation and Neuroplasticity,” *Neuroscience*, 03 2016, 325.
- Engle, Patrice L, Maureen M Black, Jere R Behrman, Meena Cabral de Mello, Paul J Gertler, Lydia Kapiriri, Reynaldo Martorell, Mary Em- ing Young, and the International Child Development Steering Group**, “Strategies to Avoid the Loss of Developmental Potential in More than 200 Million Children in the Developing World,” *The Lancet*, 2007, 369, 229–242.
- Field, Erica and Elisa M Maffioli**, “A Comparison Between Government and NGO in Delivering Cash Transfers: Who Does Perform Better?,” 2021.
- Fiszbein, Ariel, Norbert Schady, Francisco H G Ferreira, Margaret Grosh, Niall Keleher, Pedro Olinto, and Emmanuel Skoufias**, “Conditional Cash Transfers: reducing present and future poverty,” 2009.

- Fitzsimons, Emla, Bansi Malde, Alice Mesnard, and Marcos Vera-Hernandez**, “Nutrition, information and household behavior: Experimental evidence from Malawi,” *Journal of Development Economics*, 2016, *122*, 113–126.
- Gentilini, Ugo**, “Cash Transfers in Pandemic Times,” 2022.
- Gilbert, Clare**, “What is vitamin A and why do we need it?,” *Community Eye Health.*, 2013, *84* (65), 1205–1223.
- HC3, Health Communication Capacity Collaborative**, “Social and Behavior Change Communication for Emergency Preparedness Implementation Kit,” Technical Report.
- Hoddinott, John, Ishita Ahmed, Akhter Ahmed, and Shalini Roy**, “Behavior change communication activities improve infant and young child nutrition knowledge and practice of neighboring non-participants in a cluster-randomized trial in rural Bangladesh,” *PLoS One*, 2017, *12* (6).
- , – , **Jere R Behrman, Lawrence Haddad, and Susan Horton**, “The economic rationale for investing in stunting reduction,” *Maternal and Child Nutrition*, 2013, *9* (2), 69–82.
- Iannotti, Lora, Ellen Muehlhoff, and Deirdre McMahon**, “Review of milk and dairy programmes affecting nutrition,” *Journal of Development Effectiveness*, 2013, *5* (1), 82–115.
- IEG World Bank**, “What Can We Learn from Nutrition Impact Evaluations?,” 2011.
- Jeong, Dahyeon and Iva Trako**, “Cash and In-Kind Transfers in Humanitarian Settings,” 2022.
- Kandpal, Eeshani, Harold Alderman, Jed Friedman, Deon Filmer, Junko Onishi, , and Jorge Avalos**, “A Conditional Cash Transfer Program in the Philippines Reduces Severe Stunting,” *The Journal of Nutrition*, 2016.
- Kling, Jeffrey R, Jeffrey B Liebman, and Lawrence F Katz**, “Experimental analysis of neighborhood effects,” *Econometrica*, 2007, *75* (1), 83–119.
- Laplante, Mathieu and David M. Sabatini**, “mTOR signaling in growth control and disease,” *Cell*, 2012, *149*, 274–293.
- Lépine, Aurélia and Eric Strobl**, “The effect of women’s bargaining power on child nutrition in rural Senegal,” *World Development*, 2013, *45*, 17–30.
- Leroy, Jef and Edward A Frongillo**, “Perspective: What Does Stunting Really Mean? A Critical Review of the Evidence,” *Advances in Nutrition*, 2019, *10* (2), 196–204.

- Levere, Michael, Gayatri Acharya, and Prashant Bharadwaj**, “The Role of Information and Cash Transfers on Early Childhood Development, Evidence from Nepal,” *Economic Development and Cultural Change*, 2022.
- Livelihoods and Food Security Fund**, “MEDIA RELEASE: LIFT and DSW announce USD9 million in COVID-19 cash support to mothers, pregnant women and social pension beneficiaries,” Technical Report 2020.
- Luo, Renfu, Yaojiang Shi, Linxiu Zhang, Huiping Zhang, Grant Miller, Alexis Medina, and Scott Rozelle**, “The Limits of Health and Nutrition Education: Evidence from Three Randomized-Controlled Trials in Rural China,” *CESifo Economic Studies*, 2012, 58 (2), 385–404.
- Manley, James, Harold Alderman, and Ugo Gentilini**, “More evidence on cash transfers and child nutritional outcomes: a systematic review and meta-analysis,” *BMJ global health*, 2022, 7 (4), e008233.
- , **Seth Gitter, and Vanya Slavchevska**, “How effective are cash transfers at improving nutritional status?,” *World Development*, 2013, 48, 133–155.
- , **Yarlina Balarajan, Shahira Malm, Luke Harman, Jessica Owens, Sheila Murthy, David Stewart, Natalia Elena Winder-Rossi, and Atif Khurshid**, “Cash transfers and child nutritional outcomes: A systematic review and meta-analysis,” *BMJ global health*, 2020, 5 (12), e003621.
- Miguel, Edward and Michael Kremer**, “Early Stimulation and Nutrition: The Impacts of a Scalable Intervention,” *Econometrica*, 2004, 72 (1), 159–217.
- Millan, Teresa Molina, Tania Barham, Karen Macours, John A Maluccio, and Marco Stampini**, “Long-Term Impacts of Conditional Cash Transfers in Latin America: Review of the Evidence,” *The World Bank Research Observer*, 2019, 34 (1), 119–159.
- Molgaard, Christian, Anni Larnkjaer, Karina Arnberg, and Kim F Michaelsen**, “Milk and growth in children: Effects of whey and casein,” *Nestle Nutrition workshop series. Paediatric programme*, 2011, 67, 67–78.
- Newson, Roger**, “QQVALUE: Stata module to generate quasi-q-values by inverting multiple-test procedures,” 01 2011.
- and **ALSPAC Study Team**, “Multiple—test Procedures and Smile Plots,” *The Stata Journal*, 2003, 3 (2), 109–132.
- Richter, Linda M, Bernadette Daelmans, Joan Lombardi, Jody Heymann, Florencia Lopez Boo, Jere R Behrman, Chunling Lu, Jane E Lucas, Rafael Perez-Escamilla, Tarun Dua, Zulfiqar A Bhutta, Karin Stenberg, and Paul**

- Gertler**, “Investing in the Foundation of Sustainable Development: Pathways to Scale Up for Early Childhood Development,” *The Lancet*, 2017, *389*, 103–118.
- Ritcher, Linda**, “Social cash transfers to support children and families affected by HIV/AIDS. Vulnerable Children and Youth Studies,” *Vulnerable Children and Youth Studies*, 2010, *5* (1), 81–91.
- Semba, Richard D, Indi Trehan, Marta Gonzalez-Freire, Klaus Kraemer, Ruin Moaddel, M Isabel Ordiz, Luigi Ferrucci, , and Mark J Manary**, “Perspective: the potential role of essential amino acids and the mechanistic target of rapamycin complex 1 (mTORC1) pathway in the pathogenesis of child stunting,” *Advances in Nutrition*, 2016, *7* (5), 853–865.
- , **Michelle Shardell, Fayrouz A Sakr Ashour, Ruin Moaddel, Indi Trehan, Kenneth M Maleta, M Isabel Ordiz, Klaus Kraemer, Mohammed A Khadeer, Luigi Ferrucci, and Mark J Manary**, “Child stunting is associated with low circulating essential amino acids,” *EBioMedicine*, 2016, *6*, 246–252.
- , **Pingbo Zhang, Marta Gonzalez-Freire, Ruin Moaddel, Indi Trehan, Kenneth M Maleta, M Isabel Ordiz, Luigi Ferrucci, and Mark J Manary**, “The association of serum choline with linear growth failure in young children from rural Malawi,” *American Journal of Clinical Nutrition*, 2016, *104* (1), 191–197.
- Soares, Fabio Veras, Rafael Perez Ribas, and Rafael Guerreiro Osorio**, “Evaluating the impact of Brazil’s Bolsa Familia: Cash transfer programs in comparative perspective,” *Latin American Research Review*, 2010, *45* (2), 173–190.
- The Republic of The Union of Myanmar**, “Myanmar National Social Protection Strategic Plan,” Technical Report 2014.
- Tirivayia, Nyasha, Marco Knowlesb, and Benjamin Davisb**, “The interaction between social protection and agriculture: A review of evidence,” *Global Food Security*, 2016, *10*, 52–62.
- United Nation Standing Committee on Nutrition**, “Sixth Report on the World Nutrition Situation: Progress in Nutrition,” 2010.
- Vasundhara, Donugama, Rajkumar Hemalatha, Saurabh Sharma, Baru Anantha Ramalaxmi, Varanasi Bhaskar, JagJeevan Babu, Radha Krishna Kankipati Vijaya, and RajaSriswan Mamidi**, “Maternal MUAC and fetal outcome in an Indian tertiary care hospital: A prospective observational study,” *Maternal & child nutrition*, 2020, *16* (2), e12902.
- Victora, Cesar Gomes, Mercedes De Onis, Pedro Curi Hallal, Monika Blössner, and Roger Shrimpton**, “Worldwide timing of growth faltering: revisiting implications for interventions,” *Pediatrics*, 2010, *125* (3), e473–e480.

World Bank, “World development indicators,” 2017.

World Food Program, “Fill The Nutrition Gap, Myanmar: Summary Report,” Technical Report 2019.

World Health Organization, “WHO child growth standards: Length/height-for-Age, weight-for-age, weight-for-length, weight-for-height and Body Mass Index-for-age: Methods and development,” 2006.

Appendix: Additional Tables and Figures

Table A1: Balance on village-level characteristics

	T1	T2	CG	PV (T1-CG)	PV (T2-CG)	PV (T1-T2)	N(T1/T2/CG)
Tot. population (No. HH)	182.17 (128.98)	175.54 (135.99)	160.92 (106.39)	0.263	0.418	0.747	133 / 135 / 139
Tot. literacy rate	85.19 (13.45)	85.50 (12.50)	83.58 (12.87)	0.384	0.247	0.866	133 / 135 / 139
Main livelihood: Agriculture	0.92 (0.26)	0.85 (0.36)	0.91 (0.29)	0.654	0.312	0.145	133 / 135 / 139
Main livelihood: Livestock	0.29 (0.45)	0.30 (0.46)	0.24 (0.43)	0.583	0.500	0.887	133 / 135 / 139
Main livelihood: Casual Labor	0.77 (0.42)	0.79 (0.41)	0.76 (0.43)	0.779	0.714	0.885	133 / 135 / 139
Type land-dry land farming	0.53 (0.50)	0.64 (0.48)	0.55 (0.50)	0.837	0.370	0.297	133 / 135 / 139
Type land-flood plains or irrigated	0.47 (0.50)	0.35 (0.48)	0.44 (0.50)	0.780	0.322	0.229	133 / 135 / 139
Accessible by car/truck in all weather	0.79 (0.41)	0.79 (0.41)	0.75 (0.44)	0.614	0.677	0.955	133 / 135 / 139
Village has Gov electricity	0.24 (0.43)	0.22 (0.42)	0.22 (0.42)	0.839	0.992	0.837	133 / 135 / 139
Village has primary school	0.63 (0.48)	0.58 (0.50)	0.59 (0.49)	0.544	0.832	0.411	133 / 135 / 139
Village has small markets	0.02 (0.15)	0.05 (0.22)	0.03 (0.17)	0.735	0.352	0.231	133 / 135 / 139
Village has home markets	0.97 (0.17)	0.96 (0.21)	0.96 (0.19)	0.773	0.729	0.551	133 / 135 / 139
Distance to large market	34.76 (24.83)	32.96 (20.05)	40.24 (26.39)	0.310	0.150	0.687	133 / 135 / 139
Distance to small markets	24.77 (18.62)	20.46 (15.49)	28.07 (23.55)	0.487	0.084*	0.214	133 / 135 / 139
Village has health facility	0.16 (0.37)	0.19 (0.39)	0.19 (0.40)	0.269	0.803	0.472	133 / 135 / 139
Village has midwife	0.21 (0.41)	0.21 (0.41)	0.24 (0.43)	0.358	0.269	0.937	133 / 135 / 139
Water shortage past year	0.42 (0.50)	0.46 (0.50)	0.36 (0.48)	0.441	0.189	0.632	133 / 135 / 139

Notes: This table presents the balance check on village characteristics by treatment arm for the sample of villages included in the analysis. Data are from the village census conducted in February 2016. T1 (“Cash+SBCC”) refer to villages where cash transfers and SBCC activities were provided jointly; T2 (“CashOnly”) refer to villages where cash transfers only were provided; CG (control group) refer to villages where neither cash transfers nor SBCC took place. Fixed effects per geographic strata (34) are included. Standard errors are clustered at the village cluster level and reported in parenthesis. Pvalues from t-test from the difference in means are reported. *** p<0.01, ** p<0.05, * p<0.1.

Table A2: Balance on woman-level characteristics

	T1	T2	CG	PV (T1-CG)	PV (T2-CG)	PV (T1-T2)	N(T1/T2/CG)
Resp married	0.96 (0.19)	0.97 (0.17)	0.96 (0.20)	0.541	0.199	0.386	840 / 802 / 695
Resp age	32.16 (6.37)	31.42 (6.52)	31.26 (6.17)	0.013**	0.683	0.049**	840 / 802 / 696
Resp educ years	5.70 (3.19)	5.86 (3.38)	6.08 (3.36)	0.132	0.370	0.499	840 / 802 / 696
HH size	4.95 (1.71)	4.99 (1.77)	4.81 (1.64)	0.224	0.103	0.719	840 / 802 / 696
Children U5	1.12 (0.37)	1.11 (0.34)	1.13 (0.38)	0.750	0.370	0.511	840 / 802 / 696
HH head female	0.08 (0.28)	0.09 (0.28)	0.07 (0.26)	0.593	0.421	0.725	840 / 802 / 696
HH head tot yrs educ	5.49 (3.19)	5.79 (3.35)	6.06 (3.35)	0.030**	0.292	0.177	840 / 802 / 696
HH head worked past 3m	0.89 (0.66)	0.83 (0.39)	0.86 (0.58)	0.650	0.386	0.225	840 / 802 / 696
HH head income	349964.00 (478142.69)	330294.12 (486734.66)	332720.14 (466277.61)	0.627	0.949	0.588	839 / 799 / 695
Any electricity	0.42 (0.49)	0.38 (0.48)	0.47 (0.50)	0.515	0.198	0.615	840 / 802 / 696
Always electricity	0.38 (0.48)	0.33 (0.47)	0.40 (0.49)	0.775	0.350	0.561	833 / 790 / 688
Cooking fuel electricity	0.29 (0.46)	0.24 (0.43)	0.29 (0.45)	0.953	0.360	0.420	840 / 802 / 696
Tot no. rooms in house	1.15 (0.81)	1.18 (0.80)	1.14 (0.75)	0.825	0.474	0.606	838 / 796 / 693
Improved roof material	0.87 (0.34)	0.85 (0.35)	0.87 (0.34)	0.991	0.681	0.645	840 / 802 / 696
Improved wall material	0.23 (0.42)	0.22 (0.41)	0.21 (0.41)	0.390	0.704	0.663	840 / 802 / 696
Improved floor material	0.33 (0.47)	0.32 (0.47)	0.31 (0.46)	0.695	0.898	0.809	840 / 802 / 696

Notes: This table presents the balance check on women characteristics by treatment arm for the sample of all mothers who were pregnant at enrollment (2,338). Data are from the endline data collection. T1 (“Cash+SBCC”) refer to villages where cash transfers and SBCC activities were provided jointly; T2 (“CashOnly”) refer to villages where cash transfers only were provided; CG (control group) refer to villages where neither cash transfers nor SBCC took place. Fixed effects per geographic strata (34) are included. Standard errors are clustered at the village cluster level and reported in parenthesis. Pvalues from t-test from the difference in means are reported. *** p<0.01, ** p<0.05, * p<0.1.

Table A3: Nutrition-related variables at baseline

	Mean	SD	Min	Max	Obs
<i>Child level variables</i>					
Stunted	0.288	0.453	0	1	566
Severely stunted	0.0389	0.193	0	1	566
Moderate stunted	0.249	0.433	0	1	566
Wasted	0.177	0.382	0	1	565
Severely wasted	0.0336	0.180	0	1	565
Moderate wasted	0.143	0.351	0	1	565
Under weight	0.310	0.463	0	1	658
Severely under weight	0.0578	0.233	0	1	658
Moderately under weight	0.252	0.435	0	1	658
Ever breastfed	0.982	0.132	0	1	281
Received colostrum	0.947	0.225	0	1	281
Received Minimum Dietary Diversity (6-23 mo)	0.379	0.489	0	1	58
<i>Mother level variables</i>					
Under nourished by MUAC	0.063	0.24	0	1	1126
Know the meaning of exclusively breastfeeding	0.95	0.22	0	1	1190
Know the best time to initiate breastfeeding	0.99	0.100	0	1	1190
Know the best time to introduce complementary feeding	0.85	0.36	0	1	1190
Use soap ever to wash hands	0.99	0.077	0	1	1190
Use soap after baby bottom cleaning (always)	0.17	0.38	0	1	1190
Use soap after eating (always)	0.47	0.50	0	1	1190
Use soap after toilet (always)	0.78	0.42	0	1	1190
Use soap before cooking (always)	0.37	0.48	0	1	1190
Use soap before eating (always)	0.51	0.50	0	1	1190
Use soap before feeding children (always)	0.17	0.37	0	1	1190
Use soap after disposing baby feces (always)	0.31	0.46	0	1	1190
Use soap b/a handling children (always)	0.043	0.20	0	1	1190
ANC at least 4 times with skilled health personnel	0.77	0.42	0	1	1190
<i>Household level variables (stunted children)</i>					
Budget share food	0.53	0.16	0.10	0.87	158
Budget share beverage	0.017	0.021	0	0.096	158
Budget share health	0.049	0.064	0.0016	0.50	158
Budget share educ	0.023	0.043	0	0.24	158
Budget share work-related	0.068	0.11	0	0.69	158
Budget share house	0.054	0.14	0	0.70	158
Budget share donations	0.061	0.047	0.0028	0.38	158
Budget share debt	0.076	0.10	0	0.46	158
Budget share bills	0.033	0.036	0	0.22	158
Budget share remittances	0.0038	0.031	0	0.31	158
Budget share transportation	0.035	0.057	0	0.43	158
Budget share other	0.045	0.060	0	0.39	158

Notes: This table presents nutrition-related variables collected at baseline, on the sub-sample of women and children that were matched with the analysis sample (at endline): 1,190 women and 691 children. Mean, standard deviation, minimum, maximum and number of observations are presented.

Table A4: Balance on woman-level characteristics for women sample followed-up at
 endline

	T1	T2	CG	PV (T1-CG)	PV (T2-CG)	PV (T1-T2)	N(T1/T2/CG)
Resp married	0.96 (0.19)	0.97 (0.17)	0.96 (0.20)	0.581	0.189	0.311	769 / 744 / 620
Resp age	32.07 (6.38)	31.41 (6.53)	31.39 (6.18)	0.082*	0.964	0.099*	769 / 744 / 621
Resp educ years	5.71 (3.17)	5.84 (3.32)	6.09 (3.36)	0.134	0.339	0.581	769 / 744 / 621
HH size	4.92 (1.69)	4.99 (1.80)	4.83 (1.63)	0.418	0.164	0.565	769 / 744 / 621
Children U5	1.14 (0.36)	1.12 (0.33)	1.14 (0.36)	0.975	0.326	0.279	769 / 744 / 621
HH head female	0.08 (0.28)	0.08 (0.27)	0.07 (0.26)	0.585	0.699	0.869	769 / 744 / 621
HH head tot yrs educ	5.47 (3.14)	5.81 (3.33)	6.04 (3.34)	0.032**	0.371	0.142	769 / 744 / 621
HH head worked past 3m	0.89 (0.65)	0.83 (0.38)	0.86 (0.60)	0.642	0.426	0.204	769 / 744 / 621
HH head income	350743.23 (466361.87)	335991.91 (500633.62)	334404.03 (483130.36)	0.654	0.967	0.704	768 / 742 / 620
Any electricity	0.42 (0.49)	0.38 (0.49)	0.48 (0.50)	0.503	0.177	0.585	769 / 744 / 621
Always electricity	0.38 (0.49)	0.33 (0.47)	0.41 (0.49)	0.718	0.309	0.564	762 / 733 / 614
Cooking fuel electricity	0.29 (0.46)	0.24 (0.43)	0.30 (0.46)	0.881	0.253	0.428	769 / 744 / 621
Tot no. rooms in house	1.14 (0.80)	1.17 (0.79)	1.13 (0.75)	0.890	0.536	0.621	767 / 739 / 619
Improved roof material	0.86 (0.34)	0.85 (0.35)	0.87 (0.34)	0.866	0.698	0.779	769 / 744 / 621
Improved wall material	0.23 (0.42)	0.22 (0.41)	0.21 (0.41)	0.436	0.750	0.678	769 / 744 / 621
Improved floor material	0.33 (0.47)	0.31 (0.46)	0.31 (0.46)	0.642	0.989	0.657	769 / 744 / 621

Notes: This table presents the balance check on individual characteristics by treatment arm for the sample of mothers who were pregnant at enrollment included in the analysis as followed-up at endline (2,134). T1 (“Cash+SBCC”) refer to villages where cash transfers and SBCC activities were provided jointly; T2 (“CashOnly”) refer to villages where cash transfers only were provided; CG (control group) refer to villages where neither cash transfers nor SBCC took place. Fixed effects per geographic strata (34) are included. Standard errors are clustered at the village cluster level and reported in parenthesis. Pvalues from t-test from the difference in means are reported. *** p<0.01, ** p<0.05, * p<0.1.

Table A5: Fertility

	(1)	(2)
	Pregnant	Tot no. pregnancies since launch of program
Cash+SBCC	0.010 (0.009)	0.002 (0.018)
CashOnly	0.001 (0.009)	-0.028* (0.017)
Observations	2134	2134
Mean Control	.03	1.14
Clusters	102	102

Notes: The table presents OLS estimates of the effects of the maternal cash transfer program interventions on measures of fertility. “Cash+SBCC” indicates T1 villages, where cash transfers and SBCC activities were provided jointly; “CashOnly” indicates T2 villages, where cash transfers only were provided; the reference group are villages in the control group (CG) where neither cash transfers nor SBCC took place. Outcomes include whether the mother is currently pregnant at endline (1), and her total number of pregnancies between June 2016 and endline, calculated from the household roster as the sum of biological living children under 5 years old. Controls include (i) individual demographic controls, including mother’s age and education, and household head’s age and education; (ii) village-level controls, including distance to large and small markets, main source of livelihood (agriculture, livestock, or casual labor), availability of government provided electricity, and participation in a concurrent WASH intervention. Fixed effects per geographic strata (34) are included. Standard errors are clustered at the village cluster level and reported in parenthesis. *** $p < 0.01$, ** $p < 0.05$, * $p < 0.1$.

Table A6: Child stunting, by gender

	(1)	(2)	(3)	(4)
	Child stunted	Child moder- ately stunted	Child severely stunted	HAZ score (WHO)
<i>Female children</i>				
Cash+SBCC	-0.052* (0.030)	-0.058* (0.030)	0.007 (0.017)	0.087 (0.068)
CashOnly	0.000 (0.033)	-0.009 (0.033)	0.009 (0.015)	-0.020 (0.061)
Observations	1044	1044	1044	1043
Mean Control	0.30	0.24	0.05	-1.48
Clusters	101	101	101	101
<i>Male children</i>				
Cash+SBCC	-0.067** (0.030)	-0.074*** (0.028)	0.007 (0.017)	0.082 (0.060)
CashOnly	-0.026 (0.031)	-0.022 (0.029)	-0.004 (0.017)	-0.005 (0.061)
Observations	1107	1107	1107	1107
Mean Control	0.38	0.30	0.08	-1.65
Clusters	102	102	102	102

Notes: The table presents OLS estimates of the effects of the maternal cash transfer program interventions on measures of stunting for children whose mothers were pregnant at enrollment, following WHO classification, by gender. “Cash+SBCC” indicates T1 villages, where cash transfers and SBCC activities were provided jointly; “CashOnly” indicates T2 villages, where only cash transfers were provided; the reference group are villages in the control group where neither cash transfers nor SBCC took place. Outcomes include the proportion of children stunted as children with Height for Age Z score (HAZ) < -2 (1); the proportion of children moderately stunted as children with HAZ < -2 and >= -3 (2); the proportion of children severely stunted as children with HAZ < -3 (3); and, HAZ (4). Controls include (i) individual demographic controls, including child’s sex and age, mother’s age and education, and household head’s age and education; (ii) village-level controls, including distance to large and small markets, main source of livelihood (agriculture, livestock, or casual labor), availability of government provided electricity, and participation in a concurrent WASH intervention. Fixed effects per geographic strata (34) are included. Standard errors are clustered at the village cluster level and reported in parenthesis. *** p<0.01, ** p<0.05, * p<0.1.

Table A7: Children stunting - by village socio-economic status (SES)

	(1)	(2)	(3)	(4)
	Child stunted	Child moder- ately stunted	Child severely stunted	HAZ score (WHO)
Panel A: Low SES Villages				
Cash+SBCC	-0.093** (0.037)	-0.093*** (0.032)	0.001 (0.020)	0.158** (0.071)
CashOnly	-0.041 (0.038)	-0.055 (0.035)	0.014 (0.020)	-0.026 (0.070)
Observations	1051	1051	1051	1051
Mean Control	0.34	0.27	0.07	-1.57
Clusters	92	92	92	92
Panel B: High SES Villages				
Cash+SBCC	0.044 (0.031)	0.020 (0.030)	0.024 (0.016)	-0.036 (0.078)
CashOnly	0.057* (0.033)	0.048 (0.031)	0.009 (0.014)	-0.069 (0.080)
Observations	961	961	961	961
Mean Control	0.34	0.27	0.07	-1.57
Clusters	86	86	86	86

Notes: The table presents OLS estimates of the effects of the maternal cash transfer program interventions on measures of stunting for children whose mothers were pregnant at enrollment, following WHO classification. “Cash+SBCC” indicates T1 villages, where cash transfers and SBCC activities were provided jointly; “CashOnly” indicates T2 villages, where only cash transfers were provided; the reference group are villages in the control group where neither cash transfers nor SBCC took place. Outcomes include the proportion of children stunted as children with Height for Age Z score (HAZ) < -2 (1); the proportion of children moderately stunted as children with HAZ < -2 and ≥ -3 (2); the proportion of children severely stunted as children with HAZ < -3 (3); and, HAZ (4). Controls include (i) individual demographic controls, including child’s sex and age, mother’s age and education, and household head’s age and education; (ii) village-level controls, including distance to large and small markets, main source of livelihood (agriculture, livestock, or casual labor), availability of government provided electricity, and participation in a concurrent WASH intervention. The analysis excludes the sample contaminated by the imperfect implementation (139 children). Low or high SES is proxied by the average number of years of education attained by women in the sample below or above the median.

Table A8: Children stunting - by village socio-economic status (SES) and household income

	(1)	(2)	(3)	(4)
	Child stunted	Child moder- ately stunted	Child severely stunted	HAZ score (WHO)
Panel A: Low SES, Low income				
Cash+SBCC	-0.140*** (0.050)	-0.122** (0.047)	-0.017 (0.029)	0.210** (0.099)
CashOnly	-0.053 (0.056)	-0.034 (0.052)	-0.019 (0.030)	-0.010 (0.104)
Observations	544	544	544	544
Mean Control	0.43	0.35	0.08	-1.72
Clusters	89	89	89	89
Panel B: Low SES, High income				
Cash+SBCC	-0.028 (0.047)	-0.059 (0.050)	0.031 (0.030)	0.029 (0.087)
CashOnly	-0.012 (0.043)	-0.065 (0.046)	0.053* (0.030)	-0.089 (0.079)
Observations	507	507	507	507
Mean Control	0.42	0.35	0.07	-1.69
Clusters	86	86	86	86
Panel C: High SES, Low Income				
Cash+SBCC	0.039 (0.045)	0.027 (0.039)	0.012 (0.024)	0.035 (0.105)
CashOnly	0.054 (0.047)	0.059 (0.040)	-0.005 (0.021)	-0.015 (0.097)
Observations	454	454	454	454
Mean Control	0.26	0.18	0.07	-1.50
Clusters	81	81	81	81
Panel D: High SES, High Income				
Cash+SBCC	0.029 (0.045)	0.003 (0.044)	0.025 (0.024)	-0.084 (0.111)
CashOnly	0.078 (0.053)	0.044 (0.053)	0.034 (0.025)	-0.178 (0.132)
Observations	507	507	507	507
Mean Control	0.27	0.22	0.05	-1.39
Clusters	80	80	80	80

Notes: The table presents OLS estimates of the effects of the maternal cash transfer program interventions on measures of stunting for children whose mothers were pregnant at enrollment, following WHO classification. “Cash+SBCC” indicates T1 villages, where cash transfers and SBCC activities were provided jointly; “CashOnly” indicates T2 villages, where only cash transfers were provided; the reference group are villages in the control group where neither cash transfers nor SBCC took place. Outcomes include the proportion of children stunted as children with Height for Age Z score (HAZ) < -2 (1); the proportion of children moderately stunted as children with HAZ < -2 and >= -3 (2); the proportion of children severely stunted as children with HAZ < -3 (3); and, HAZ (4). Controls include (i) individual demographic controls, including child’s sex and age, mother’s age and education, and household head’s age and education; (ii) village-level controls, including distance to large and small markets, main source of livelihood (agriculture, livestock, or casual labor), availability of government provided electricity, and participation in a concurrent WASH intervention. The analysis excludes the sample contaminated by the imperfect implementation (139 children). Low or high SES is proxied by the average number of years of education attained by women in the sample below or above the median. Low or high income is defined as below or above the household median income. Fixed effects per geographic strata (34) are included. Standard errors are clustered at the village cluster level and reported in parenthesis. *** p<0.01, ** p<0.05, * p<0.1.

Table A9: Maternal health knowledge

	(1)	(2)	(3)	(4)	(5)
	Mother knows child food diversity is important	Mother knows the meaning of exclusive breastfeeding	Mother knows the best time to initiate breastfeeding	Mother knows the best time to introduce complemen- tary feeding	Knowledge index
Cash+SBCC	0.033** (0.015)	0.031* (0.016)	0.013*** (0.005)	0.089*** (0.022)	0.143*** (0.033)
CashOnly	0.030** (0.014)	-0.000 (0.019)	0.009* (0.005)	0.033 (0.025)	0.063* (0.034)
Observations	2134	2134	2134	2134	2134
Mean Control	0.91	0.94	0.98	0.81	0
Clusters	102	102	102	102	102
Cash+SBCC=CashOnly	0.87	0.11	0.39	0.01	0.03

Notes: The table presents OLS estimates of the effects of the maternal cash transfer program interventions on measures of knowledge covered by the education sessions in SBCC activities. Outcomes include: the proportion of mothers who know the importance of food diversity in their children diet (1); the proportion of mothers who know the meaning of exclusive breastfeeding (2); the proportion of mothers who know the best time to initiate breastfeeding (3); the proportion of mothers who know the best time to introduce complementary feeding (4); an index of knowledge constructed following [Kling et al. 2007](#). “Cash+SBCC” indicates T1 villages, where cash transfers and SBCC activities were provided jointly; “CashOnly” indicates T2 villages, where cash transfers only were provided; the reference group are villages in the control group where neither cash transfers nor SBCC took place. Controls include (i) individual demographic controls, including child’s sex and age, mother’s age and education, and household head’s age and education for child-level analysis; mother’s age and education, and household head’s age and education for mother-level analysis; (ii) village-level controls, including distance to large and small markets, main source of livelihood (agriculture, livestock, or casual labor), availability of government provided electricity, and participation in a concurrent WASH intervention. Fixed effects per geographic strata (34) are included. Standard errors are clustered at the village cluster level and reported in parenthesis. Pvalues from t-test from the difference in means are reported. *** $p < 0.01$, ** $p < 0.05$, * $p < 0.1$.

Table A10: Women’s decision-making

	(1)	(2)	(3)	(4)	(5)
	Mother decides on own earnings	Mother decides on spouse earnings	Mother decides on health	Mother decides on major purchases	Mother decides on visit relatives
Cash+SBCC	0.027 (0.031)	-0.004 (0.024)	0.047 (0.031)	0.043* (0.024)	0.022 (0.027)
CashOnly	0.045* (0.027)	0.017 (0.024)	0.029 (0.028)	0.012 (0.025)	-0.014 (0.029)
Observations	1958	2088	2130	2130	2132
Mean Control	0.34	0.23	0.24	0.18	0.23
Clusters	102	102	102	102	102
Cash+SBCC=CashOnly	0.51	0.40	0.59	0.17	0.19

Notes: The table presents OLS estimates of the effects of the maternal cash transfer program interventions on measures of decision-making. Outcomes include: the proportion of mothers who decide on own (1) or spouse’s earnings (2); on health (3); or on major household purchases (4) or visiting relatives (5). “Cash+SBCC” indicates T1 villages, where cash transfers and SBCC activities were provided jointly; “CashOnly” indicates T2 villages, where cash transfers only were provided; the reference group are villages in the control group where neither cash transfers nor SBCC took place. Controls include (i) individual demographic controls, including child’s sex and age, mother’s age and education, and household head’s age and education for child-level analysis; mother’s age and education, and household head’s age and education for mother-level analysis; (ii) village-level controls, including distance to large and small markets, main source of livelihood (agriculture, livestock, or casual labor), availability of government provided electricity, and participation in a concurrent WASH intervention. Fixed effects per geographic strata (34) are included. Standard errors are clustered at the village cluster level and reported in parenthesis. Pvalues from t-test from the difference in means are reported. *** p<0.01, ** p<0.05, * p<0.1.

Table A11: Children stunting - addressing contamination

	(1)	(2)	(3)	(4)
	Child stunted	Child moder- ately stunted	Child severely stunted	HAZ score (WHO)
Cash+SBCC	-0.050** (0.022)	-0.057*** (0.018)	0.007 (0.011)	0.074 (0.047)
CashOnly	-0.002 (0.025)	-0.006 (0.022)	0.004 (0.012)	-0.031 (0.044)
Observations	2012	2012	2012	2012
Mean Control	0.34	0.27	0.07	-1.57
Clusters	102	102	102	102

Notes: The table presents OLS estimates of the effects of the maternal cash transfer program interventions on measures of stunting for children whose mothers were pregnant at enrollment, following WHO classification. ‘Cash+SBCC’ indicates T1 villages, where cash transfers and SBCC activities were provided jointly; ‘CashOnly’ indicates T2 villages, where only cash transfers were provided; the reference group are villages in the control group where neither cash transfers nor SBCC took place. Outcomes include the proportion of children stunted as children with Height for Age Z score (HAZ) < -2 (1); the proportion of children moderately stunted as children with HAZ < -2 and \geq -3 (2); the proportion of children severely stunted as children with HAZ < -3 (3); and, HAZ (4). Controls include (i) individual demographic controls, including child’s sex and age, mother’s age and education, and household head’s age and education; (ii) village-level controls, including distance to large and small markets, main source of livelihood (agriculture, livestock, or casual labor), availability of government provided electricity, and participation in a concurrent WASH intervention. The analysis excludes the sample contaminated by the imperfect implementation (139 children). Fixed effects per geographic strata (34) are included. Standard errors are clustered at the village cluster level and reported in parenthesis. *** p<0.01, ** p<0.05, * p<0.1.

Table A12: Children stunting - endline full sample

	(1)	(2)	(3)	(4)
	Child stunted	Child moder- ately stunted	Child severely stunted	HAZ score (WHO)
Cash+SBCC	-0.038** (0.016)	-0.042*** (0.015)	0.004 (0.009)	0.060* (0.035)
CashOnly	-0.006 (0.017)	-0.009 (0.016)	0.003 (0.009)	-0.014 (0.035)
Observations	3176	3176	3176	3176
Mean Control	0.35	0.28	0.06	-1.58
Clusters	102	102	102	102

Notes: The table presents OLS estimates of the effects of the maternal cash transfer program interventions on measures of stunting for children of the full endline sample, following WHO classification. “Cash+SBCC” indicates T1 villages, where cash transfers and SBCC activities were provided jointly; “CashOnly” indicates T2 villages, where only cash transfers were provided; the reference group are villages in the control group where neither cash transfers nor SBCC took place. Outcomes include the proportion of children stunted as children with Height for Age Z score (HAZ) < -2 (1); the proportion of children moderately stunted as children with HAZ < -2 and \geq -3 (2); the proportion of children severely stunted as children with HAZ < -3 (3); and, HAZ (4). Controls include (i) individual demographic controls, including child’s sex and age, mother’s age and education, and household head’s age and education; (ii) village-level controls, including distance to large and small markets, main source of livelihood (agriculture, livestock, or casual labor), availability of government provided electricity, and participation in a concurrent WASH intervention. Fixed effects per geographic strata (34) are included. Standard errors are clustered at the village cluster level and reported in parenthesis. *** p<0.01, ** p<0.05, * p<0.1.

Table A13: Children stunting - standard errors clustered at village level


	(1)	(2)	(3)	(4)
	Child stunted	Child moder- ately stunted	Child severely stunted	HAZ score (WHO)
Cash+SBCC	-0.046* (0.025)	-0.053** (0.023)	0.007 (0.014)	0.074 (0.054)
CashOnly	-0.004 (0.026)	-0.008 (0.024)	0.004 (0.014)	-0.017 (0.051)
Observations	2151	2151	2151	2151
Mean Control	0.34	0.27	0.07	-1.57
Clusters	407	407	407	407

Notes: The table presents OLS estimates of the effects of the maternal cash transfer program interventions on measures of stunting for children whose mothers were pregnant at enrollment, following WHO classification. ‘Cash+SBCC’ indicates T1 villages, where cash transfers and SBCC activities were provided jointly; ‘CashOnly’ indicates T2 villages, where only cash transfers were provided; the reference group are villages in the control group where neither cash transfers nor SBCC took place. Outcomes include the proportion of children stunted as children with Height for Age Z score (HAZ) < -2 (1); the proportion of children moderately stunted as children with HAZ < -2 and >= -3 (2); the proportion of children severely stunted as children with HAZ < -3 (3); and, HAZ (4). Controls include (i) individual demographic controls, including child’s sex and age, mother’s age and education, and household head’s age and education; (ii) village-level controls, including distance to large and small markets, main source of livelihood (agriculture, livestock, or casual labor), availability of government provided electricity, and participation in a concurrent WASH intervention. Fixed effects per geographic strata (34) are included. Standard errors are clustered at the village level. *** p<0.01, ** p<0.05, * p<0.1.

Table A14: Child stunting (alternative controls)


	(1)	(2)	(3)	(4)
	Child stunted	Child moder- ately stunted	Child severely stunted	HAZ score (WHO)
Panel A: Basic controls				
CASH+SBCC	-0.040 (0.025)	-0.051** (0.020)	0.011 (0.011)	0.051 (0.058)
CASH	-0.008 (0.026)	-0.012 (0.022)	0.004 (0.011)	-0.025 (0.054)
Observations	2151	2151	2151	2151
Mean Control	0.34	0.27	0.07	-1.57
Clusters	102	102	102	102
Panel B: No controls				
Cash+SBCC	-0.033 (0.026)	-0.046** (0.021)	0.013 (0.011)	0.038 (0.059)
CashOnly	-0.006 (0.027)	-0.010 (0.022)	0.004 (0.012)	-0.030 (0.056)
Observations	2151	2151	2151	2151
Mean Control	0.34	0.27	0.07	-1.57
Clusters	102	102	102	102

Notes: The table presents OLS estimates of the effects of the maternal cash transfer program interventions on measures of stunting for children whose mothers were pregnant at enrollment, following WHO classification. “Cash+SBCC” indicates T1 villages, where cash transfers and SBCC activities were provided jointly; “CashOnly” indicates T2 villages, where only cash transfers were provided; the reference group are villages in the control group where neither cash transfers nor SBCC took place. Outcomes include the proportion of children stunted as children with Height for Age Z score (HAZ) < -2 (1); the proportion of children moderately stunted as children with HAZ < -2 and \geq -3 (2); the proportion of children severely stunted as children with HAZ < -3 (3); and, HAZ (4). Panel A includes only unbalanced covariates (respondent’s age and head of the household’s years of education, Appendix Table 2); Panel B does not include any covariate. Fixed effects per geographic strata (34) are included. Standard errors are clustered at the village cluster level and reported in parenthesis. *** p<0.01, ** p<0.05, * p<0.1.




Notes: This figure presents the profile of the randomized controlled trial.

Figure A1: Profile of the Randomized Controlled Trial



Notes: This figure presents the timeline of the data collection rounds (survey activity) and the maternal cash transfer program rollout (program activity).

Figure A2: Timeline



Notes: This figure plots the relationship between HAZ score and age of the children (in months) in the sample.

Figure A3: Relationship between HAZ and age of children