



The Optimal Age to Stop Endoscopic Surveillance of Patients With Barrett's Esophagus Based on Sex and Comorbidity: A Comparative Cost-Effectiveness Analysis

Amir-Houshang Omidvari,¹ William D. Hazelton,² Brianna N. Lauren,³ Steffie K. Naber,¹ Minyi Lee,⁴ Ayman Ali,⁵ Claudia Seguin,⁴ Chun Yin Kong,⁴ Ellen Richmond,⁶ Joel H. Rubenstein,^{7,8} Georg E. Luebeck,² John M. Inadomi,⁹ Chin Hur,³ and Iris Lansdorp-Vogelaar¹

¹Department of Public Health, Erasmus MC University Medical Center Rotterdam, The Netherlands; ²Program in Computational Biology, Fred Hutchinson Cancer Research Center, Seattle, Washington; ³Department of Medicine, Columbia University, New York, New York; ⁴Institute for Technology Assessment, Massachusetts General Hospital, Boston, Massachusetts; ⁵Tulane University School of Medicine, New Orleans, Louisiana; ⁶Division of Cancer Prevention, National Cancer Institute, Rockville, Maryland; ⁷Barrett's Esophagus Program, Division of Gastroenterology, University of Michigan, Ann Arbor, Michigan; ⁸Veterans Affairs Center for Clinical Management Research, Ann Arbor, Michigan; and ⁹Department of Internal Medicine, University of Utah School of Medicine, Salt Lake City, Utah

See Covering the Cover synopsis on page 379.

BACKGROUND AND AIMS: Current guidelines recommend surveillance for patients with nondysplastic Barrett's esophagus (NDBE) but do not include a recommended age for discontinuing surveillance. This study aimed to determine the optimal age for last surveillance of NDBE patients stratified by sex and level of comorbidity. **METHODS:** We used 3 independently developed models to simulate patients diagnosed with NDBE, varying in age, sex, and comorbidity level (no, mild, moderate, and severe). All patients had received regular surveillance until their current age. We calculated incremental costs and quality-adjusted life-years (QALYs) gained from 1 additional endoscopic surveillance at the current age versus not performing surveillance at that age. We determined the optimal age to end surveillance as the age at which incremental cost-effectiveness ratio of 1 more surveillance was just less than the willingness-to-pay threshold of \$100,000/QALY. **RESULTS:** The benefit of having 1 more surveillance endoscopy strongly depended on age, sex, and comorbidity. For men with NDBE and severe comorbidity, 1 additional surveillance at age 80 years provided 4 more QALYs per 1000 patients with BE at an additional cost of \$1.2 million, whereas for women with severe comorbidity the benefit at that age was 7 QALYs at a cost of \$1.3 million. For men with no, mild, moderate, and severe comorbidity, the optimal ages of last surveillance were 81, 80, 77, and 73 years, respectively. For women, these ages were younger: 75, 73, 73, and 69 years, respectively. **CONCLUSIONS:** Our comparative modeling analysis illustrates the importance of considering comorbidity status and sex when deciding on the age to discontinue surveillance in patients with NDBE.

Keywords: EAC; Esophageal Cancer; Stop Age; CEA.

Barrett's esophagus (BE) is the only known precursor lesion for esophageal adenocarcinoma (EAC).¹ Patients with BE have a 10--55-fold higher risk of developing EAC than the general population. Fortunately, BE

surveillance and early detection and treatment of dysplasia may avert EAC development.² Generally, guidelines in the United States recommend endoscopic eradication therapy (EET) for high-risk patients (ie, patients with low-grade dysplasia [LGD] or high-grade dysplasia [HGD]). Furthermore, they recommend endoscopic surveillance every 3–5 years for nondysplastic BE (NDBE) patients, who are at a lower risk of developing EAC than those with dysplasia. However, there is no recommendation for the age to discontinue surveillance.^{3–6}

The expected benefits of surveillance diminish with advancing age and greater comorbidity due to lower life expectancy. For example, US men without comorbidities at age 68 have a life expectancy of 14.7 years, whereas US men with severe comorbidities at age 80 have a life expectancy of 5.3 years (Table 1).^{7,8} Therefore, the harms of endoscopic surveillance (eg, complications, false-positive results, and overtreatment) might outweigh the benefits (eg, deaths averted) for some patient populations. Patients with NDBE constitute about 90% of the total BE population.⁹ Additional surveillance endoscopies, particularly for this population, increase the cost of the surveillance program considerably, and continuation at older ages may, therefore, not be cost-effective.

To the best of our knowledge, no prior studies on BE surveillance have investigated the optimal age to

Abbreviations used in this paper: BE, Barrett's esophagus; CISNET, Cancer Intervention and Surveillance Modeling Network; EAC, esophageal adenocarcinoma; EACMo, Esophageal AdenoCarcinoma Model; EET, endoscopic eradication therapy; HGD, high-grade dysplasia; ICER, incremental cost-effectiveness ratio; LGD, low-grade dysplasia; LY, life year; MISCAN, Microsimulation Screening Analysis; MSCE, Multistage Clonal Expansion; NDBE, nondysplastic Barrett's esophagus; QALY, quality-adjusted life year; RFA, radiofrequency ablation; WTP, willingness-to-pay.

Most current article

© 2021 by the AGA Institute. Published by Elsevier Inc. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

0016-5085

<https://doi.org/10.1053/j.gastro.2021.05.003>

WHAT YOU NEED TO KNOW**BACKGROUND AND CONTEXT**

Current guidelines do not specify a recommended age for discontinuing surveillance endoscopy of patients with nondysplastic Barrett's esophagus (NDBE). This study aimed to determine an optimal age for last surveillance of patients with NDBE.

NEW FINDINGS

The optimal age for last surveillance of patients with NDBE without comorbidities for women is 75 years and for men is 81 years. However, it may be up to 6–8 years earlier if patients have severe comorbidities.

LIMITATIONS

Due to limited data, the prognosis of cancer was not varied by patient comorbidity.

IMPACT

Our analysis has important implications for surveillance of patients with NDBE. In addition to chronological age, the comorbidity status and sex of patients are important factors to inform the decision to discontinue surveillance.

discontinue surveillance of patients with NDBE with regard to the comorbidity level of patients. Evaluating the harms and benefits of many different stop ages in a clinical study would both be very costly and very time consuming. Therefore, modeling studies are required to estimate the effectiveness and cost-effectiveness of different stop ages.

In this study, we aimed to determine the optimal age of last surveillance for patients with NDBE by level of comorbidity using a comparative modeling approach.

Methods

We used 3 independently developed simulation models of EAC screening and surveillance that are part of the Cancer Intervention and Surveillance Modeling Network (CISNET) of the National Cancer Institute.

CISNET-EAC Models

We used the following models: (1) Microsimulation Screening Analysis model for esophageal adenocarcinoma (MISCAN-EAC) from Erasmus MC University Medical Center Rotterdam and the University of Utah; (2) Esophageal Adeno-Carcinoma Model (EACMo) from the Columbia University Medical Center and Massachusetts General Hospital; and (3) Multistage Clonal Expansion for EAC model from the Fred Hutchinson Cancer Research Center (MSCE-EAC). Each model was independently calibrated to common calibration targets based on US Surveillance, Epidemiology, and End Results cancer registry data until 2014.¹⁰

In all 3 models, it was assumed that EAC only develops in patients with BE. Healthy (asymptomatic) individuals and individuals with symptomatic gastroesophageal reflux disease may develop NDBE, which can progress to LGD and then HGD. Patients with BE with HGD may develop preclinical EAC, which can then progress to clinical EAC as symptoms develop. Individuals with clinical EAC may die of the disease with probabilities dependent on age and stage. More details on the

structure and quantifications of the models have been published and are available online.^{9,11,12}

Simulated Population and Intervention

For the base case, we simulated 200 cohorts of US patients diagnosed with NDBE, and followed them until death or age 100 years. Each cohort was defined by a unique combination of starting age (66–90 years), sex (man or woman), and comorbidity level (none, mild, moderate, or severe) (Table 1). EACMo and MSCE-EAC models simulated approximately one million patients in each cohort, and the MISCAN model simulated 265,000 patients. However, the results are reported per 1000 patients diagnosed with NDBE.

We used the cancer-free age, sex, and comorbidity-specific life tables from Lansdorp-Vogelaar et al^{7,8} and adjusted them to additionally include age and sex-specific mortality due to all cancers except esophageal cancers from CDC Wonder.¹³ Surveillance for patients with NDBE occurred every 3 years after the initial diagnosis, which was assumed to occur between 60 and 62 years of age.

For each cohort, we simulated an additional surveillance at the current age, or no further surveillance. For example, a 70-year-old patient with NDBE with a mild comorbidity level either did or did not receive 1 more surveillance at age 70.

Patients who were diagnosed with LGD received a repeat endoscopy with biopsies after 2 months of treatment with high-dose proton pump inhibitor to confirm LGD.^{14,15} Patients with HGD or confirmed LGD received EET followed by surveillance until death. In case of recurrence, patients received radio-frequency ablation (RFA) touch-ups followed by surveillance. The post-treatment surveillance strategies were simulated according to the outcome of initial EET or RFA touch-ups (Supplementary Table 1). Patients with treatment failure or recurrences more than 3 times did not receive additional RFA touch-ups and underwent surveillance until cancer diagnosis or death. Treatment and surveillance assumptions are presented in detail in Supplementary Table 2.

Costs and Utilities

The costs of endoscopies and EETs in \$US were estimated using the 2015 reimbursement rates from Centers for Medicare and Medicaid Services.¹⁶ The costs and utilities of cancer care by stage at diagnosis and those of complications due to endoscopy and EET were estimated using published literature (Supplementary Table 2).^{17–23} All costs and utilities were discounted at an annual rate of 3%.²⁴

Outcomes and Analysis

Using the average results of the 3 models for every cohort, we calculated the number of EAC cases, EAC deaths, life years (LYs), and quality-adjusted life years (QALYs) with and without 1 more surveillance. To estimate the total costs, we calculated the cost of cancer care, surveillance endoscopies, EETs, RFA touch-ups, and treatment of complications (ie, bleeding, perforation, and stricture) from a third-party payer perspective.

Subsequently, we calculated incremental costs and QALYs gained from 1 additional endoscopic surveillance at the current age versus not performing surveillance at that age, using the average results. The incremental cost-effectiveness ratio (ICER) of performing a last surveillance was calculated for all 25

Table 1. Overview of Comorbidity Levels, Associated Conditions, and Life Expectancies at Selected Ages 68, 74, and 80 Years in Men

Comorbidity level	Conditions included	Life expectancy, y		
		At age 68	At age 74	At age 80
No	None of the conditions listed for mild, moderate, or severe	14.7	11.5	8.5
Mild	History of myocardial infarction, ulcer, or rheumatologic disease	13.7	11.0	8.0
Moderate	Peripheral vascular disease, cerebrovascular disease, paralysis, diabetes, or combinations of mild conditions	12.8	9.8	6.9
Severe	AIDS, chronic obstructive pulmonary disease, cirrhosis, chronic hepatitis, chronic renal failure, dementia, congestive heart failure, or combinations of at least 1 moderate condition (except diabetes) with any mild or moderate condition	9.7	7.3	5.3

AIDS, acquired immunodeficiency syndrome.

potential stopping ages (66–90 years), and the age with the highest ICER just less than the willingness-to-pay (WTP) threshold of \$100,000 per QALY gained was considered the optimal age of last surveillance.

Sensitivity Analysis

Separate results of each model function as an independent sensitivity analysis of underlying assumptions for the natural history of EAC. In addition, we simulated cohorts of patients diagnosed with NDBE at ages 50, 51, and 52 or 70, 71, and 72 years in addition to 60, 61, and 62 to evaluate the robustness of our findings. We also simulated cohorts of patients with NDBE who underwent surveillance every 5 years (instead of 3 years) after the initial diagnosis at ages 60–62 years. Furthermore, we considered EAC survival probabilities, endoscopy and EET complication rates, and disutility scores depending on the comorbidity level of patients. For patients without comorbidity, we considered 50% lower complication rates and disutilities, and 10% higher EAC survival than base case, whereas for patients with mild comorbidity, we considered the same values assumed in the base case. For patients with moderate or severe comorbidity, we considered 50% or 100% higher complication rates and disutilities, and 10% or 20% lower EAC survival than base case, respectively.

All authors had access to the study data and reviewed and approved the final manuscript.

Results

Results for Men

Table 2 presents lifetime net benefits and costs of 1 additional endoscopic surveillance at selected stopping ages of 68, 74, 80, and 86 years. One more surveillance at age 68 in 1000 patients with NDBE without comorbidity prevented 10 more EAC cases than not performing surveillance at that age. Overall, 56 more QALYs were gained at an incremental cost of more than \$1 million, resulting in an ICER of \$23,600 per QALY, which was well less than the WTP threshold. The same comparison for patients with NDBE with comorbidities showed that 1 additional surveillance at age 68 years

prevented fewer EAC cases and deaths, which led to higher net costs and lower QALYs. Nonetheless, the ICERs remained less than the WTP threshold, and surveillance at age 68 was considered cost-effective for patients with NDBE of all comorbidity levels.

By increasing the age of the patients with NDBE, the net benefits of 1 additional surveillance decreased, and the ICERs increased accordingly. The ICERs of 1 additional surveillance versus not performing surveillance at ages 74, 80, or 86 years were higher than at age 68 years irrespective of comorbidity level (Table 2). An additional surveillance at age 74 for patients with NDBE with severe comorbidities was not cost-effective, with an ICER greater than \$100,000/QALY. Similarly, an additional surveillance at age 80 was not cost-effective for patients with NDBE with moderate or severe comorbidities. At age 86, 1 more surveillance was not cost-effective for any level of comorbidity.

For men without comorbidity, 1 additional surveillance at age 82 years in comparison with not performing surveillance at that age resulted in an ICER of \$116,300 per QALY, whereas the same comparison at age 81 years resulted in an ICER of \$99,000 per QALY. Therefore, 81 years was considered the optimal age of last surveillance for individuals without comorbidity. For individuals with mild, moderate, and severe comorbidity, the optimal ages of last surveillance using the average results of the 3 models were 80, 77, and 73 years, respectively (Figure 1A).

Results for Women

Similar to men, the net benefits of 1 additional surveillance of women with NDBE decreased with increasing age and comorbidity. However, the ICERs of 1 more surveillance in women were generally higher than those for men of similar age and comorbidity status (Supplementary Table 3). For example, surveillance of women aged >75 years was not cost-effective (ICERs >\$101,800/QALY) for any level of comorbidity.

Consequently, the optimal ages of last surveillance were lower in women than in men. For women without comorbidity, 75 years was the optimal age of last surveillance with

Table 2. Lifetime Net Benefits and Costs of 1 Additional Endoscopic Surveillance, For Example, at Ages 68, 74, 80, and 86 vs Not Performing Surveillance at That Age Per 1000 Men With NDBE

Age	Comorbidity level ^a	EAC prevented	EAC death prevented ^b	QALYs gained	Endoscopies	EET and touch-ups	Net cost (\$)	ICER
68	No	10	11	56	1952	81	1,328,609	23,620
	Mild	9	10	49	1910	79	1,343,689	27,183
	Moderate	8	9	44	1875	78	1,360,570	30,927
	Severe	5	7	28	1732	73	1,393,730	49,673
74	No	6	8	31	1670	70	1,269,878	41,302
	Mild	6	7	28	1650	70	1,275,048	45,230
	Moderate	5	6	22	1591	68	1,290,369	59,030
	Severe	3	4	13	1465	64	1,296,268	101,966
80	No	3	5	14	1401	62	1,192,137	83,986
	Mild	3	4	12	1381	61	1,195,159	96,407
	Moderate	2	3	8	1322	60	1,195,875	143,993
	Severe	1	2	4	1220	57	1,175,899	269,344
86	No	1	2	4	1128	55	1,083,739	254,074
	Mild	1	2	4	1112	54	1,081,072	295,144
	Moderate	1	2	2	1068	53	1,072,173	482,703
	Severe	0	1	0	981	51	1,033,270	2,352,232

^aOverview of comorbidity levels and associated conditions can be found in [Table 1](#).

^bSurveillance of patients with NDBE can prevent EAC deaths in 2 ways: (1) by finding and treating patients with BE with LGD and HGD, this way preventing both EAC incidence and thus EAC death; and 2) by finding and treating EAC early, and this way preventing death from EAC but not EAC incidence. Therefore, the number of prevented deaths due to EAC can be higher than the number of prevented EAC cases.

an ICER of \$84,200/QALY. Surveillance of patients with higher comorbidity levels resulted in higher ICERs and lower optimal stopping ages. For women with mild and moderate comorbidity, the optimal age of last surveillance was the same: 73 years; however, the ICERs were different (\$88,000 vs. \$98,700 per QALY, respectively). For women with severe comorbidity, the optimal stopping age was 69 years ([Figure 1B](#)).

Sensitivity Analysis

The separate results of each model consistently showed that women had lower optimal ages for last surveillance than men. All 3 models also showed lower optimal stopping ages for patients with higher comorbidity levels. However, the results from the EACMo suggested earlier optimal ages for last surveillance compared with other models, particularly for women with NDBE ([Table 3](#)).

Our results were also robust to surveillance every 5 years, different diagnosis ages, as well as variation in the assumed complication rates, EAC survival probabilities, and utility values by comorbidity level. Only small changes in the optimal age of last surveillance of NDBE patients by these sensitivity analyses were observed ([Table 3](#), [Figure 2](#)).

Discussion

Our comparative modeling analysis indicates that the optimal age for last surveillance of patients with NDBE depends on the sex and the comorbidity level of patients. We found that for men with NDBE without comorbidity, the optimal age for last surveillance is 81 years, whereas it may be up to 8 years earlier for those with comorbidity. For women, we found that without comorbidity, the optimal age for last surveillance of patients with NDBE is 75 years but can be up to 6 years earlier if patients have comorbidities.

Generally, by increasing the age and level of comorbidity of patients, the life expectancy is decreased and consequently, the benefit of surveillance is decreased. Therefore, after a certain age, surveillance of patients with NDBE is no longer cost-effective. Despite having a longer life expectancy, women have a lower optimal age for last surveillance due to a lower lifetime risk of EAC in women than in men. The separate results of each model showed the same patterns. However, the EACMo suggests earlier ages to discontinue surveillance in men and women compared with other models. This discrepancy can be explained by different natural history assumptions between the models. EAC incidence varies by age across the models. At older




Figure 1. Incremental cost-effectiveness ratio of surveillance of patients with NDBE at different ages by level of comorbidity, men (A) and women (B).

ages, the cumulative incidence of EAC in the EACMo is lower than the other 2 models. Therefore, patients with NDBE in the EACMo are more likely to die of other causes before progression to EAC occurs. This is the reason that surveillance of patients with NDBE at later ages in the EACMo was not cost-effective, unlike in the other 2 models.

None of the previous analyses examining the cost-effectiveness of surveillance of patients with NDBE evaluated the optimal age to discontinue surveillance.^{15,25,26} However, we can compare our findings to a previous study evaluating the age of colorectal, prostate, and breast

cancer screening cessation based on comorbidity level. This study found that people with higher comorbidity level gained less benefits from cancer screening and suggested to discontinue screening earlier.⁸ A prior cost-effectiveness analysis also showed that the comorbidity status of individuals undergoing colorectal cancer screening had a large impact on the effectiveness of the screening program. Screening was, therefore, cost-effective up to a lower age for people with comorbidities compared with those without.²⁷

In our base case analysis, we simulated cohorts of patients with NDBE diagnosed at age 60 years, because

Table 3. The Optimal Age of Last Surveillance Based on Sensitivity Analysis and Comorbidity Level

Comorbidity/analysis	Men				Women			
	No	Mild	Moderate	Severe	No	Mild	Moderate	Severe
Base case	81	80	77	73	75	73	73	69
Models								
MISCAN-EAC	83	82	79	75	77	74	74	71
EACMo	78	77	74	70	69	66	≤65	≤65
MSCE-EAC	81	80	77	75	78	75	74	72
NDBE by diagnosis age								
Age 50/51/52	81	80	77	74	76	74	73	70
Age 70/71/72	80	79	77	75	75	75	74	≤70
Surveillance every 5 y	81	81	78	75	77	75	74	71
Complication ^a rate	81	80	77	73	76	73	72	69
Utility values ^a	81	80	77	73	76	73	72	69
EAC survival ^a	80	79	77	74	75	73	73	69

^aFor patients without comorbidity, we considered 50% less complications and disutilities, and 10% less EAC mortality than base case. For patients with moderate comorbidity, we considered the same as what we assumed in the base case. For patients with moderate or severe comorbidity, we assumed 50% or 100% higher complication rate and disutilities, and 10% or 20% higher EAC mortality rate than base case, respectively.

the mean age of patients with BE at diagnosis has been reported to be older than 60 years.²⁸⁻³⁰ However, in sensitivity analyses, we assumed a longer interval for surveillance of patients with NDBE, and younger and older ages of diagnosis and varied utility values, complication, and EAC survival probabilities based on the comorbidity status of patients. Our results were quite robust for these external model parameters. However, they depended quite heavily on the model used (ie, on structural model assumptions). The main differences between the models are the time it takes to progress from NDBE to EAC and when BE develops in a patient (ie, how long a patient has lived with BE at the time of diagnosis). Because these patterns are still unknown, future linkage studies with long-term

follow-up might help to address these issues. Nevertheless, all 3 models in our study show that surveillance should not continue indefinitely, even in patients without any comorbidity.

Our study has some limitations. We are unaware of life tables for patients younger than 66 years of age with different comorbidity levels and, therefore, we could not determine the optimal age of last surveillance if it was younger than 66 years (for the EACMo, this was the case for women with moderate or severe comorbidity). However, this limitation did not affect our combined results. In addition, due to the limited data, we could not apply the impact of patient comorbidity level on the prognosis of cancer. Furthermore, the utility values used in our analysis are




Figure 2. The optimal age of last surveillance for men (A) and women (B) with NDBE. *The error bars present the ranges of surveillance stopping ages between models. The large error bars for women are the result of differences in natural history assumptions between the models. At older ages, the cumulative incidence of EAC in the EACMo is lower than the other 2 models, resulting in an earlier stopping age for women compared with the other 2 models (see Discussion section for more details on model differences).

derived from limited available literature that may not accurately represent the value or quality of patients' lives.

Despite these limitations, our findings have many strengths. The 3 independent models were developed under the auspices of the National Cancer Institute CISNET modeling consortium during the past 10 years with regular meetings lending support to the credibility and prior validation of the models and the comparative modeling process. The largest limitation in simulation modeling is the uncertainty in both model parameter estimates and structure. Our analysis used 3 models, which may provide some reassurance as opposed to the use of 1 model.

In addition, our results have important clinical implications for personalized management of patients with NDBE because none of the gastroenterology scientific societies recommend any stopping age for BE surveillance. For example, our results suggest that performing 1 more surveillance might not be appropriate from a cost-effectiveness perspective for a 76-year-old man with NDBE and severe comorbidity such as congestive heart failure. However, a 76-year-old man without comorbidity may be considered for 1 more surveillance at that age. It is worth mentioning that in addition to monetary costs, surveillance itself can become harmful and preclude increases in QALYs. For example, surveillance of an 85-year-old woman with NDBE and severe comorbidity can result in QALY loss. Empirical evidence has demonstrated that advancing age and more severe comorbidity have very little effect on the decision of whether to perform surveillance endoscopy in Medicare patients with BE.³¹

Our study was conducted in the US setting, but our findings can be applied to other settings with similarly high incidences of BE and EAC, such as countries in Northern and Western Europe and Oceania, and can inform international guidelines on the optimal age for last surveillance of patients with NDBE.

In conclusion, our comparative modeling approach shows that, in addition to chronological age, sex and the comorbidity status of patients with NDBE are important factors to inform the decision when to discontinue surveillance. Our analysis finds that the optimal age for last surveillance of patients with NDBE without comorbidity for women is 75 years and for men is 81 years. However, it may be up to 6 years earlier for women and up to 8 years earlier for men if patients have severe comorbidities.

Supplementary Material

Note: To access the supplementary material accompanying this article, visit the online version of *Gastroenterology* at www.gastrojournal.org, and at <http://doi.org/10.1053/j.gastro.2021.05.003>.

References

1. Rustgi AK, El-Serag HB. Esophageal carcinoma. *N Engl J Med* 2014;371:2499–2509.
2. Thrift AP. Barrett's esophagus and esophageal adenocarcinoma: how common are they really? *Dig Dis Sci* 2018;63:1988–1996.
3. Standards of Practice Committee, C, Wani S, Qumseya B, et al. Endoscopic eradication therapy for patients with Barrett's esophagus-associated dysplasia and intramucosal cancer. *Gastrointest Endosc* 2018; 87:907–931.e9.
4. Shaheen NJ, Falk GW, Iyer PG, et al. ACG clinical guideline: diagnosis and management of Barrett's esophagus. *Am J Gastroenterol* 2016;111:30–50; quiz 51.
5. American Gastroenterological Association, Spechler SJ, Sharma P, et al. American Gastroenterological Association medical position statement on the management of Barrett's esophagus. *Gastroenterology* 2011;140:1084–1091.
6. Fitzgerald RC, di Pietro M, Ragunath K, et al. British Society of Gastroenterology guidelines on the diagnosis and management of Barrett's oesophagus. *Gut* 2014; 63:7–42.
7. Cho H, Klabunde CN, Yabroff KR, et al. Comorbidity-adjusted life expectancy: a new tool to inform recommendations for optimal screening strategies. *Ann Intern Med* 2013;159:667–676.
8. Lansdorp-Vogelaar I, Gulati R, Mariotto AB, et al. Personalizing age of cancer screening cessation based on comorbid conditions: model estimates of harms and benefits. *Ann Intern Med* 2014;161:104–112.
9. Kroep S, Lansdorp-Vogelaar I, Rubenstein JH, et al. An accurate cancer incidence in Barrett's esophagus: a best estimate using published data and modeling. *Gastroenterology* 2015;149:577–585.e4; quiz e14–e15.
10. NCI. Surveillance, Epidemiology, and End Results (SEER) program population (1969–2013). National Cancer Institute, DCCPS, Surveillance Research program, Surveillance Systems branch. Available at: <https://seer.cancer.gov/>. Accessed June 15, 2021.
11. CISNET esophagus cancer collaborators. Esophageal Cancer Model Profiles 2018. NIH Cancer Intervention and Surveillance Modeling Network (CISNET), 2018. Available at: <https://cisnet.cancer.gov/esophagus/profiles.html>. Accessed June 15, 2021.
12. Kong CY, Kroep S, Curtius K, et al. Exploring the recent trend in esophageal adenocarcinoma incidence and mortality using comparative simulation modeling. *Cancer Epidemiol Biomarkers Prev* 2014;23:997–1006.
13. Centers for Disease Control and Prevention. Underlying cause of death, 1999–2017. Available at: <https://wonder.cdc.gov/Deaths-by-Underlying-Cause.html>. Accessed June 15, 2021.
14. Wani S, Rubenstein JH, Vieth M, et al. Diagnosis and management of low-grade dysplasia in Barrett's esophagus: expert review from the Clinical Practice Updates Committee of the American Gastroenterological Association. *Gastroenterology* 2016;151:822–835.
15. Omidvari AH, Ali A, Hazelton WD, et al. Optimizing management of patients with Barrett's esophagus and low-grade or no dysplasia based on comparative modeling. *Clin Gastroenterol Hepatol* 2020;18:1961–1969.
16. 2015 GI Endoscopy Coding and Reimbursement Guide. Cook Medical, 2015. Available at: <https://www.cookmedical.com/>. Accessed June 15, 2021.

17. Hur C, Choi SE, Rubenstein JH, et al. The cost effectiveness of radiofrequency ablation for Barrett's esophagus. *Gastroenterology* 2012;143:567–575.
18. Cryer BL, Wilcox CM, Henk HJ, et al. The economics of upper gastrointestinal bleeding in a US managed-care setting: a retrospective, claims-based analysis. *J Med Econ* 2010;13:70–77.
19. Yabroff KR, Lamont EB, Mariotto A, et al. Cost of care for elderly cancer patients in the United States. *J Natl Cancer Inst* 2008;100:630–641.
20. Kroep S, Heberle CR, Curtius K, et al. Radiofrequency ablation of Barrett's esophagus reduces esophageal adenocarcinoma incidence and mortality in a comparative modeling analysis. *Clin Gastroenterol Hepatol* 2017;15:1471–1474.
21. Hur C, Nishioka NS, Gazelle GS. Cost-effectiveness of aspirin chemoprevention for Barrett's esophagus. *J Natl Cancer Inst* 2004;96:316–325.
22. de Boer AG, Stalmeier PF, Sprangers MA, et al. Transhiatal vs extended transthoracic resection in oesophageal carcinoma: patients' utilities and treatment preferences. *Br J Cancer* 2002;86:851–857.
23. Garside R, Pitt M, Somerville M, et al. Surveillance of Barrett's oesophagus: exploring the uncertainty through systematic review, expert workshop and economic modelling. *Health Technol Assess* 2006;10:1–142; iii–iv.
24. Sanders GD, Neumann PJ, Basu A, et al. Recommendations for conduct, methodological practices, and reporting of cost-effectiveness analyses: second panel on cost-effectiveness in health and medicine. *JAMA* 2016;316:1093–1103.
25. Gordon LG, Mayne GC, Hirst NG, et al. Cost-effectiveness of endoscopic surveillance of non-dysplastic Barrett's esophagus. *Gastrointest Endosc* 2014;79:242–256.e6.
26. Kastelein F, van Olphen S, Steyerberg EW, et al. Surveillance in patients with long-segment Barrett's oesophagus: a cost-effectiveness analysis. *Gut* 2015;64:864–871.
27. van Hees F, Saini SD, Lansdorp-Vogelaar I, et al. Personalizing colonoscopy screening for elderly individuals based on screening history, cancer risk, and comorbidity status could increase cost effectiveness. *Gastroenterology* 2015;149:1425–1437.
28. Runge TM, Abrams JA, Shaheen NJ. Epidemiology of Barrett's esophagus and esophageal adenocarcinoma. *Gastroenterol Clin North Am* 2015;44:203–231.
29. Gatenby P, Caygill C, Wall C, et al. Lifetime risk of esophageal adenocarcinoma in patients with Barrett's esophagus. *World J Gastroenterol* 2014;20:9611–9617.
30. Rex DK, Cummings OW, Shaw M, et al. Screening for Barrett's esophagus in colonoscopy patients with and without heartburn. *Gastroenterology* 2003;125:1670–1677.
31. Rubenstein JH, Noureldin M, Tavakkoli A, et al. Utilization of surveillance endoscopy for Barrett's esophagus in Medicare enrollees. *Gastroenterology* 2020;158:773–775.e1.

Received September 20, 2020. Accepted May 1, 2021.

Correspondence

Address correspondence to: Amir-Houshang Omidvari, MD, PhD, PO Box 2040, 3000 CA Rotterdam, The Netherlands. e-mail: a.omidvari@erasmusmc.nl

Acknowledgment

The authors thank Dr Amy Knudsen, of the Institute for Technology Assessment, Massachusetts General Hospital, for her help on adjustment of life tables.

CRedit Authorship Contributions

Amir-Houshang Omidvari, MD, PhD (Conceptualization: Equal; Formal analysis: Lead; Project administration: Lead; Validation: Equal; Writing – original draft: Lead).

William D. Hazelton, PhD (Formal analysis: Equal; Validation: Equal; Writing – review & editing: Equal).

Brianna N. Lauren, BS (Formal analysis: Equal; Project administration: Equal; Validation: Equal; Writing – review & editing: Equal).

Steffie K. Naber, PhD (Formal analysis: Supporting; Supervision: Supporting; Visualization: Equal; Writing – review & editing: Equal).

Minyi Lee, BS (Formal analysis: Equal; Writing – review & editing: Equal).

Ayman Ali, BS (Formal analysis: Equal; Writing – review & editing: Equal).

Claudia Seguin, BS (Formal analysis: Equal; Writing – review & editing: Equal).

Chun Yin Kong, PhD (Conceptualization: Equal; Formal analysis: Supporting; Writing – review & editing: Equal).

Ellen Richmond, MS (Formal analysis: Supporting; Resources: Supporting; Writing – review & editing: Equal).

Joel H. Rubenstein, MD, MSc (Conceptualization: Equal; Formal analysis: Supporting; Writing – review & editing: Equal).

Georg E. Luebeck, PhD (Formal analysis: Supporting; Funding acquisition: Equal; Supervision: Equal; Writing – review & editing: Equal).

John M. Inadomi, MD (Conceptualization: Equal; Formal analysis: Supporting; Funding acquisition: Equal; Supervision: Equal; Writing – review & editing: Equal).

Chin Hur, MD, MPH (Conceptualization: Equal; Formal analysis: Supporting; Funding acquisition: Equal; Supervision: Equal; Writing – review & editing: Equal).

Iris Lansdorp-Vogelaar, PhD (Conceptualization: Lead; Formal analysis: Supporting; Funding acquisition: Equal; Supervision: Equal; Writing – review & editing: Equal).

Conflicts of interest

All authors received funding from the National Institutes of Health/National Cancer Institute for conducting this study.

Funding

This study has been supported by the National Institutes of Health/National Cancer Institute (grant numbers U01CA152926 and U01CA199336). The authors also thank the Scientific Computing Infrastructure at Fred Hutchinson Cancer Research Center funded by Office of Research Infrastructure Programs grant S10OD028685.

Supplementary Table 1. Post-Treatment (EET or Touch-Up RFA) Surveillance Strategies

Patients characteristics and treatment outcome	Strategy
Recurrent NDBE patient (initial LGD)	
After CE-IM (state = normal)	Surveillance at 6, 12, 24, and 36 mo, then every 3 y
After non-CE-IM (state = ND)	
Recurrent NDBE patient (Initial HGD)	
After CE-IM (state = normal)	Surveillance at 3, 6, 9, 12, 18, and 24 mo, then every 1 y
After non-CE-IM (state = ND)	
Initial or recurrent LGD patients	
After CE-IM (state = normal)	Surveillance at 12 and 24 mo, then every 3 y
After CE-D, non-CE-IM (state = ND)	Surveillance at 6, 12, 24, and 36 mo, then every 3 y
After non-CE-D, non-CE-IM (state = LGD)	Surveillance at 6 and 12 mo, then every 1 y
Initial or recurrent HGD patients	
After CE-IM (state = normal)	Surveillance at 3, 6, 9, 12, 18, and 24 mo, then every 1 y
After CE-D, non-CE-IM (state = ND)	
After non-CE-D, non-CE-IM (state = LGD/HGD)	Surveillance every 3 mo

CE, complete eradication; D, dysplasia; IM, intestinal metaplasia; ND, no dysplasia.

Supplementary Table 2. Model Inputs

Parameter/definition	Value	Source
Maximum number of touch-ups RFA	3	1
Duration of initial EET	2 y	1
No. of endoscopies during initial EET	4	1
No. of RFA sessions during initial EET	3.55	1
Proportion of patients receiving EMR treatments before RFA	0.55	2
Complication rates		
Perforation due to surveillance endoscopy	0.00025	3-7
Bleeding due to surveillance endoscopy	0.00026	3-6
Perforation due to EET (per procedure)	0.002 ^a	8
Bleeding due to EET (per procedure)	0.004 ^a	8,9
Stricture rate due to EET (per procedure)	0.019 ^a	8,9
Perforation rate resulting from stricture treatment	0.0009	10
Bleeding rate resulting from stricture treatment	0.0009	10
Success probabilities of EET ^b		
In HGD patients		
CE-IM and CE-D	88.9%	11
Non-CE-IM, CE-D	3.7%	11
Non-CE-IM and non-CE-D	7.4%	11
In LGD patients		
CE-IM and CE-D	98.1%	11
Non-CE-IM, CE-D	0	11
Non-CE-IM and non-CE-D	1.9%	11
Recurrence rates by baseline histologic grade and grade of recurrence		
Annual recurrence rates after CE-IM		
Pretreatment misdiagnosed NDBE ^b	7%	12,13
Pretreatment IND/LGD	8.3%	11
Pretreatment HGD	13.5%	11
Recurrent histology of misdiagnosed NDBE ^b after CE-IM		
NDBE	92%	12,13
IND/LGD	6%	12,13
HGD	2%	12,13
EAC	0%	12,13
Recurrent histology of IND/LGD after CE-IM		
NDBE	50%	11
IND/LGD	25%	11
HGD	25%	11
EAC	0	11

Supplementary Table 2. Continued

Parameter/definition	Value	Source
Recurrent histology of HGD after CE-IM		
NDBE	50%	11-13
IND/LGD	15%	11
HGD	25%	11
EAC	10%	11
Costs		
Endoscopy	\$745	14
Initial EET treatment (EMR and RFA)	\$5,630	14
RFA touch-up	\$1,012	14
Complications		
Stricture	\$1,012	14
Bleeding	\$11,815	15
Perforation	\$28,533	16
Localized EAC initial care	\$58,997	17
Localized EAC Terminal care	\$64,704	17
Regional EAC initial care	\$75,295	17
Regional EAC terminal care	\$77,742	17
Distant EAC initial care	\$57,169	17
Distant EAC terminal care	\$85,212	17
Unstaged ^c EAC initial care	\$63,820	17
Unstaged ^c EAC terminal care	\$75,886	17
EAC continuous care	\$4,080	17
Utility		
Short term		
Endoscopy with or without EET (1 d)	0.70	18
After EET treatment (1 wk) ^d	0.70	1
After RFA touch-up (1 wk)	0.70	1
Stricture (4 wk)	0.70	1, expert opinion
Perforation (4 wk)	0.70	1, expert opinion
Bleeding (1 wk)	0.70	1
Long term (until death)		
Localized EAC initial care (yearly)	0.84	19,20
Localized EAC continuous and terminal care (yearly)	0.96	19,20
Regional EAC care (yearly)	0.65	19,20
Distant EAC care (yearly)	0.40	19,20
Unstaged ^c EAC care (yearly)	0.63	19,20

EMR, endoscopic mucosal resection; IND, indefinite dysplasia.
^aThe complication rate per patient due to RFA was adjusted to average RFA sessions to compute the complication rate per procedure.⁹

^bRecurrent NDBE patients or NDBE patients who are misdiagnosed as having LGD/HGD receive EET as well. For NDBE, we assumed the same EET success probability rate that we assumed for LGD patients, but we assumed a different recurrence rate after CE-IM as described in the table.

^cUnknown.

^dPatients were assumed to receive an average of 3.55 RFA sessions and 0.55 EMR treatments in the first 2 y after initial EET, therefore (3.55 + 0.55 =) 4.1 weeks with utility of 0.7.

Supplementary Table 3. Life-Time Net Benefits and Costs of 1 Additional Endoscopic Surveillance at Ages 68, 74, 80, and 86 Vs Not Performing Surveillance at That Age Per 1000 Women With NDBE Diagnosed at Age 60

Age	Comorbidity level ^a	EAC prevented	EAC death prevented	QALYs gained	Net endoscopies	Net EET and touch-ups	Net cost (\$)	ICER
68	No	5	7	34	1956	75	1,510,218	44,873
	Mild	5	6	28	1885	73	1,502,862	54,376
	Moderate	4	6	25	1861	72	1,498,669	59,010
	Severe	3	4	19	1753	68	1,468,876	78,148
74	No	3	5	18	1699	66	1,387,573	79,020
	Mild	3	4	13	1631	64	1,371,970	102,016
	Moderate	2	3	12	1603	63	1,365,868	116,662
	Severe	2	3	8	1500	60	1,325,333	168,689
80	No	2	3	7	1447	58	1,265,942	181,671
	Mild	1	2	5	1399	56	1,248,497	248,183
	Moderate	1	2	4	1356	55	1,235,250	336,392
	Severe	1	1	2	1265	53	1,196,610	737,790
86	No	1	1	1	1202	52	1,131,114	778,692
	Mild	1	1	1	1180	52	1,123,488	1,171,891
	Moderate	0	1	0	1134	51	1,102,231	5,010,660
	Severe	0	1	-1	1049	49	1,054,641	Dominated

^aOverview of comorbidity levels and associated conditions can be found in [Table 1](#) of the main text.

Supplementary References

1. Kroep S, Heberle CR, Curtius K, et al. Radiofrequency ablation of Barrett's esophagus reduces esophageal adenocarcinoma incidence and mortality in a comparative modeling analysis. *Clin Gastroenterol Hepatol* 2017; 15:1471–1474.
2. Gupta M, Iyer PG, Lutzke L, et al. Recurrence of esophageal intestinal metaplasia after endoscopic mucosal resection and radiofrequency ablation of Barrett's esophagus: results from a US Multicenter Consortium. *Gastroenterology* 2013;145:79–86.e1.
3. Silvis SE, Nebel O, Rogers G, et al. Endoscopic complications. Results of the 1974 American Society for Gastrointestinal Endoscopy Survey. *JAMA* 1976; 235:928–930.
4. Sieg A, Hachmoeller-Eisenbach U, Eisenbach T. Prospective evaluation of complications in outpatient GI endoscopy: a survey among German gastroenterologists. *Gastrointest Endosc* 2001;53:620–627.
5. Quine MA, Bell GD, McCloy RF, et al. Prospective audit of upper gastrointestinal endoscopy in two regions of England: safety, staffing, and sedation methods. *Gut* 1995;36:462–467.
6. Davis RE, Graham DY. Endoscopic complications: the Texas experience. *Gastrointest Endosc* 1979;25:146–149.
7. Dawson J, Cockel R. Oesophageal perforation at fiberoptic gastroscopy. *Br Med J (Clin Res Ed)* 1981; 283:583.
8. Qumseya BJ, Wani S, Desai M, et al. Adverse events after radiofrequency ablation in patients with Barrett's esophagus: a systematic review and meta-analysis. *Clin Gastroenterol Hepatol* 2016;14:1086–1095.e6.
9. Orman ES, Li N, Shaheen NJ. Efficacy and durability of radiofrequency ablation for Barrett's esophagus: systematic review and meta-analysis. *Clin Gastroenterol Hepatol* 2013;11:1245–1255.
10. Piotet E, Escher A, Monnier P. Esophageal and pharyngeal strictures: report on 1,862 endoscopic dilatations using the Savary-Gilliard technique. *Eur Arch Otorhinolaryngol* 2008;265:357–364.
11. Cotton CC, Wolf WA, Overholt BF, et al. Late recurrence of Barrett's esophagus after complete eradication of intestinal metaplasia is rare: final report from ablation in intestinal metaplasia containing dysplasia trial. *Gastroenterology* 2017;153:681–688.e2.
12. Asher Wolf W, Overholt B, Nan L, et al. Durability of radiofrequency ablation (RFA) in Barrett's esophagus with dysplasia: the AIM Dysplasia Trial at five years. *Gastroenterology* 2014;146:S131.
13. Asher Wolf W, Pruitt R, Ertan A, et al. Predictors of esophageal adenocarcinoma in patients with prior radiofrequency ablation (RFA) for treatment of Barrett's esophagus: results from the U.S. RFA Registry. *Gastrointest Endosc* 2014;79:AB217.
14. Arnold M, Soerjomataram I, Ferlay J, et al. Global incidence of oesophageal cancer by histological subtype in 2012. *Gut* 2015;64:381–387.
15. Cryer BL, Wilcox CM, Henk HJ, et al. The economics of upper gastrointestinal bleeding in a US managed-care setting: a retrospective, claims-based analysis. *J Med Econ* 2010;13:70–77.
16. Hur C, Choi SE, Rubenstein JH, et al. The cost effectiveness of radiofrequency ablation for Barrett's esophagus. *Gastroenterology* 2012;143:567–575.
17. Yabroff KR, Lamont EB, Mariotto A, et al. Cost of care for elderly cancer patients in the United States. *J Natl Cancer Inst* 2008;100:630–641.
18. Hur C, Nishioka NS, Gazelle GS. Cost-effectiveness of aspirin chemoprevention for Barrett's esophagus. *J Natl Cancer Inst* 2004;96:316–325.
19. de Boer AG, Stalmeier PF, Sprangers MA, et al. Transhiatal vs extended transthoracic resection in oesophageal carcinoma: patients' utilities and treatment preferences. *Br J Cancer* 2002;86:851–857.
20. Garside R, Pitt M, Somerville M, et al. Surveillance of Barrett's oesophagus: exploring the uncertainty through systematic review, expert workshop and economic modelling. *Health Technol Assess* 2006;10:1–142; iii–iv.