

Extra twisted connected sums and their ν -invariants


Sebastian Goette

Mathematisches Institut der Universität Freiburg


SCSHGAP Second Annual Meeting
September 13–14, 2018

- ▶ G_2 -geometry
Intro, properties, examples, questions
- ▶ The ν -invariant
Differential topology, definition of ν , properties, first examples
- ▶ Extra twisted connected sums
Construction, properties, problems
- ▶ Computation of the ν -invariant
Computations with η -invariants, examples, questions


Consider parallel translation along a spherical triangle




Consider parallel translation along a spherical triangle



Consider parallel translation along a spherical triangle




Consider parallel translation along a spherical triangle



Consider parallel translation along a spherical triangle

A vector is rotated by an angle equal to the spherical area of the triangle
([Gauß-Bonnet theorem](#))




Consider parallel translation along a spherical triangle

A vector is rotated by an angle equal to the spherical area of the triangle
(Gauß-Bonnet theorem)

The subgroup of $\text{Aut}(T_p S^2)$ of all these parallel translations is the **holonomy group**

$$\text{Hol}(S^2, g^{\text{rnd}}) \cong SO(2)$$



Consider parallel translation along a spherical triangle


A vector is rotated by an angle equal to the spherical area of the triangle
([Gauß-Bonnet theorem](#))

The subgroup of $\text{Aut}(T_p S^2)$ of all these parallel translations is the **holonomy group**

$$\text{Hol}(S^2, g^{\text{rnd}}) \cong SO(2)$$

Related: the spherical area formula

$$A(\Delta) = \alpha + \beta + \gamma - \pi .$$



Theorem (Berger)

The only possible holonomy groups of complete, simply connected Riemannian manifolds that are neither a product nor a symmetric space are

Theorem (Berger)

The only possible holonomy groups of complete, simply connected Riemannian manifolds that are neither a product nor a symmetric space are

<i>Holonomy group</i>	<i>dim</i>	<i>ric</i>	<i>Structure</i>	<i>Parallel spinors</i>	<i>Name</i>
$SO(n)$	n				<i>generic case</i>
$U(k)$	$2k$		J		<i>Kähler</i>
$SU(k)$	$2k$	0	J, Ω	2	<i>Calabi-Yau</i>
$Sp(\ell) \cdot Sp(1)$	4ℓ	const	$\langle I, J, K \rangle$		<i>Quat. Kähler</i>
$Sp(\ell)$	4ℓ	0	I, J, K, Ω	$\ell + 1$	<i>hyper Kähler</i>
G_2	7	0	$\varphi \in \Omega^3$	1	<i>exceptional</i>
$Spin(7)$	8	0	$\psi \in \Omega^4$	1	<i>exceptional</i>

Theorem (Berger)

The only possible holonomy groups of complete, simply connected Riemannian manifolds that are neither a product nor a symmetric space are

<i>Holonomy group</i>	<i>dim</i>	<i>ric</i>	<i>Structure</i>	<i>Parallel spinors</i>	<i>Name</i>
$SO(n)$	n				<i>generic case</i>
$U(k)$	$2k$		J		<i>Kähler</i>
$SU(k)$	$2k$	0	J, Ω	2	<i>Calabi-Yau</i>
$Sp(\ell) \cdot Sp(1)$	4ℓ	const	$\langle I, J, K \rangle$		<i>Quat. Kähler</i>
$Sp(\ell)$	4ℓ	0	I, J, K, Ω	$\ell + 1$	<i>hyper Kähler</i>
G_2	7	0	$\varphi \in \Omega^3$	1	<i>exceptional</i>
$Spin(7)$	8	0	$\psi \in \Omega^4$	1	<i>exceptional</i>

Why consider G_2 -manifolds?

Why consider G_2 -manifolds?

Mathematical motivation

- ▶ Only special holonomy group for odd dimensional manifolds
- ▶ Only G_2 and $\text{Spin}(7)$ holonomy have no direct relation to algebraic geometry

Hence, new methods are needed

Why consider G_2 -manifolds?

Mathematical motivation

- ▶ Only special holonomy group for odd dimensional manifolds
- ▶ Only G_2 and $\text{Spin}(7)$ holonomy have no direct relation to algebraic geometry

Hence, new methods are needed

Physical motivation

- ▶ In string theory, spacetime takes the form $\mathbb{R}^{3,1} \times V$, where V is Calabi-Yau
- ▶ In **M-theory**, spacetime takes the form $\mathbb{R}^{3,1} \times M$, where M is a G_2 -manifold
- ▶ Possible relations to other physical theories

Hence, many fruitful interactions possible

Characterisations of the Lie group G_2 give characterisations of G_2 -manifolds

▶ $G_2 = \text{Aut}(\mathbb{O})$

Hence G_2 preserves $\varphi = \langle \cdot \times \cdot, \cdot \rangle \in \Lambda^3 \text{Im } \mathbb{O} \cong \Lambda^3 \mathbb{R}^7$

Characterisations of the Lie group G_2 give characterisations of G_2 -manifolds

- ▶ $G_2 = \text{Aut}(\mathbb{O})$

Hence G_2 preserves $\varphi = \langle \cdot \times \cdot, \cdot \rangle \in \Lambda^3 \text{Im } \mathbb{O} \cong \Lambda^3 \mathbb{R}^7$

- ▶ The stabiliser of φ in $GL(7, \mathbb{R})$ is G_2

The $GL(7, \mathbb{R})$ -orbit of $\varphi \in \Lambda^3 \mathbb{R}^7$ is open (not dense)

Forms in this orbit are called **positive**

Characterisations of the Lie group G_2 give characterisations of G_2 -manifolds

▶ $G_2 = \text{Aut}(\mathbb{O})$

Hence G_2 preserves $\varphi = \langle \cdot \times \cdot, \cdot \rangle \in \Lambda^3 \text{Im } \mathbb{O} \cong \Lambda^3 \mathbb{R}^7$

▶ The stabiliser of φ in $GL(7, \mathbb{R})$ is G_2

The $GL(7, \mathbb{R})$ -orbit of $\varphi \in \Lambda^3 \mathbb{R}^7$ is open (not dense)

Forms in this orbit are called **positive**

▶ A positive 3-form on M determines a G_2 -structure and a metric g_φ

Call φ **torsion free** if $d\varphi = d_{g_\varphi}^* \varphi = 0$ (nonlinear condition)

Then (M, g_φ) has $\text{Hol}(M, g_\varphi) \subset G_2$ if and only if φ is torsion-free

Characterisations of the Lie group G_2 give characterisations of G_2 -manifolds

▶ $G_2 = \text{Aut}(\mathbb{O})$

Hence G_2 preserves $\varphi = \langle \cdot \times \cdot, \cdot \rangle \in \Lambda^3 \text{Im } \mathbb{O} \cong \Lambda^3 \mathbb{R}^7$ and $1 \in \mathbb{R} \subset \mathbb{O}$

▶ The stabiliser of φ in $GL(7, \mathbb{R})$ is G_2

The $GL(7, \mathbb{R})$ -orbit of $\varphi \in \Lambda^3 \mathbb{R}^7$ is open (not dense)

Forms in this orbit are called **positive**

▶ A positive 3-form on M determines a G_2 -structure and a metric g_φ

Call φ **torsion free** if $d\varphi = d_{g_\varphi}^* \varphi = 0$ (nonlinear condition)

Then (M, g_φ) has $\text{Hol}(M, g_\varphi) \subset G_2$ if and only if φ is torsion-free

▶ G_2 is compact, simply connected, so G_2 -manifolds are Riemannian and spin

The stabiliser of a nonzero spinor in $\text{Spin}(7)$ is G_2

A Riemannian 7-manifold (M, g) has $\text{Hol}(M, g) \subset G_2$

if and only if it is spin and there exists a nonzero **parallel spinor**

Let $\varphi \in \Omega^3(M)$ be positive and closed

There exists a bilinear form B on $H^2(M; \mathbb{R})$ given for closed forms $\alpha, \beta \in \Omega^2(M)$ by

$$B([\alpha], [\beta]) = ([\alpha] \smile [\beta] \smile [\varphi])[M] = \int_M \alpha \wedge \beta \wedge \varphi$$

Let $p_1(TM) \in H^4(M; \mathbb{R})$ denote the first Pontryagin class

¹Spiro Karigiannis pointed out after the talk that another requirement is almost formality, see recent work by Chan, Karigiannis and Tsang, arXiv:1801.06410

Let $\varphi \in \Omega^3(M)$ be positive and closed

There exists a bilinear form B on $H^2(M; \mathbb{R})$ given for closed forms $\alpha, \beta \in \Omega^2(M)$ by

$$B([\alpha], [\beta]) = ([\alpha] \smile [\beta] \smile [\varphi])[M] = \int_M \alpha \wedge \beta \wedge \varphi$$

Let $p_1(TM) \in H^4(M; \mathbb{R})$ denote the first Pontryagin class

If M is closed and $\text{Hol}(M, \varphi) \subset G_2$ then

- ▶ M is **oriented** and **spin** and $b_3(M) = \dim H^3(M; \mathbb{R}) \geq 1$

¹Spiro Karigiannis pointed out after the talk that another requirement is almost formality, see recent work by Chan, Karigiannis and Tsang, arXiv:1801.06410

Let $\varphi \in \Omega^3(M)$ be positive and closed

There exists a bilinear form B on $H^2(M; \mathbb{R})$ given for closed forms $\alpha, \beta \in \Omega^2(M)$ by

$$B([\alpha], [\beta]) = ([\alpha] \smile [\beta] \smile [\varphi])[M] = \int_M \alpha \wedge \beta \wedge \varphi$$

Let $p_1(TM) \in H^4(M; \mathbb{R})$ denote the first Pontryagin class

If M is closed and $\text{Hol}(M, \varphi) \subset G_2$ then

- ▶ M is **oriented** and **spin** and $b_3(M) = \dim H^3(M; \mathbb{R}) \geq 1$
- ▶ $\text{Hol}(M, g) = G_2 \iff \#\pi_1(M) < \infty$

¹Spiro Karigiannis pointed out after the talk that another requirement is almost formality, see recent work by Chan, Karigiannis and Tsang, arXiv:1801.06410

Let $\varphi \in \Omega^3(M)$ be positive and closed

There exists a bilinear form B on $H^2(M; \mathbb{R})$ given for closed forms $\alpha, \beta \in \Omega^2(M)$ by

$$B([\alpha], [\beta]) = ([\alpha] \smile [\beta] \smile [\varphi])[M] = \int_M \alpha \wedge \beta \wedge \varphi$$

Let $p_1(TM) \in H^4(M; \mathbb{R})$ denote the first Pontryagin class

If M is closed and $\text{Hol}(M, \varphi) \subset G_2$ then

- ▶ M is **oriented** and **spin** and $b_3(M) = \dim H^3(M; \mathbb{R}) \geq 1$
- ▶ $\text{Hol}(M, g) = G_2 \iff \#\pi_1(M) < \infty$
- ▶ $\text{Hol}(M, g) = G_2 \implies B$ is negative definite

¹Spiro Karigiannis pointed out after the talk that another requirement is almost formality, see recent work by Chan, Karigiannis and Tsang, arXiv:1801.06410

Let $\varphi \in \Omega^3(M)$ be positive and closed

There exists a bilinear form B on $H^2(M; \mathbb{R})$ given for closed forms $\alpha, \beta \in \Omega^2(M)$ by

$$B([\alpha], [\beta]) = ([\alpha] \smile [\beta] \smile [\varphi])[M] = \int_M \alpha \wedge \beta \wedge \varphi$$

Let $p_1(TM) \in H^4(M; \mathbb{R})$ denote the first Pontryagin class

If M is closed and $\text{Hol}(M, \varphi) \subset G_2$ then

- ▶ M is **oriented** and **spin** and $b_3(M) = \dim H^3(M; \mathbb{R}) \geq 1$
- ▶ $\text{Hol}(M, g) = G_2 \iff \#\pi_1(M) < \infty$
- ▶ $\text{Hol}(M, g) = G_2 \implies B$ is negative definite
- ▶ $\text{Hol}(M, g) = G_2 \implies (p_1(TM) \smile [\varphi])[M] < 0$

¹Spiro Karigiannis pointed out after the talk that another requirement is almost formality, see recent work by Chan, Karigiannis and Tsang, arXiv:1801.06410

Let $\varphi \in \Omega^3(M)$ be positive and closed

There exists a bilinear form B on $H^2(M; \mathbb{R})$ given for closed forms $\alpha, \beta \in \Omega^2(M)$ by

$$B([\alpha], [\beta]) = ([\alpha] \smile [\beta] \smile [\varphi])[M] = \int_M \alpha \wedge \beta \wedge \varphi$$

Let $p_1(TM) \in H^4(M; \mathbb{R})$ denote the first Pontryagin class

If M is closed and $\text{Hol}(M, \varphi) \subset G_2$ then

- ▶ M is **oriented** and **spin** and $b_3(M) = \dim H^3(M; \mathbb{R}) \geq 1$
- ▶ $\text{Hol}(M, g) = G_2 \iff \#\pi_1(M) < \infty$
- ▶ $\text{Hol}(M, g) = G_2 \implies B$ is negative definite
- ▶ $\text{Hol}(M, g) = G_2 \implies (p_1(TM) \smile [\varphi])[M] < 0$

These are all **known**¹ obstructions against holonomy G_2

¹Spiro Karigiannis pointed out after the talk that another requirement is almost formality, see recent work by Chan, Karigiannis and Tsang, arXiv:1801.06410

Let M be a compact oriented spin 7-manifold and define

$$\mathcal{X} = \{ \varphi \in \Omega^3(M) \mid \varphi \text{ is positive and torsion free} \}$$

Let M be a compact oriented spin 7-manifold and define

$$\mathcal{X} = \{ \varphi \in \Omega^3(M) \mid \varphi \text{ is positive and torsion free} \}$$

Let $\mathcal{D} \subset \text{Diff}(M)$ be the connected component of id_M . Then

$$\mathcal{M} = \mathcal{X}/\mathcal{D}$$

is called the G_2 -moduli space of M

Let M be a compact oriented spin 7-manifold and define

$$\mathcal{X} = \{ \varphi \in \Omega^3(M) \mid \varphi \text{ is positive and torsion free} \}$$

Let $\mathcal{D} \subset \text{Diff}(M)$ be the connected component of id_M . Then

$$\mathcal{M} = \mathcal{X}/\mathcal{D}$$

is called the G_2 -moduli space of M

Theorem (Joyce)

The G_2 -moduli space is a manifold, and the map

$$\mathcal{M} \longrightarrow H^3(M; \mathbb{R}) \quad \text{with} \quad [\varphi] \longmapsto [\varphi]$$

*is a **local** diffeomorphism*

Let M be a compact oriented spin 7-manifold and define

$$\mathcal{X} = \{ \varphi \in \Omega^3(M) \mid \varphi \text{ is positive and torsion free} \}$$

Let $\mathcal{D} \subset \text{Diff}(M)$ be the connected component of id_M . Then

$$\mathcal{M} = \mathcal{X}/\mathcal{D}$$

is called the G_2 -moduli space of M

Theorem (Joyce)

The G_2 -moduli space is a manifold, and the map

$$\mathcal{M} \longrightarrow H^3(M; \mathbb{R}) \quad \text{with} \quad [\varphi] \longmapsto [\varphi]$$

is a local diffeomorphism

Not much is known about the global structure of \mathcal{M}

To construct subsets of the G_2 -moduli space means to construct deformation families of G_2 -manifolds first

To construct subsets of the G_2 -moduli space means to construct deformation families of G_2 -manifolds first

- ▶ Bryant '87: first **non-complete** examples
- ▶ Bryant and Salamon '89: first **complete** examples
- ▶ Joyce '96: first **closed** examples

To construct subsets of the G_2 -moduli space means to construct deformation families of G_2 -manifolds first

- ▶ Bryant '87: first **non-complete** examples
- ▶ Bryant and Salamon '89: first **complete** examples
- ▶ Joyce '96: first **closed** examples

Joyce's construction: let a “rich enough” finite subgroup $\Gamma \subset G_2$ act on **flat** T^7 with “sufficiently many” fixpoints, preserving a parallel positive 3-form $\varphi \in \Omega^3(T^7)$

To construct subsets of the G_2 -moduli space means to construct deformation families of G_2 -manifolds first

- ▶ Bryant '87: first **non-complete** examples
- ▶ Bryant and Salamon '89: first **complete** examples
- ▶ Joyce '96: first **closed** examples

Joyce's construction: let a “rich enough” finite subgroup $\Gamma \subset G_2$ act on **flat** T^7 with “sufficiently many” fixpoints, preserving a parallel positive 3-form $\varphi \in \Omega^3(T^7)$
The stabilisers of fixpoints $p \in T^7$ are isomorphic to subgroups of $SU(2)$ or $SU(3)$

To construct subsets of the G_2 -moduli space means to construct deformation families of G_2 -manifolds first

- ▶ Bryant '87: first **non-complete** examples
- ▶ Bryant and Salamon '89: first **complete** examples
- ▶ Joyce '96: first **closed** examples

Joyce's construction: let a “rich enough” finite subgroup $\Gamma \subset G_2$ act on **flat** T^7 with “sufficiently many” fixpoints, preserving a parallel positive 3-form $\varphi \in \Omega^3(T^7)$
The stabilisers of fixpoints $p \in T^7$ are isomorphic to subgroups of $SU(2)$ or $SU(3)$
By gluing in suitable bundles of **noncompact Calabi-Yau manifolds** in place of the singularities, Joyce constructs a desingularisation

$$M \longrightarrow T^7/\Gamma$$

To construct subsets of the G_2 -moduli space means to construct deformation families of G_2 -manifolds first

- ▶ Bryant '87: first **non-complete** examples
- ▶ Bryant and Salamon '89: first **complete** examples
- ▶ Joyce '96: first **closed** examples


Joyce's construction: let a “rich enough” finite subgroup $\Gamma \subset G_2$ act on **flat** T^7 with “sufficiently many” fixpoints, preserving a parallel positive 3-form $\varphi \in \Omega^3(T^7)$
The stabilisers of fixpoints $p \in T^7$ are isomorphic to subgroups of $SU(2)$ or $SU(3)$
By gluing in suitable bundles of **noncompact Calabi-Yau manifolds** in place of the singularities, Joyce constructs a desingularisation

$$M \longrightarrow T^7/\Gamma$$

The closed G_2 -structure on M obtained by gluing is close to a torsion-free one


- ▶ Kovalev '03, Corti-Haskins-Nordström-Pacini '15: **Twisted connected sums**

- ▶ Kovalev '03, Corti-Haskins-Nordström-Pacini '15: **Twisted connected sums**




Let V_+ , V_- be **asymptotically cylindrical Calabi-Yau threefolds**

- ▶ Kovalev '03, Corti-Haskins-Nordström-Pacini '15: **Twisted connected sums**




Let V_+ , V_- be asymptotically cylindrical Calabi-Yau threefolds with ends asymptotic to $\Sigma_{\pm} \times S^1 \times \mathbb{R}$, where Σ_{\pm} are **K3 surfaces**

- Kovalev '03, Corti-Haskins-Nordström-Pacini '15: **Twisted connected sums**



Let V_+ , V_- be asymptotically cylindrical Calabi-Yau threefolds with ends asymptotic to $\Sigma_{\pm} \times S^1 \times \mathbb{R}$, where Σ_{\pm} are K3 surfaces
 Glue $V_- \times S^1$ to $V_+ \times S^1$, flipping the circles

- ▶ Kovalev '03, Corti-Haskins-Nordström-Pacini '15: **Twisted connected sums**



Let V_+ , V_- be asymptotically cylindrical Calabi-Yau threefolds
with ends asymptotic to $\Sigma_{\pm} \times S^1 \times \mathbb{R}$, where Σ_{\pm} are K3 surfaces

Glue $V_- \times S^1$ to $V_+ \times S^1$, flipping the circles

The closed G_2 -structure on M obtained by gluing is close to a torsion-free one

Let us summarise first

- ▶ Only a few obstructions against G_2 -holonomy are known

Let us summarise first

- ▶ Only a few obstructions against G_2 -holonomy are known
- ▶ Only a few compact examples are known

Let us summarise first

- ▶ Only a few obstructions against G_2 -holonomy are known
- ▶ Only a few compact examples are known—only $\sim 10^8$ deformation families

Let us summarise first

- ▶ Only a few obstructions against G_2 -holonomy are known
- ▶ Only a few compact examples are known—only $\sim 10^8$ deformation families
- ▶ Known compact examples represent points close to the boundary of the moduli space

Let us summarise first

- ▶ Only a few obstructions against G_2 -holonomy are known
- ▶ Only a few compact examples are known—only $\sim 10^8$ deformation families
- ▶ Known compact examples represent points close to the boundary of the moduli space—
the G_2 -metric is locally close to metrics with holonomy groups $SU(2)$ or $SU(3)$

Let us summarise first

- ▶ Only a few obstructions against G_2 -holonomy are known
- ▶ Only a few compact examples are known—only $\sim 10^8$ deformation families
- ▶ Known compact examples represent points close to the boundary of the moduli space—
the G_2 -metric is locally close to metrics with holonomy groups $SU(2)$ or $SU(3)$

Important open problems / questions

- ▶ Find more invariants for / obstructions against G_2 -metrics

Let us summarise first

- ▶ Only a few obstructions against G_2 -holonomy are known
- ▶ Only a few compact examples are known—only $\sim 10^8$ deformation families
- ▶ Known compact examples represent points close to the boundary of the moduli space—
the G_2 -metric is locally close to metrics with holonomy groups $SU(2)$ or $SU(3)$

Important open problems / questions

- ▶ Find more invariants for / obstructions against G_2 -metrics
- ▶ Construct G_2 -metrics far away from the boundary of the moduli space

Let us summarise first

- ▶ Only a few obstructions against G_2 -holonomy are known
- ▶ Only a few compact examples are known—only $\sim 10^8$ deformation families
- ▶ Known compact examples represent points close to the boundary of the moduli space—
the G_2 -metric is locally close to metrics with holonomy groups $SU(2)$ or $SU(3)$

Important open problems / questions

- ▶ Find more invariants for / obstructions against G_2 -metrics
- ▶ Construct G_2 -metrics far away from the boundary of the moduli space
- ▶ How can families of G_2 -metrics become singular?

Let us summarise first

- ▶ Only a few obstructions against G_2 -holonomy are known
- ▶ Only a few compact examples are known—only $\sim 10^8$ deformation families
- ▶ Known compact examples represent points close to the boundary of the moduli space—
the G_2 -metric is locally close to metrics with holonomy groups $SU(2)$ or $SU(3)$

Important open problems / questions

- ▶ Find more invariants for / obstructions against G_2 -metrics
- ▶ Construct G_2 -metrics far away from the boundary of the moduli space
- ▶ How can families of G_2 -metrics become singular?
How far can one deform a given G_2 -metric?

Let us summarise first

- ▶ Only a few obstructions against G_2 -holonomy are known
- ▶ Only a few compact examples are known—only $\sim 10^8$ deformation families
- ▶ Known compact examples represent points close to the boundary of the moduli space—
the G_2 -metric is locally close to metrics with holonomy groups $SU(2)$ or $SU(3)$

Important open problems / questions

- ▶ Find more invariants for / obstructions against G_2 -metrics
- ▶ Construct G_2 -metrics far away from the boundary of the moduli space
- ▶ How can families of G_2 -metrics become singular?
How far can one deform a given G_2 -metric?
- ▶ Construct G_2 -metrics with prescribed singularities

Let us summarise first

- ▶ Only a few obstructions against G_2 -holonomy are known
- ▶ Only a few compact examples are known—only $\sim 10^8$ deformation families
- ▶ Known compact examples represent points close to the boundary of the moduli space—
the G_2 -metric is locally close to metrics with holonomy groups $SU(2)$ or $SU(3)$

Important open problems / questions

- ▶ Find more invariants for / obstructions against G_2 -metrics
- ▶ Construct G_2 -metrics far away from the boundary of the moduli space
- ▶ How can families of G_2 -metrics become singular?
How far can one deform a given G_2 -metric?
- ▶ Construct G_2 -metrics with prescribed singularities
Singularities allow massless particles to appear in M -theory

We want to describe G_2 -manifolds using differential topology

Definition

A G_2 -structure on a seven-manifold M is a reduction of the $GL(7, \mathbb{R})$ -frame bundle to a bundle with structure group G_2

We want to describe G_2 -manifolds using differential topology

Definition

A G_2 -structure on a seven-manifold M is a reduction of the $GL(7, \mathbb{R})$ -frame bundle to a bundle with structure group G_2

Equivalent descriptions

- ▶ Positive three form φ on M

We want to describe G_2 -manifolds using differential topology

Definition

A G_2 -structure on a seven-manifold M is a reduction of the $GL(7, \mathbb{R})$ -frame bundle to a bundle with structure group G_2

Equivalent descriptions

- ▶ Positive three form φ on M
- ▶ Riemannian metric, spin structure, and a unit spinor (up to sign)

We want to describe G_2 -manifolds using differential topology

Definition

A G_2 -structure on a seven-manifold M is a reduction of the $GL(7, \mathbb{R})$ -frame bundle to a bundle with structure group G_2

Equivalent descriptions

- ▶ Positive three form φ on M
- ▶ Riemannian metric, spin structure, and a unit spinor (up to sign)

Idea. Use nowhere vanishing spinors to describe and distinguish G_2 -structures

When we (Diarmuid Crowley, Johannes Nordström and myself) started our project, we asked the following questions

When we (Diarmuid Crowley, Johannes Nordström and myself) started our project, we asked the following questions

Can different constructions of G_2 -holonomy metrics

- ▶ lead to the same closed 7-manifold up to diffeomorphism?

When we (Diarmuid Crowley, Johannes Nordström and myself) started our project, we asked the following questions

Can different constructions of G_2 -holonomy metrics

- ▶ lead to the same closed 7-manifold up to diffeomorphism?
- ▶ if so, are the underlying G_2 -structures the same up to homotopy and spin diffeomorphism?

When we (Diarmuid Crowley, Johannes Nordström and myself) started our project, we asked the following questions

Can different constructions of G_2 -holonomy metrics

- ▶ lead to the same closed 7-manifold up to diffeomorphism?
- ▶ if so, are the underlying G_2 -structures the same up to homotopy and spin diffeomorphism?
- ▶ if so, do the two metrics belong to the same connected component of the G_2 -moduli space?

When we (Diarmuid Crowley, Johannes Nordström and myself) started our project, we asked the following questions

Can different constructions of G_2 -holonomy metrics

- ▶ lead to the same closed 7-manifold up to diffeomorphism?
- ▶ if so, are the underlying G_2 -structures the same up to homotopy and spin diffeomorphism?
- ▶ if so, do the two metrics belong to the same connected component of the G_2 -moduli space?

Diarmuid Crowley and Johannes Nordström also asked

- ▶ Are there pairs of G_2 -manifolds that are homeomorphic but not diffeomorphic?

Assume that M is a simply connected oriented spin 7-manifold with $H^4(M)$ torsion-free

Let $d \in \mathbb{N}$ be the divisibility of $p_1(TM)$, then $4 \mid d$ (Wu)

Let $\tilde{d} = \text{lcm}(4, d/2)$

Assume that M is a simply connected oriented spin 7-manifold with $H^4(M)$ torsion-free

Let $d \in \mathbb{N}$ be the divisibility of $p_1(TM)$, then $4 \mid d$ (Wu)

Let $\tilde{d} = \text{lcm}(4, d/2)$

Let s be a nowhere vanishing spinor on M , defining a G_2 -structure

Assume that M is a simply connected oriented spin 7-manifold with $H^4(M)$ torsion-free

Let $d \in \mathbb{N}$ be the divisibility of $p_1(TM)$, then $4 \mid d$ (Wu)

Let $\tilde{d} = \text{lcm}(4, d/2)$

Let s be a nowhere vanishing spinor on M , defining a G_2 -structure

Relevant differential topological invariants

$$\mu(M) \in \mathbb{Z}/\text{gcd}(28, \tilde{d}/4) \quad \text{generalised Eells-Kuiper invariant}$$

Assume that M is a simply connected oriented spin 7-manifold with $H^4(M)$ torsion-free

Let $d \in \mathbb{N}$ be the divisibility of $p_1(TM)$, then $4 \mid d$ (Wu)

Let $\tilde{d} = \text{lcm}(4, d/2)$

Let s be a nowhere vanishing spinor on M , defining a G_2 -structure

Relevant differential topological invariants

$\mu(M) \in \mathbb{Z}/\text{gcd}(28, \tilde{d}/4)$ generalised Eells-Kuiper invariant

$\xi(M, s) \in \mathbb{Z}/3\tilde{d}$

Assume that M is a simply connected oriented spin 7-manifold with $H^4(M)$ torsion-free

Let $d \in \mathbb{N}$ be the divisibility of $p_1(TM)$, then $4 \mid d$ (Wu)

Let $\tilde{d} = \text{lcm}(4, d/2)$

Let s be a nowhere vanishing spinor on M , defining a G_2 -structure

Relevant differential topological invariants

$\mu(M) \in \mathbb{Z}/\text{gcd}(28, \tilde{d}/4)$ generalised Eells-Kuiper invariant

$\xi(M, s) \in \mathbb{Z}/3\tilde{d}$

$\nu(M, s) \in \mathbb{Z}/48$ see below

Assume that M is a simply connected oriented spin 7-manifold with $H^4(M)$ torsion-free

Let $d \in \mathbb{N}$ be the divisibility of $p_1(TM)$, then $4 \mid d$ (Wu)

Let $\tilde{d} = \text{lcm}(4, d/2)$

Let s be a nowhere vanishing spinor on M , defining a G_2 -structure

Relevant differential topological invariants

$\mu(M) \in \mathbb{Z}/\text{gcd}(28, \tilde{d}/4)$ generalised Eells-Kuiper invariant

$\xi(M, s) \in \mathbb{Z}/3\tilde{d}$

$\nu(M, s) \in \mathbb{Z}/48$ see below

Important relations

$$\xi(M, s) \equiv 7\nu(M, s) \pmod{12}$$

$$\frac{\xi(M, s) - 7\nu(M, s)}{12} \equiv \mu(M) \pmod{\text{gcd}(28, \tilde{d}/4)}$$

Let σ_0, σ_1 be two nowhere vanishing spinors. Extend to $\bar{\sigma} \in \Gamma(\mathcal{S}^+(M \times [0, 1]))$
A generic $\bar{\sigma}$ will have nondegenerate isolated zeros because

$$\text{rk } \mathcal{S}^+(M \times [0, 1]) = 8 = \dim(M \times [0, 1])$$

Let σ_0, σ_1 be two nowhere vanishing spinors. Extend to $\bar{\sigma} \in \Gamma(\mathcal{S}^+(M \times [0, 1]))$
A generic $\bar{\sigma}$ will have nondegenerate isolated zeros because

$$\text{rk } \mathcal{S}^+(M \times [0, 1]) = 8 = \dim(M \times [0, 1])$$

Orient $\mathcal{S}^+(M \times [0, 1])$ and count with signs

$$\Delta\nu(M; \sigma_0, \sigma_1) = 2 \cdot \#\bar{\sigma}^{-1}(0) = 2 \cdot \sum_{p \in \bar{\sigma}^{-1}(0)} \text{sign}(d_p \bar{\sigma})$$

Let σ_0, σ_1 be two nowhere vanishing spinors. Extend to $\bar{\sigma} \in \Gamma(\mathcal{S}^+(M \times [0, 1]))$
A generic $\bar{\sigma}$ will have nondegenerate isolated zeros because

$$\text{rk } \mathcal{S}^+(M \times [0, 1]) = 8 = \dim(M \times [0, 1])$$

Orient $\mathcal{S}^+(M \times [0, 1])$ and count with signs

$$\Delta\nu(M; \sigma_0, \sigma_1) = 2 \cdot \#\bar{\sigma}^{-1}(0) = 2 \cdot \sum_{p \in \bar{\sigma}^{-1}(0)} \text{sign}(d_p \bar{\sigma})$$

Theorem (Crowley-Nordström)

Let $F: M \rightarrow M$ be a spin diffeomorphism, then

$$\Delta\nu(M; \sigma, F^* \sigma) \in 48\mathbb{Z}$$

Let σ_0, σ_1 be two nowhere vanishing spinors. Extend to $\bar{\sigma} \in \Gamma(\mathcal{S}^+(M \times [0, 1]))$
 A generic $\bar{\sigma}$ will have nondegenerate isolated zeros because

$$\text{rk } \mathcal{S}^+(M \times [0, 1]) = 8 = \dim(M \times [0, 1])$$

Orient $\mathcal{S}^+(M \times [0, 1])$ and count with signs

$$\Delta\nu(M; \sigma_0, \sigma_1) = 2 \cdot \#\bar{\sigma}^{-1}(0) = 2 \cdot \sum_{p \in \bar{\sigma}^{-1}(0)} \text{sign}(d_p \bar{\sigma})$$

Theorem (Crowley-Nordström)

Let $F: M \rightarrow M$ be a spin diffeomorphism, then

$$\Delta\nu(M; \sigma, F^* \sigma) \in 48\mathbb{Z}$$

Can we write $\Delta\nu(M; \sigma_0, \sigma_1) = \nu(M, \sigma_0) - \nu(M, \sigma_1) \in \mathbb{Z}/48$?

Idea. If M is spin, then M is the spin boundary of some compact 8-manifold W .
Extend σ to $\bar{\sigma} \in \Gamma(S^+W)$, then $\#\bar{\sigma}^{-1}(0)$ depends on W —**not well-defined yet!**

Idea. If M is spin, then M is the spin boundary of some compact 8-manifold W .
Extend σ to $\bar{\sigma} \in \Gamma(\mathcal{S}^+ W)$, then $\#\bar{\sigma}^{-1}(0)$ depends on W —**not well-defined yet!**

- ▶ $\chi(W)$ —Euler characteristic of W
- ▶ $\text{sign}(W)$ —signature of W

Definition (Crowley-Nordström)

Assume that $M = \partial W$ with W spin, compact. Define

$$\nu(M, \sigma) = \chi(W) - 3 \text{sign}(W) - 2\#\bar{\sigma}^{-1}(0) \pmod{48}$$

Idea. If M is spin, then M is the spin boundary of some compact 8-manifold W . Extend σ to $\bar{\sigma} \in \Gamma(\mathcal{S}^+ W)$, then $\#\bar{\sigma}^{-1}(0)$ depends on W —**not well-defined yet!**

- ▶ $\chi(W)$ —Euler characteristic of W
- ▶ $\text{sign}(W)$ —signature of W

Definition (Crowley-Nordström)

Assume that $M = \partial W$ with W spin, compact. Define

$$\nu(M, \sigma) = \chi(W) - 3 \text{sign}(W) - 2\#\bar{\sigma}^{-1}(0) \pmod{48}$$

Theorem (Crowley-Nordström)

$$\Delta\nu(M; \sigma_0, \sigma_1) = \nu(M, \sigma_0) - \nu(M, \sigma_1) \in \mathbb{Z}/48$$

Idea. If M is spin, then M is the spin boundary of some compact 8-manifold W . Extend σ to $\bar{\sigma} \in \Gamma(\mathcal{S}^+ W)$, then $\#\bar{\sigma}^{-1}(0)$ depends on W —**not well-defined yet!**

- ▶ $\chi(W)$ —Euler characteristic of W
- ▶ $\text{sign}(W)$ —signature of W

Definition (Crowley-Nordström)

Assume that $M = \partial W$ with W spin, compact. Define

$$\nu(M, \sigma) = \chi(W) - 3 \text{sign}(W) - 2\#\bar{\sigma}^{-1}(0) \pmod{48}$$

Theorem (Crowley-Nordström)

$$\Delta\nu(M; \sigma_0, \sigma_1) = \nu(M, \sigma_0) - \nu(M, \sigma_1) \in \mathbb{Z}/48$$

Problem. Given M , how to determine W with $M = \partial W$?

Idea. Use the APS-index theorem and Mathai-Quillen currents

Idea. Use the APS-index theorem and Mathai-Quillen currents

- ▶ $\psi(\nabla^{SM}, g^{SM})$ —Mathai-Quillen form in $\Omega^\bullet(SM)$
- ▶ D_M —spin Dirac operator on $\Gamma(SM)$
- ▶ B_M —odd signature operator $*d \pm d*$ on $\Omega^{\text{ev}}(M)$
- ▶ h —dimension of the kernel
- ▶ η —Atiyah-Patodi-Singer η -invariant

Idea. Use the APS-index theorem and Mathai-Quillen currents

- ▶ $\psi(\nabla^{SM}, g^{SM})$ —Mathai-Quillen form in $\Omega^\bullet(SM)$
- ▶ D_M —spin Dirac operator on $\Gamma(SM)$
- ▶ B_M —odd signature operator $*d \pm d*$ on $\Omega^{\text{ev}}(M)$
- ▶ h —dimension of the kernel
- ▶ η —Atiyah-Patodi-Singer η -invariant

Theorem (Crowley-G-Nordström)

$$\nu(M, \sigma) = 2 \int_M \sigma^* \psi(\nabla^{SM}, g^{SM}) - 24(\eta + h)(D_M) + 3\eta(B_M) \in \mathbb{Z}/48$$

Proof.

Use $2e(\nabla^{S^+W}) = e(\nabla) + 48\hat{A}(\nabla)^{[8]} - 3L(\nabla)^{[8]} \in \Omega^8(W)$



In the case of G_2 -holonomy, things simplify

- ▶ σ is parallel, so $\sigma^*\psi(\nabla^{SM}, g^{SM}) = 0$

In the case of G_2 -holonomy, things simplify

- ▶ σ is parallel, so $\sigma^*\psi(\nabla^{SM}, g^{SM}) = 0$
- ▶ Harmonic spinors are parallel, so $h(D_M) = 1 + b_1(M)$

In the case of G_2 -holonomy, things simplify

- ▶ σ is parallel, so $\sigma^*\psi(\nabla^{SM}, g^{SM}) = 0$
- ▶ Harmonic spinors are parallel, so $h(D_M) = 1 + b_1(M)$
- ▶ $\eta(D_M) \in \mathbb{R}$ is smooth on the G_2 -moduli space \mathcal{M}

In the case of G_2 -holonomy, things simplify

- ▶ σ is parallel, so $\sigma^*\psi(\nabla^{SM}, g^{SM}) = 0$
- ▶ Harmonic spinors are parallel, so $h(D_M) = 1 + b_1(M)$
- ▶ $\eta(D_M) \in \mathbb{R}$ is smooth on the G_2 -moduli space \mathcal{M}

Definition (Crowley-G-Nordström)

Let (M, g) be a compact manifold with $\text{Hol}(M, g) \subset G_2$. Put

$$\bar{\nu}(M, g) = 3\eta(B_M) - 24\eta(D_M) \in \mathbb{Z}$$

In the case of G_2 -holonomy, things simplify

- ▶ σ is parallel, so $\sigma^*\psi(\nabla^{SM}, g^{SM}) = 0$
- ▶ Harmonic spinors are parallel, so $h(D_M) = 1 + b_1(M)$
- ▶ $\eta(D_M) \in \mathbb{R}$ is smooth on the G_2 -moduli space \mathcal{M}

Definition (Crowley-G-Nordström)

Let (M, g) be a compact manifold with $\text{Hol}(M, g) \subset G_2$. Put

$$\bar{\nu}(M, g) = 3\eta(B_M) - 24\eta(D_M) \in \mathbb{Z}$$

- ▶ $\nu(M, \sigma) \equiv \bar{\nu}(M, g) - 24(1 + b_1(M)) \pmod{48}$

In the case of G_2 -holonomy, things simplify

- ▶ σ is parallel, so $\sigma^*\psi(\nabla^{SM}, g^{SM}) = 0$
- ▶ Harmonic spinors are parallel, so $h(D_M) = 1 + b_1(M)$
- ▶ $\eta(D_M) \in \mathbb{R}$ is smooth on the G_2 -moduli space \mathcal{M}

Definition (Crowley-G-Nordström)

Let (M, g) be a compact manifold with $\text{Hol}(M, g) \subset G_2$. Put

$$\bar{\nu}(M, g) = 3\eta(B_M) - 24\eta(D_M) \in \mathbb{Z}$$

- ▶ $\nu(M, \sigma) \equiv \bar{\nu}(M, g) - 24(1 + b_1(M)) \pmod{48}$
- ▶ $\bar{\nu}(M, g)$ is locally constant on \mathcal{M}

In the case of G_2 -holonomy, things simplify

- ▶ σ is parallel, so $\sigma^*\psi(\nabla^{SM}, g^{SM}) = 0$
- ▶ Harmonic spinors are parallel, so $h(D_M) = 1 + b_1(M)$
- ▶ $\eta(D_M) \in \mathbb{R}$ is smooth on the G_2 -moduli space \mathcal{M}

Definition (Crowley-G-Nordström)

Let (M, g) be a compact manifold with $\text{Hol}(M, g) \subset G_2$. Put

$$\bar{\nu}(M, g) = 3\eta(B_M) - 24\eta(D_M) \in \mathbb{Z}$$

- ▶ $\nu(M, \sigma) \equiv \bar{\nu}(M, g) - 24(1 + b_1(M)) \pmod{48}$
- ▶ $\bar{\nu}(M, g)$ is locally constant on \mathcal{M}
- ▶ $\bar{\nu}(M, g) = 0$ if M admits an orientation reversing isometry

What about the known examples by Joyce and Kovalev?

- ▶ $\bar{\nu}(M, g) = 0$ for all twisted connected sums
(but $\xi(M, s) \neq 0$ is possible—Wallis, arXiv:1808.09443)

What about the known examples by Joyce and Kovalev?

- ▶ $\bar{\nu}(M, g) = 0$ for all twisted connected sums
(but $\xi(M, s) \neq 0$ is possible—Wallis, arXiv:1808.09443)
- ▶ $\bar{\nu}(M, g) = 0$ for some of Joyce's examples—some are twisted connected sums, some admit orientation reversing isometries

What about the known examples by Joyce and Kovalev?

- ▶ $\bar{\nu}(M, g) = 0$ for all twisted connected sums
(but $\xi(M, s) \neq 0$ is possible—Wallis, arXiv:1808.09443)
- ▶ $\bar{\nu}(M, g) = 0$ for some of Joyce's examples—some are twisted connected sums, some admit orientation reversing isometries

Question. Is $\bar{\nu}(M, g) = 0$ whenever $\text{Hol}(M, g) = G_2$?

What about the known examples by Joyce and Kovalev?

- ▶ $\bar{\nu}(M, g) = 0$ for all twisted connected sums
(but $\xi(M, s) \neq 0$ is possible—Wallis, arXiv:1808.09443)
- ▶ $\bar{\nu}(M, g) = 0$ for some of Joyce's examples—some are twisted connected sums, some admit orientation reversing isometries

Question. Is $\bar{\nu}(M, g) = 0$ whenever $\text{Hol}(M, g) = G_2$?

- ▶ If **yes**, then $\bar{\nu}(M, g) \neq 0$ or $\nu(M, \sigma) \neq 24$ is a new obstruction against G_2 -holonomy

What about the known examples by Joyce and Kovalev?

- ▶ $\bar{\nu}(M, g) = 0$ for all twisted connected sums
(but $\xi(M, s) \neq 0$ is possible—Wallis, arXiv:1808.09443)
- ▶ $\bar{\nu}(M, g) = 0$ for some of Joyce's examples—some are twisted connected sums, some admit orientation reversing isometries

Question. Is $\bar{\nu}(M, g) = 0$ whenever $\text{Hol}(M, g) = G_2$?

- ▶ If **yes**, then $\bar{\nu}(M, g) \neq 0$ or $\nu(M, \sigma) \neq 24$ is a new obstruction against G_2 -holonomy
- ▶ If **no**, then $\bar{\nu}(M, g)$ is a non-trivial new invariant

What about the known examples by Joyce and Kovalev?

- ▶ $\bar{\nu}(M, g) = 0$ for all twisted connected sums
(but $\xi(M, s) \neq 0$ is possible—Wallis, arXiv:1808.09443)
- ▶ $\bar{\nu}(M, g) = 0$ for some of Joyce's examples—some are twisted connected sums, some admit orientation reversing isometries

Question. Is $\bar{\nu}(M, g) = 0$ whenever $\text{Hol}(M, g) = G_2$?

- ▶ If **yes**, then $\bar{\nu}(M, g) \neq 0$ or $\nu(M, \sigma) \neq 24$ is a new obstruction against G_2 -holonomy
- ▶ If **no**, then $\bar{\nu}(M, g)$ is a non-trivial new invariant

Answer. We will construct examples with $\bar{\nu}(M, g) \neq 0$

What about the known examples by Joyce and Kovalev?

- ▶ $\bar{\nu}(M, g) = 0$ for all twisted connected sums
(but $\xi(M, s) \neq 0$ is possible—Wallis, arXiv:1808.09443)
- ▶ $\bar{\nu}(M, g) = 0$ for some of Joyce's examples—some are twisted connected sums, some admit orientation reversing isometries


Question. Is $\bar{\nu}(M, g) = 0$ whenever $\text{Hol}(M, g) = G_2$?

- ▶ If **yes**, then $\bar{\nu}(M, g) \neq 0$ or $\nu(M, \sigma) \neq 24$ is a new obstruction against G_2 -holonomy
- ▶ If **no**, then $\bar{\nu}(M, g)$ is a non-trivial new invariant


Answer. We will construct examples with $\bar{\nu}(M, g) \neq 0$

Using $\bar{\nu}(M, g)$, we will show that for some particular M , the G_2 -moduli space \mathcal{M} has several connected components


Recall twisted connected sums



Gluing of tori at angle $\vartheta = \frac{\pi}{2}$ between exterior circles



Extra twisted connected sums




Assume that $\Gamma_{\pm} \cong \mathbb{Z}/k_{\pm}$ acts both on V_{\pm} and on $S^1_{\pm,ext}$

The induced action on ∂V_{\pm} has to fix Σ_{\pm} pointwise

The actions on $S^1_{\pm,int}$ and $S^1_{\pm,ext}$ have to be free

Extra twisted connected sums




Assume that $\Gamma_{\pm} \cong \mathbb{Z}/k_{\pm}$ acts both on V_{\pm} and on $S^1_{\pm,ext}$

The induced action on ∂V_{\pm} has to fix Σ_{\pm} pointwise

The actions on $S^1_{\pm,int}$ and $S^1_{\pm,ext}$ have to be free

Then $(S^1_{\pm,int} \times S^1_{\pm,ext})/\Gamma_{\pm}$ is again a flat 2-torus

Extra twisted connected sums



Assume that $\Gamma_{\pm} \cong \mathbb{Z}/k_{\pm}$ acts both on V_{\pm} and on $S^1_{\pm,ext}$

The induced action on ∂V_{\pm} has to fix Σ_{\pm} pointwise


The actions on $S^1_{\pm,int}$ and $S^1_{\pm,ext}$ have to be free

Then $(S^1_{\pm,int} \times S^1_{\pm,ext})/\Gamma_{\pm}$ is again a flat 2-torus


If both the tori and the K3 surfaces are isometric,

we can glue $M_{\pm} = (V_{\pm} \times S^1_{\pm,ext})/\Gamma_{\pm}$ at various angles ϑ


Extra twisted connected sums




Modified gluing of tori at angle $\vartheta = \frac{3}{4}\pi$




Extra twisted connected sums




Modified gluing of tori at angle $\vartheta = \frac{2}{3}\pi$



Extra twisted connected sums



Modified gluing of tori at angle $\vartheta = \arccos(\frac{1}{\sqrt{6}})$



The CY structures on V_{\pm} induce G_2 -structures on M_{\pm} and on $\Sigma_{\pm} \times T_{\pm}^2 \times \mathbb{R}$

Then we need an **isomorphism of G_2 -manifolds** $\Sigma_+ \times T_+^2 \times \mathbb{R} \xrightarrow{\cong} \Sigma_- \times T_-^2 \times \mathbb{R}$

The CY structures on V_{\pm} induce G_2 -structures on M_{\pm} and on $\Sigma_{\pm} \times T_{\pm}^2 \times \mathbb{R}$

Then we need an **isomorphism of G_2 -manifolds** $\Sigma_+ \times T_+^2 \times \mathbb{R} \xrightarrow{\cong} \Sigma_- \times T_-^2 \times \mathbb{R}$

Let u_{\pm}, v_{\pm} be coordinates on the interior and exterior circles, respectively

$$u_- = -u_+ \cos \vartheta + v_+ \sin \vartheta \quad \text{and} \quad v_- = u_+ \sin \vartheta + v_+ \cos \vartheta$$

Let $t = t_- = -t_+$ be the coordinate in the cylinder direction

The CY structures on V_{\pm} induce G_2 -structures on M_{\pm} and on $\Sigma_{\pm} \times T_{\pm}^2 \times \mathbb{R}$

Then we need an **isomorphism of G_2 -manifolds** $\Sigma_+ \times T_+^2 \times \mathbb{R} \xrightarrow{\cong} \Sigma_- \times T_-^2 \times \mathbb{R}$

Let u_{\pm}, v_{\pm} be coordinates on the interior and exterior circles, respectively

$$u_- = -u_+ \cos \vartheta + v_+ \sin \vartheta \quad \text{and} \quad v_- = u_+ \sin \vartheta + v_+ \cos \vartheta$$

Let $t = t_- = -t_+$ be the coordinate in the cylinder direction

Let $\omega_1^{\pm}, \omega_2^{\pm}, \omega_3^{\pm} \in \Omega^{2,+}(\Sigma)$ be hyperkähler triples, ω_1^{\pm} is the Kähler form from V_{\pm}

The CY structures on V_{\pm} induce G_2 -structures on M_{\pm} and on $\Sigma_{\pm} \times T_{\pm}^2 \times \mathbb{R}$

Then we need an **isomorphism of G_2 -manifolds** $\Sigma_+ \times T_+^2 \times \mathbb{R} \xrightarrow{\cong} \Sigma_- \times T_-^2 \times \mathbb{R}$

Let u_{\pm}, v_{\pm} be coordinates on the interior and exterior circles, respectively

$$u_- = -u_+ \cos \vartheta + v_+ \sin \vartheta \quad \text{and} \quad v_- = u_+ \sin \vartheta + v_+ \cos \vartheta$$

Let $t = t_- = -t_+$ be the coordinate in the cylinder direction

Let $\omega_1^{\pm}, \omega_2^{\pm}, \omega_3^{\pm} \in \Omega^{2,+}(\Sigma)$ be hyperkähler triples, ω_1^{\pm} is the Kähler form from V_{\pm}


$$\begin{aligned} \varphi &= dv_{\pm} \wedge \omega_1^{\pm} + du_{\pm} \wedge \omega_2^{\pm} + dt_{\pm} \wedge \omega_3^{\pm} + dt_{\pm} \wedge du_{\pm} \wedge dv_{\pm} \\ \omega_1^- &= \cos \vartheta \omega_1^+ + \sin \vartheta \omega_2^+ & \omega_2^- &= \sin \vartheta \omega_1^+ - \cos \vartheta \omega_2^+ & \omega_3^- &= -\omega_3^+ \end{aligned}$$

Both the torus matching and the K3 matching depend on the **gluing angle** ϑ

Assume that $\Gamma_{\pm} \cong \mathbb{Z}/k_{\pm}$ acts on $V_{\pm} \times S^1_{\pm, \text{ext}}$

A **torus matching** is described by

- ▶ A number $\varepsilon_+ \in (\mathbb{Z}/k_+)^{\times}$ if $k_+ > 1$




Assume that $\Gamma_{\pm} \cong \mathbb{Z}/k_{\pm}$ acts on $V_{\pm} \times S^1_{\pm, \text{ext}}$

A **torus matching** is described by

- ▶ A number $\varepsilon_+ \in (\mathbb{Z}/k_+)^{\times}$ if $k_+ > 1$
- ▶ A **gluing matrix** $G = \begin{pmatrix} m & p \\ n & q \end{pmatrix}$
with $\det G = -k_+ k_-$ and $mq \leq 0, np \geq 0$


satisfying some extra conditions (only finitely many choices for ε_+, G possible)



Assume that $\Gamma_{\pm} \cong \mathbb{Z}/k_{\pm}$ acts on $V_{\pm} \times S^1_{\pm, \text{ext}}$

A **torus matching** is described by

- ▶ A number $\varepsilon_+ \in (\mathbb{Z}/k_+)^{\times}$ if $k_+ > 1$
- ▶ A **gluing matrix** $G = \begin{pmatrix} m & p \\ n & q \end{pmatrix}$
with $\det G = -k_+k_-$ and $mq \leq 0, np \geq 0$



satisfying some extra conditions (only finitely many choices for ε_+ , G possible)

From G , recover

- ▶ The **aspect ratios** $s_+ = \frac{\ell(S^1_{+, \text{ext}})}{\ell(S^1_{+, \text{int}})} = \sqrt{-\frac{nq}{mp}}$ and $s_- = \frac{\ell(S^1_{-, \text{ext}})}{\ell(S^1_{-, \text{int}})} = \sqrt{-\frac{mn}{pq}}$
- ▶ The **gluing angle** $\vartheta = \arg(ms_+ + in) \in (-\pi, \pi]$
- ▶ The **fundamental group** $\pi_1(M) \cong \mathbb{Z}/p$

If M is an extra twisted connect sum, identify $\Sigma = \Sigma_+ \cong \Sigma_-$, let

$$N_{\pm} = \text{im}(H^2(V_{\pm}) \rightarrow H^2(\Sigma)) \subset L = H^2(\Sigma) \cong E_8^2 \oplus U^3$$

The pair (N_+, N_-) of sublattices of L (up to $\text{Aut}(L)$) is called a **configuration**

If M is an extra twisted connect sum, identify $\Sigma = \Sigma_+ \cong \Sigma_-$, let

$$N_{\pm} = \text{im}(H^2(V_{\pm}) \rightarrow H^2(\Sigma)) \subset L = H^2(\Sigma) \cong E_8^2 \oplus U^3$$

The pair (N_+, N_-) of sublattices of L (up to $\text{Aut}(L)$) is called a **configuration**

We get positive orthonormal triples $[\omega_1^{\pm}] \in N_{\pm, \mathbb{R}}$ and $[\omega_2^{\pm}], [\omega_3^{\pm}] \in N_{\pm, \mathbb{R}}^{\perp} \subset L_{\mathbb{R}}$

If M is an extra twisted connect sum, identify $\Sigma = \Sigma_+ \cong \Sigma_-$, let

$$N_{\pm} = \text{im}(H^2(V_{\pm}) \rightarrow H^2(\Sigma)) \subset L = H^2(\Sigma) \cong E_8^2 \oplus U^3$$

The pair (N_+, N_-) of sublattices of L (up to $\text{Aut}(L)$) is called a **configuration**

We get positive orthonormal triples $[\omega_1^{\pm}] \in N_{\pm, \mathbb{R}}$ and $[\omega_2^{\pm}], [\omega_3^{\pm}] \in N_{\pm, \mathbb{R}}^{\perp} \subset L_{\mathbb{R}}$

Let $A_{N_{\pm}}$ denote the reflections of $L_{\mathbb{R}} \cong \mathbb{R}^{3,19}$ in $N_{\pm, \mathbb{R}}$. Because

$$[\omega_1^-] = \cos \vartheta [\omega_1^+] + \sin \vartheta [\omega_2^+] \quad \text{and} \quad [\omega_2^-] = \sin \vartheta [\omega_1^+] - \cos \vartheta [\omega_2^+] \quad (*)$$

the classes $[\omega_1^{\pm}], [\omega_2^{\pm}]$ lie in the subspace $L_{2\vartheta} \subset L_{\mathbb{R}}$ that $A_{N_+} A_{N_-}$ rotates through 2ϑ

If M is an extra twisted connect sum, identify $\Sigma = \Sigma_+ \cong \Sigma_-$, let

$$N_{\pm} = \text{im}(H^2(V_{\pm}) \rightarrow H^2(\Sigma)) \subset L = H^2(\Sigma) \cong E_8^2 \oplus U^3$$

The pair (N_+, N_-) of sublattices of L (up to $\text{Aut}(L)$) is called a **configuration**

We get positive orthonormal triples $[\omega_1^{\pm}] \in N_{\pm, \mathbb{R}}$ and $[\omega_2^{\pm}], [\omega_3^{\pm}] \in N_{\pm, \mathbb{R}}^{\perp} \subset L_{\mathbb{R}}$

Let $A_{N_{\pm}}$ denote the reflections of $L_{\mathbb{R}} \cong \mathbb{R}^{3,19}$ in $N_{\pm, \mathbb{R}}$. Because

$$[\omega_1^-] = \cos \vartheta [\omega_1^+] + \sin \vartheta [\omega_2^+] \quad \text{and} \quad [\omega_2^-] = \sin \vartheta [\omega_1^+] - \cos \vartheta [\omega_2^+] \quad (*)$$

the classes $[\omega_1^{\pm}], [\omega_2^{\pm}]$ lie in the subspace $L_{2\vartheta} \subset L_{\mathbb{R}}$ that $A_{N_+} A_{N_-}$ rotates through 2ϑ

Matching Problem. To construct M , find $N_+, N_- \subset L$ and positive classes $[\omega_1^{\pm}] \in L_{2\vartheta} \cap N_{\pm, \mathbb{R}}$ and $[\omega_2^{\pm}] \in L_{2\vartheta} \cap N_{\pm, \mathbb{R}}^{\perp}$ of length 1 satisfying (*)

If M is an extra twisted connect sum, identify $\Sigma = \Sigma_+ \cong \Sigma_-$, let

$$N_{\pm} = \text{im}(H^2(V_{\pm}) \rightarrow H^2(\Sigma)) \subset L = H^2(\Sigma) \cong E_8^2 \oplus U^3$$

The pair (N_+, N_-) of sublattices of L (up to $\text{Aut}(L)$) is called a **configuration**

We get positive orthonormal triples $[\omega_1^{\pm}] \in N_{\pm, \mathbb{R}}$ and $[\omega_2^{\pm}], [\omega_3^{\pm}] \in N_{\pm, \mathbb{R}}^{\perp} \subset L_{\mathbb{R}}$

Let $A_{N_{\pm}}$ denote the reflections of $L_{\mathbb{R}} \cong \mathbb{R}^{3,19}$ in $N_{\pm, \mathbb{R}}$. Because

$$[\omega_1^-] = \cos \vartheta [\omega_1^+] + \sin \vartheta [\omega_2^+] \quad \text{and} \quad [\omega_2^-] = \sin \vartheta [\omega_1^+] - \cos \vartheta [\omega_2^+] \quad (*)$$

the classes $[\omega_1^{\pm}], [\omega_2^{\pm}]$ lie in the subspace $L_{2\vartheta} \subset L_{\mathbb{R}}$ that $A_{N_+} A_{N_-}$ rotates through 2ϑ

Matching Problem. To construct M , find $N_+, N_- \subset L$ and positive classes $[\omega_1^{\pm}] \in L_{2\vartheta} \cap N_{\pm, \mathbb{R}}$ and $[\omega_2^{\pm}] \in L_{2\vartheta} \cap N_{\pm, \mathbb{R}}^{\perp}$ of length 1 satisfying (*)

Then find V_{\pm} with \mathbb{Z}/k_{\pm} -actions and Σ that “realise” $N_{\pm} \subset L$ and $[\omega_1^{\pm}], \dots, [\omega_3^{\pm}]$

If M is an extra twisted connect sum, identify $\Sigma = \Sigma_+ \cong \Sigma_-$, let

$$N_{\pm} = \text{im}(H^2(V_{\pm}) \rightarrow H^2(\Sigma)) \subset L = H^2(\Sigma) \cong E_8^2 \oplus U^3$$

The pair (N_+, N_-) of sublattices of L (up to $\text{Aut}(L)$) is called a **configuration**

We get positive orthonormal triples $[\omega_1^{\pm}] \in N_{\pm, \mathbb{R}}$ and $[\omega_2^{\pm}], [\omega_3^{\pm}] \in N_{\pm, \mathbb{R}}^{\perp} \subset L_{\mathbb{R}}$

Let $A_{N_{\pm}}$ denote the reflections of $L_{\mathbb{R}} \cong \mathbb{R}^{3,19}$ in $N_{\pm, \mathbb{R}}$. Because

$$[\omega_1^-] = \cos \vartheta [\omega_1^+] + \sin \vartheta [\omega_2^+] \quad \text{and} \quad [\omega_2^-] = \sin \vartheta [\omega_1^+] - \cos \vartheta [\omega_2^+] \quad (*)$$

the classes $[\omega_1^{\pm}], [\omega_2^{\pm}]$ lie in the subspace $L_{2\vartheta} \subset L_{\mathbb{R}}$ that $A_{N_+} A_{N_-}$ rotates through 2ϑ

Matching Problem. To construct M , find $N_+, N_- \subset L$ and positive classes $[\omega_1^{\pm}] \in L_{2\vartheta} \cap N_{\pm, \mathbb{R}}$ and $[\omega_2^{\pm}] \in L_{2\vartheta} \cap N_{\pm, \mathbb{R}}^{\perp}$ of length 1 satisfying (*)

Then find V_{\pm} with \mathbb{Z}/k_{\pm} -actions and Σ that “realise” $N_{\pm} \subset L$ and $[\omega_1^{\pm}], \dots, [\omega_3^{\pm}]$

All this works for “easy” V_{\pm} if $\text{rk } N_+ = \text{rk } N_-$ and $N_{\pm, \mathbb{R}} \subset L_{2\vartheta}$

To construct an extra twisted connected sum

- ▶ Pick deformation types V_{\pm} of asymptotically cylindrical Calabi-Yau threefolds with \mathbb{Z}/k_{\pm} -actions (can be constructed from weak Fano threefolds Z_{\pm})

To construct an extra twisted connected sum

- ▶ Pick deformation types V_{\pm} of asymptotically cylindrical Calabi-Yau threefolds with \mathbb{Z}/k_{\pm} -actions (can be constructed from weak Fano threefolds Z_{\pm})
- ▶ Find a configuration (N_+, N_-) for the sublattices $N_{\pm} \subset L$ induced by V_{\pm}

To construct an extra twisted connected sum

- ▶ Pick deformation types V_{\pm} of asymptotically cylindrical Calabi-Yau threefolds with \mathbb{Z}/k_{\pm} -actions (can be constructed from weak Fano threefolds Z_{\pm})
- ▶ Find a configuration (N_+, N_-) for the sublattices $N_{\pm} \subset L$ induced by V_{\pm}
- ▶ Determine hyperkähler triples with gluing angle ϑ induced by actual varieties V_{\pm} (requires knowledge of moduli spaces of Σ_{\pm} and Z_{\pm})

To construct an extra twisted connected sum

- ▶ Pick deformation types V_{\pm} of asymptotically cylindrical Calabi-Yau threefolds with \mathbb{Z}/k_{\pm} -actions (can be constructed from weak Fano threefolds Z_{\pm})
- ▶ Find a configuration (N_+, N_-) for the sublattices $N_{\pm} \subset L$ induced by V_{\pm}
- ▶ Determine hyperkähler triples with gluing angle ϑ induced by actual varieties V_{\pm} (requires knowledge of moduli spaces of Σ_{\pm} and Z_{\pm})
- ▶ Find $\varepsilon_+ \in (\mathbb{Z}/k_+)^{\times}$ if $k_+ > 1$ and a gluing matrix G with gluing angle ϑ

To construct an extra twisted connected sum

- ▶ Pick deformation types V_{\pm} of asymptotically cylindrical Calabi-Yau threefolds with \mathbb{Z}/k_{\pm} -actions (can be constructed from weak Fano threefolds Z_{\pm})
- ▶ Find a configuration (N_+, N_-) for the sublattices $N_{\pm} \subset L$ induced by V_{\pm}
- ▶ Determine hyperkähler triples with gluing angle ϑ induced by actual varieties V_{\pm} (requires knowledge of moduli spaces of Σ_{\pm} and Z_{\pm})
- ▶ Find $\varepsilon_+ \in (\mathbb{Z}/k_+)^{\times}$ if $k_+ > 1$ and a gluing matrix G with gluing angle ϑ
- ▶ Determine the asymptotically cylindrical Calabi-Yau metrics on V_{\pm}

To construct an extra twisted connected sum

- ▶ Pick deformation types V_{\pm} of asymptotically cylindrical Calabi-Yau threefolds with \mathbb{Z}/k_{\pm} -actions (can be constructed from weak Fano threefolds Z_{\pm})
- ▶ Find a configuration (N_+, N_-) for the sublattices $N_{\pm} \subset L$ induced by V_{\pm}
- ▶ Determine hyperkähler triples with gluing angle ϑ induced by actual varieties V_{\pm} (requires knowledge of moduli spaces of Σ_{\pm} and Z_{\pm})
- ▶ Find $\varepsilon_+ \in (\mathbb{Z}/k_+)^{\times}$ if $k_+ > 1$ and a gluing matrix G with gluing angle ϑ
- ▶ Determine the asymptotically cylindrical Calabi-Yau metrics on V_{\pm}
- ▶ Glue $(V_+ \times S_{\xi_+}^1)/\Gamma_+$ to $(V_- \times S_{\xi_-}^1)/\Gamma_-$ as specified by the data above

To construct an extra twisted connected sum

- ▶ Pick deformation types V_{\pm} of asymptotically cylindrical Calabi-Yau threefolds with \mathbb{Z}/k_{\pm} -actions (can be constructed from weak Fano threefolds Z_{\pm})
- ▶ Find a configuration (N_+, N_-) for the sublattices $N_{\pm} \subset L$ induced by V_{\pm}
- ▶ Determine hyperkähler triples with gluing angle ϑ induced by actual varieties V_{\pm} (requires knowledge of moduli spaces of Σ_{\pm} and Z_{\pm})
- ▶ Find $\varepsilon_+ \in (\mathbb{Z}/k_+)^{\times}$ if $k_+ > 1$ and a gluing matrix G with gluing angle ϑ
- ▶ Determine the asymptotically cylindrical Calabi-Yau metrics on V_{\pm}
- ▶ Glue $(V_+ \times S_{\xi_+}^1)/\Gamma_+$ to $(V_- \times S_{\xi_-}^1)/\Gamma_-$ as specified by the data above
- ▶ Find a torsion-free G_2 -structure close to the G_2 -structure obtained by gluing

To construct an extra twisted connected sum

- ▶ Pick deformation types V_{\pm} of asymptotically cylindrical Calabi-Yau threefolds with \mathbb{Z}/k_{\pm} -actions (can be constructed from weak Fano threefolds Z_{\pm})
- ▶ Find a configuration (N_+, N_-) for the sublattices $N_{\pm} \subset L$ induced by V_{\pm}
- ▶ Determine hyperkähler triples with gluing angle ϑ induced by actual varieties V_{\pm} (requires knowledge of moduli spaces of Σ_{\pm} and Z_{\pm})
- ▶ Find $\varepsilon_+ \in (\mathbb{Z}/k_+)^{\times}$ if $k_+ > 1$ and a gluing matrix G with gluing angle ϑ
- ▶ Determine the asymptotically cylindrical Calabi-Yau metrics on V_{\pm}
- ▶ Glue $(V_+ \times S_{\xi_+}^1)/\Gamma_+$ to $(V_- \times S_{\xi_-}^1)/\Gamma_-$ as specified by the data above
- ▶ Find a torsion-free G_2 -structure close to the G_2 -structure obtained by gluing

In the following, we will only need the configuration (N_+, N_-) , the gluing matrix G , and the number ε_+

With respect to modified Atiyah-Patodi-Singer boundary conditions, define

$$\bar{\nu}(M_{\pm}, g) = 3\eta_{\text{APS}}(B_{M_{\pm}}; L_{B, \pm}) - 24\eta_{\text{APS}}(D_{M_{\pm}}; L_{D, \pm}) \in \mathbb{R}$$

With respect to modified Atiyah-Patodi-Singer boundary conditions, define

$$\bar{\nu}(M_{\pm}, g) = 3\eta_{\text{APS}}(B_{M_{\pm}}; L_{B, \pm}) - 24\eta_{\text{APS}}(D_{M_{\pm}}; L_{D, \pm}) \in \mathbb{R}$$

The isometry $A_{N_+} A_{N_-}$ respects the decomposition $H^{2,+}(\Sigma) \oplus H^{2,-}(\Sigma)$

Let $\alpha_1^+, \dots, \alpha_3^+, \alpha_1^-, \dots, \alpha_{19}^- \in (-\pi, \pi]$ be the angles through which $A_{N_+} A_{N_-} \otimes \mathbb{C}$ rotates $H^{2,+}(\Sigma, \mathbb{C})$ and $H^{2,-}(\Sigma, \mathbb{C})$, respectively

With respect to modified Atiyah-Patodi-Singer boundary conditions, define

$$\bar{\nu}(M_{\pm}, g) = 3\eta_{\text{APS}}(B_{M_{\pm}}; L_{B, \pm}) - 24\eta_{\text{APS}}(D_{M_{\pm}}; L_{D, \pm}) \in \mathbb{R}$$

The isometry $A_{N_+} A_{N_-}$ respects the decomposition $H^{2,+}(\Sigma) \oplus H^{2,-}(\Sigma)$

Let $\alpha_1^+, \dots, \alpha_3^+, \alpha_1^-, \dots, \alpha_{19}^- \in (-\pi, \pi]$ be the angles through which $A_{N_+} A_{N_-} \otimes \mathbb{C}$ rotates $H^{2,+}(\Sigma, \mathbb{C})$ and $H^{2,-}(\Sigma, \mathbb{C})$, respectively

Assume $\vartheta \in (0, \pi)$, put $\rho = \pi - 2\vartheta \in (-\pi, \pi)$, and define

$$m_{\rho}(L; N_+, N_-) = \text{sign } \rho \left(\#\{j \mid \alpha_j^- \in \{\pi - |\rho|, \pi\}\} - 1 + 2 \#\{j \mid \alpha_j^- \in (\pi - |\rho|, \pi)\} \right)$$

Put $\text{sign } 0 = 0$, then $m_{\rho}(L; N_+, N_-) = 0$ for ordinary twisted connected sums

With respect to modified Atiyah-Patodi-Singer boundary conditions, define

$$\bar{\nu}(M_{\pm}, g) = 3\eta_{\text{APS}}(B_{M_{\pm}}; L_{B, \pm}) - 24\eta_{\text{APS}}(D_{M_{\pm}}; L_{D, \pm}) \in \mathbb{R}$$

The isometry $A_{N_+} A_{N_-}$ respects the decomposition $H^{2,+}(\Sigma) \oplus H^{2,-}(\Sigma)$

Let $\alpha_1^+, \dots, \alpha_3^+, \alpha_1^-, \dots, \alpha_{19}^- \in (-\pi, \pi]$ be the angles through which $A_{N_+} A_{N_-} \otimes \mathbb{C}$ rotates $H^{2,+}(\Sigma, \mathbb{C})$ and $H^{2,-}(\Sigma, \mathbb{C})$, respectively

Assume $\vartheta \in (0, \pi)$, put $\rho = \pi - 2\vartheta \in (-\pi, \pi)$, and define

$$m_{\rho}(L; N_+, N_-) = \text{sign } \rho \left(\#\{j \mid \alpha_j^- \in \{\pi - |\rho|, \pi\}\} - 1 + 2 \#\{j \mid \alpha_j^- \in (\pi - |\rho|, \pi)\} \right)$$

Put $\text{sign } 0 = 0$, then $m_{\rho}(L; N_+, N_-) = 0$ for ordinary twisted connected sums

From the gluing formulas for η -invariants by Bunke, Kirk-Lesch and others, we get

Theorem (Crowley-G-Nordström)

$$\bar{\nu}(M, g) = \bar{\nu}(M_+, g) + \bar{\nu}(M_-, g) - 72 \frac{\rho}{\pi} + 3m_{\rho}(L; N_+, N_-)$$

If $\Gamma_{\pm} \cong \{0\}$ or $\mathbb{Z}/2$ then $\bar{\nu}(M_{\pm}, g) = 0$

If $\Gamma_{\pm} \cong \{0\}$ or $\mathbb{Z}/2$ then $\bar{\nu}(M_{\pm}, g) = 0$

Example (Crowley-G-Nordström)

There exists a spin 7-manifold M with

$$H^1(M) \cong H^2(M) \cong 0, \quad H^4(M) \cong \mathbb{Z}^{97}, \quad \text{div } p_1(TM) = 4$$

admitting three different G_2 -holonomy metrics g_1, g_2, g_3 with

$$\bar{\nu}(M, g_1) = 0, \quad \bar{\nu}(M, g_2) = 36, \quad \bar{\nu}(M, g_3) = -36$$

If $\Gamma_{\pm} \cong \{0\}$ or $\mathbb{Z}/2$ then $\bar{\nu}(M_{\pm}, g) = 0$

Example (Crowley-G-Nordström)

There exists a spin 7-manifold M with

$$H^1(M) \cong H^2(M) \cong 0, \quad H^4(M) \cong \mathbb{Z}^{97}, \quad \text{div } p_1(TM) = 4$$

admitting three different G_2 -holonomy metrics g_1, g_2, g_3 with

$$\bar{\nu}(M, g_1) = 0, \quad \bar{\nu}(M, g_2) = 36, \quad \bar{\nu}(M, g_3) = -36$$

Hence, the G_2 -moduli space of M is disconnected

If $\Gamma_{\pm} \cong \{0\}$ or $\mathbb{Z}/2$ then $\bar{\nu}(M_{\pm}, g) = 0$

Example (Crowley-G-Nordström)

There exists a spin 7-manifold M with

$$H^1(M) \cong H^2(M) \cong 0, \quad H^4(M) \cong \mathbb{Z}^{97}, \quad \text{div } p_1(TM) = 4$$

admitting three different G_2 -holonomy metrics g_1, g_2, g_3 with

$$\bar{\nu}(M, g_1) = 0, \quad \bar{\nu}(M, g_2) = 36, \quad \bar{\nu}(M, g_3) = -36$$

Hence, the G_2 -moduli space of M is disconnected

The metric g_1 comes from a rectangular twisted connected sum

For g_2, g_3 , take $\Gamma_+ \cong \mathbb{Z}/2, \Gamma_- \cong \{0\}$ and $\vartheta = \frac{\pi}{4}$

Example (Crowley-G-Nordström)

There exists a spin 7-manifold M with

$$H^1(M) \cong H^2(M) \cong 0, \quad H^4(M) \cong \mathbb{Z}^{109}, \quad \text{div } p_1(TM) = 4$$

admitting three different G_2 -holonomy metrics g_1, g_2, g_3 with

$$\bar{\nu}(M, g_1) = 0, \quad \bar{\nu}(M, g_2) = 48, \quad \bar{\nu}(M, g_3) = -48$$

In particular, the ν -invariants agree and the G_2 -structures are homotopic

Example (Crowley-G-Nordström)

There exists a spin 7-manifold M with

$$H^1(M) \cong H^2(M) \cong 0, \quad H^4(M) \cong \mathbb{Z}^{109}, \quad \text{div } p_1(TM) = 4$$

admitting three different G_2 -holonomy metrics g_1, g_2, g_3 with

$$\bar{\nu}(M, g_1) = 0, \quad \bar{\nu}(M, g_2) = 48, \quad \bar{\nu}(M, g_3) = -48$$

In particular, the ν -invariants agree and the G_2 -structures are homotopic. Hence, one homotopy class of G_2 -structures can give rise to several connected components of the G_2 -moduli space.

Example (Crowley-G-Nordström)

There exists a spin 7-manifold M with

$$H^1(M) \cong H^2(M) \cong 0, \quad H^4(M) \cong \mathbb{Z}^{109}, \quad \text{div } p_1(TM) = 4$$

admitting three different G_2 -holonomy metrics g_1, g_2, g_3 with

$$\bar{\nu}(M, g_1) = 0, \quad \bar{\nu}(M, g_2) = 48, \quad \bar{\nu}(M, g_3) = -48$$

In particular, the ν -invariants agree and the G_2 -structures are homotopic. Hence, one homotopy class of G_2 -structures can give rise to several connected components of the G_2 -moduli space.

The metric g_1 comes from a rectangular twisted connected sum.

For g_2, g_3 , take $\Gamma_+ \cong \Gamma_- \cong \mathbb{Z}/2$ and $\vartheta = \frac{\pi}{6}$.

Let (M, g) be an extra twisted connected sum with $\Gamma_{\pm} \cong \mathbb{Z}/k_{\pm}$
where $k_+, k_- \in \{1, 2\}$ and $\vartheta \in \{\pm\frac{\pi}{6}, \pm\frac{\pi}{4}, \pm\frac{\pi}{3}, \frac{\pi}{2}\}$
In this case, 3 divides $\bar{\nu}(M, g) = -72\frac{\rho}{\pi} + 3m_{\rho}(L; N_+, N_-)$

Let (M, g) be an extra twisted connected sum with $\Gamma_{\pm} \cong \mathbb{Z}/k_{\pm}$
 where $k_+, k_- \in \{1, 2\}$ and $\vartheta \in \{\pm\frac{\pi}{6}, \pm\frac{\pi}{4}, \pm\frac{\pi}{3}, \frac{\pi}{2}\}$

In this case, 3 divides $\bar{\nu}(M, g) = -72\frac{\rho}{\pi} + 3m_{\rho}(L; N_+, N_-)$

Note that the G_2 -bordism group in dimension 7 is $\Omega_{G_2}^7 \cong \mathbb{Z}/3$

The ν -invariant mod 3 is an isomorphism $\nu: \Omega_{G_2}^7 \rightarrow \mathbb{Z}/3$

Hence (M, σ) is **G_2 -nullbordant** if and only if $3 \mid \nu(M, \sigma)$

Let (M, g) be an extra twisted connected sum with $\Gamma_{\pm} \cong \mathbb{Z}/k_{\pm}$
 where $k_+, k_- \in \{1, 2\}$ and $\vartheta \in \{\pm\frac{\pi}{6}, \pm\frac{\pi}{4}, \pm\frac{\pi}{3}, \frac{\pi}{2}\}$

In this case, 3 divides $\bar{\nu}(M, g) = -72\frac{\rho}{\pi} + 3m_{\rho}(L; N_+, N_-)$

Note that the G_2 -bordism group in dimension 7 is $\Omega_{G_2}^7 \cong \mathbb{Z}/3$

The ν -invariant mod 3 is an isomorphism $\nu: \Omega_{G_2}^7 \rightarrow \mathbb{Z}/3$

Hence (M, σ) is **G_2 -nullbordant** if and only if $3 \mid \nu(M, \sigma)$

Question. Is the G_2 -bordism class of (M, σ) an obstruction against holonomy G_2 ?

Let (M, g) be an extra twisted connected sum with $\Gamma_{\pm} \cong \mathbb{Z}/k_{\pm}$
 where $k_+, k_- \in \{1, 2\}$ and $\vartheta \in \{\pm\frac{\pi}{6}, \pm\frac{\pi}{4}, \pm\frac{\pi}{3}, \frac{\pi}{2}\}$

In this case, 3 divides $\bar{\nu}(M, g) = -72\frac{\rho}{\pi} + 3m_{\rho}(L; N_+, N_-)$

Note that the G_2 -bordism group in dimension 7 is $\Omega_{G_2}^7 \cong \mathbb{Z}/3$

The ν -invariant mod 3 is an isomorphism $\nu: \Omega_{G_2}^7 \rightarrow \mathbb{Z}/3$

Hence (M, σ) is **G_2 -nullbordant** if and only if $3 \mid \nu(M, \sigma)$

Question. Is the G_2 -bordism class of (M, σ) an obstruction against holonomy G_2 ?

Answer. No, there are examples with $3 \nmid \nu(M, \sigma)$.

This is our motivation to consider more complicated extra twisted connected sums

Let (M, g) be an extra twisted connected sum with $\Gamma_{\pm} \cong \mathbb{Z}/k_{\pm}$
 where $k_+, k_- \in \{1, 2\}$ and $\vartheta \in \{\pm\frac{\pi}{6}, \pm\frac{\pi}{4}, \pm\frac{\pi}{3}, \frac{\pi}{2}\}$

In this case, 3 divides $\bar{\nu}(M, g) = -72\frac{\rho}{\pi} + 3m_{\rho}(L; N_+, N_-)$

Note that the G_2 -bordism group in dimension 7 is $\Omega_{G_2}^7 \cong \mathbb{Z}/3$

The ν -invariant mod 3 is an isomorphism $\nu: \Omega_{G_2}^7 \rightarrow \mathbb{Z}/3$

Hence (M, σ) is **G_2 -nullbordant** if and only if $3 \mid \nu(M, \sigma)$

Question. Is the G_2 -bordism class of (M, σ) an obstruction against holonomy G_2 ?

Answer. No, there are examples with $3 \nmid \nu(M, \sigma)$.

This is our motivation to consider more complicated extra twisted connected sums

Note. Recall that with $k_+ \geq 3$ or $k_- \geq 3$, we can get gluing angles $\vartheta \notin \mathbb{Q}\pi$

Because $\bar{\nu}(M, g) \in \mathbb{Z}$, expect $\bar{\nu}(M_{\pm}, g) \neq 0$ if $k_{\pm} > 2$

Let $M_{\pm} = V_{\pm} \times S^1_{\pm, \text{ext}}$, rescale $S^1_{\pm, \text{ext}}$ by $a > 0$ to get $M_{\pm, a}$
The limit $a \rightarrow 0$ is called **adiabatic limit**

Let $M_{\pm} = V_{\pm} \times S^1_{\pm, \text{ext}}$, rescale $S^1_{\pm, \text{ext}}$ by $a > 0$ to get $M_{\pm, a}$

The limit $a \rightarrow 0$ is called **adiabatic limit**

From the adiabatic limit theorems of Bismut-Cheeger, Dai, G, we deduce

Theorem (G-Nordström)

Let $\gamma \in \Gamma_{\pm}$ be the generator that acts by $\frac{2\pi}{k_{\pm}}$ on $S^1_{\pm, \text{ext}}$

Let $V_{\pm, j}$ be the set of isolated fixpoints of γ^j on V_{\pm}

Let $e^{i\alpha_{j,1}(p)}$, $e^{i\alpha_{j,2}(p)}$, $e^{i\alpha_{j,3}(p)}$ be the eigenvalues of γ^j on $T_p V_{\pm}$
with $\alpha_{j,1}(p) + \alpha_{j,2}(p) + \alpha_{j,3}(p) = 0$. Then

$$\lim_{a \rightarrow 0} \bar{\nu}(M_{\pm, a}) = \frac{3}{k_{\pm}} \sum_{j=1}^{k_{\pm}-1} \cot \frac{\pi j}{k_{\pm}} \sum_{p \in V_{\pm, j}} \frac{\cos \frac{\alpha_{j,1}(p)}{2} \cos \frac{\alpha_{j,2}(p)}{2} \cos \frac{\alpha_{j,3}(p)}{2} - 1}{\sin \frac{\alpha_{j,1}(p)}{2} \sin \frac{\alpha_{j,2}(p)}{2} \sin \frac{\alpha_{j,3}(p)}{2}} \in \mathbb{Q}$$

Let $M_{\pm} = V_{\pm} \times S^1_{\pm, \text{ext}}$, rescale $S^1_{\pm, \text{ext}}$ by $a > 0$ to get $M_{\pm, a}$

The limit $a \rightarrow 0$ is called **adiabatic limit**

From the adiabatic limit theorems of Bismut-Cheeger, Dai, G, we deduce

Theorem (G-Nordström)

Let $\gamma \in \Gamma_{\pm}$ be the generator that acts by $\frac{2\pi}{k_{\pm}}$ on $S^1_{\pm, \text{ext}}$

Let $V_{\pm, j}$ be the set of isolated fixpoints of γ^j on V_{\pm}

Let $e^{i\alpha_{j,1}(p)}$, $e^{i\alpha_{j,2}(p)}$, $e^{i\alpha_{j,3}(p)}$ be the eigenvalues of γ^j on $T_p V_{\pm}$
with $\alpha_{j,1}(p) + \alpha_{j,2}(p) + \alpha_{j,3}(p) = 0$. Then

$$\lim_{a \rightarrow 0} \bar{\nu}(M_{\pm, a}) = \frac{3}{k_{\pm}} \sum_{j=1}^{k_{\pm}-1} \cot \frac{\pi j}{k_{\pm}} \sum_{p \in V_{\pm, j}} \frac{\cos \frac{\alpha_{j,1}(p)}{2} \cos \frac{\alpha_{j,2}(p)}{2} \cos \frac{\alpha_{j,3}(p)}{2} - 1}{\sin \frac{\alpha_{j,1}(p)}{2} \sin \frac{\alpha_{j,2}(p)}{2} \sin \frac{\alpha_{j,3}(p)}{2}} \in \mathbb{Q}$$

We will focus on examples without isolated fixpoints

By theorems of Bismut-Cheeger, Dai-Freed, the variational formula for the η -invariant of a Dirac type operator a manifold with boundary consists of

- ▶ the integral of a Chern-Simons form over the interior
- ▶ the degree-1-component of an η -form on the boundary

By theorems of Bismut-Cheeger, Dai-Freed, the variational formula for the η -invariant of a Dirac type operator a manifold with boundary consists of

- ▶ the integral of a Chern-Simons form over the interior
- ▶ the degree-1-component of an η -form on the boundary

The interior contribution vanishes because $M_{\pm,a}$ is locally a product

By theorems of Bismut-Cheeger, Dai-Freed, the variational formula for the η -invariant of a Dirac type operator a manifold with boundary consists of

- ▶ the integral of a Chern-Simons form over the interior
- ▶ the degree-1-component of an η -form on the boundary

The interior contribution vanishes because $M_{\pm,a}$ is locally a product

The boundary contribution can be expressed in terms of the η -form $\tilde{\eta}(\mathbb{A})$ of the family of tori $(S_{\pm,\text{int}}^1 \times aS_{\pm,\text{ext}}^1)/\Gamma_{\pm}$ for $a \in (0, s_{\pm})$

Theorem (G-Nordström)

$$\bar{\nu}(M_{\pm}) - \lim_{a \rightarrow 0} \bar{\nu}(M_{\pm,a}) = F_{k_{\pm}, \epsilon_{\pm}}(s_{\pm}) = 288 \int_0^{s_{\pm}} \tilde{\eta}(\mathbb{A})$$

By theorems of Bismut-Cheeger, Dai-Freed, the variational formula for the η -invariant of a Dirac type operator a manifold with boundary consists of

- ▶ the integral of a Chern-Simons form over the interior
- ▶ the degree-1-component of an η -form on the boundary

The interior contribution vanishes because $M_{\pm,a}$ is locally a product

The boundary contribution can be expressed in terms of the η -form $\tilde{\eta}(\mathbb{A})$ of the family of tori $(S_{\pm,\text{int}}^1 \times aS_{\pm,\text{ext}}^1)/\Gamma_{\pm}$ for $a \in (0, s_{\pm})$

Theorem (G-Nordström)

$$\bar{\nu}(M_{\pm}) - \lim_{a \rightarrow 0} \bar{\nu}(M_{\pm,a}) = F_{k_{\pm}, \epsilon_{\pm}}(s_{\pm}) = 288 \int_0^{s_{\pm}} \tilde{\eta}(\mathbb{A})$$

Bismut-Cheeger also give a formula for $\tilde{\eta}(\mathbb{A})$ as a sum over \mathbb{Z}^2 depending on a

Represent the torus $\mathbb{C}/(\mathbb{Z} + \tau\mathbb{Z})$ by $\tau \in \mathcal{H} = \{z \in \mathbb{C} \mid \text{Im}(z) > 0\}$

Then $\tilde{\eta} \in \Omega^1(\mathcal{H})$ is $SL(2, \mathbb{Z})$ -invariant and satisfies $d\tilde{\eta} = -\frac{1}{4\pi} dA_{\text{hyp}}$ by Bismut's family index theorem

Represent the torus $\mathbb{C}/(\mathbb{Z} + \tau\mathbb{Z})$ by $\tau \in \mathcal{H} = \{z \in \mathbb{C} \mid \text{Im}(z) > 0\}$

Then $\tilde{\eta} \in \Omega^1(\mathcal{H})$ is $SL(2, \mathbb{Z})$ -invariant and satisfies $d\tilde{\eta} = -\frac{1}{4\pi} dA_{\text{hyp}}$ by Bismut's family index theorem


Idea. Use hyperbolic geometry
to compute $F_{k_-, \varepsilon_-}(s_-) + F_{k_+, \varepsilon_+}(s_+)$

Represent the torus $\mathbb{C}/(\mathbb{Z} + \tau\mathbb{Z})$ by $\tau \in \mathcal{H} = \{z \in \mathbb{C} \mid \text{Im}(z) > 0\}$

Then $\tilde{\eta} \in \Omega^1(\mathcal{H})$ is $SL(2, \mathbb{Z})$ -invariant and satisfies $d\tilde{\eta} = -\frac{1}{4\pi} dA_{\text{hyp}}$ by Bismut's family index theorem

Idea. Use hyperbolic geometry to compute $F_{k_-, \varepsilon_-}(s_-) + F_{k_+, \varepsilon_+}(s_+)$

Adiabatic limits—geodesic rays




Represent the torus $\mathbb{C}/(\mathbb{Z} + \tau\mathbb{Z})$ by $\tau \in \mathcal{H} = \{z \in \mathbb{C} \mid \text{Im}(z) > 0\}$

Then $\tilde{\eta} \in \Omega^1(\mathcal{H})$ is $SL(2, \mathbb{Z})$ -invariant and satisfies $d\tilde{\eta} = -\frac{1}{4\pi} dA_{\text{hyp}}$ by Bismut's family index theorem

Idea. Use hyperbolic geometry to compute $F_{k_-, \varepsilon_-}(s_-) + F_{k_+, \varepsilon_+}(s_+)$

Adiabatic limits—geodesic rays

$\tilde{\eta}(\mathbb{A}) = 0$ along families of rectangular ($k = 1$) or rhombic ($k = 2$) tori



Represent the torus $\mathbb{C}/(\mathbb{Z} + \tau\mathbb{Z})$ by $\tau \in \mathcal{H} = \{z \in \mathbb{C} \mid \text{Im}(z) > 0\}$

Then $\tilde{\eta} \in \Omega^1(\mathcal{H})$ is $SL(2, \mathbb{Z})$ -invariant and satisfies $d\tilde{\eta} = -\frac{1}{4\pi} dA_{\text{hyp}}$ by Bismut's family index theorem

Idea. Use hyperbolic geometry to compute $F_{k_-, \varepsilon_-}(s_-) + F_{k_+, \varepsilon_+}(s_+)$


Adiabatic limits—geodesic rays

$\tilde{\eta}(\mathbb{A}) = 0$ along families of rectangular

($k = 1$) or rhombic ($k = 2$) tori

Cusps—families of adiabatic limits

Their contribution can be computed using a formula by Bunke and Ma



Represent the torus $\mathbb{C}/(\mathbb{Z} + \tau\mathbb{Z})$ by $\tau \in \mathcal{H} = \{z \in \mathbb{C} \mid \text{Im}(z) > 0\}$

Then $\tilde{\eta} \in \Omega^1(\mathcal{H})$ is $SL(2, \mathbb{Z})$ -invariant and satisfies $d\tilde{\eta} = -\frac{1}{4\pi} dA_{\text{hyp}}$ by Bismut's family index theorem

Idea. Use hyperbolic geometry to compute $F_{k_-, \varepsilon_-}(s_-) + F_{k_+, \varepsilon_+}(s_+)$

Adiabatic limits—geodesic rays

$\tilde{\eta}(\mathbb{A}) = 0$ along families of rectangular


($k = 1$) or rhombic ($k = 2$) tori

Cusps—families of adiabatic limits

Their contribution can be computed using a formula by Bunke and Ma

Compute $F_{k_-, \varepsilon_-}(s_-) + F_{k_+, \varepsilon_+}(s_+)$ using Stokes' theorem from the cusp contributions and the hyperbolic area formula

The angle 2ϑ at the finite corner cancels $-72\frac{\rho}{\pi}$ in the gluing formula



The logarithm of the **Dedekind η -function** is given by

$$L(\tau) = \frac{\pi i \tau}{12} - \sum_{n=1}^{\infty} \sum_{d|n} d^{-1} e^{2\pi i n \tau}$$

Theorem (G-Nordström-Zagier)

There exists a constant $c_{k_{\pm}, \varepsilon_{\pm}} \in \mathbb{Q}$ such that

$$F_{k_{\pm}, \varepsilon_{\pm}}(s_{\pm}) = \frac{144}{\pi} \left(iL\left(\frac{s_{\pm} i + \varepsilon_{\pm}}{k_{\pm}}\right) - iL\left(\frac{s_{\pm} i - \varepsilon_{\pm}}{k_{\pm}}\right) + c_{k_{\pm}, \varepsilon_{\pm}} \right)$$

The logarithm of the **Dedekind η -function** is given by

$$L(\tau) = \frac{\pi i \tau}{12} - \sum_{n=1}^{\infty} \sum_{d|n} d^{-1} e^{2\pi i n \tau}$$

Theorem (G-Nordström-Zagier)

There exists a constant $c_{k_{\pm}, \varepsilon_{\pm}} \in \mathbb{Q}$ such that

$$F_{k_{\pm}, \varepsilon_{\pm}}(\mathbf{s}_{\pm}) = \frac{144}{\pi} \left(iL\left(\frac{\mathbf{s}_{\pm} i + \varepsilon_{\pm}}{k_{\pm}}\right) - iL\left(\frac{\mathbf{s}_{\pm} i - \varepsilon_{\pm}}{k_{\pm}}\right) + c_{k_{\pm}, \varepsilon_{\pm}} \right)$$

Compute the variational term using the functional equations

$$L(\tau + 1) = \frac{\pi i}{12} + L(\tau) \quad \text{and} \quad L\left(-\frac{1}{\tau}\right) = \frac{1}{2} \log \frac{\tau}{i} + L(\tau)$$

Example. The example with $\cos \vartheta = \frac{1}{\sqrt{6}}$ has $\bar{\nu}(M, g) = -65$
In particular, $3 \nmid \bar{\nu}(M, g)$, so it is not G_2 -nullbordant

Example. The example with $\cos \vartheta = \frac{1}{\sqrt{6}}$ has $\bar{\nu}(M, g) = -65$
In particular, $3 \nmid \bar{\nu}(M, g)$, so it is not G_2 -nullbordant

Conjecture

All values in $\mathbb{Z}/48$ occur as ν -invariants of G_2 -holonomy metrics

Example. The example with $\cos \vartheta = \frac{1}{\sqrt{6}}$ has $\bar{\nu}(M, g) = -65$
In particular, $3 \nmid \bar{\nu}(M, g)$, so it is not G_2 -nullbordant

Conjecture

All values in $\mathbb{Z}/48$ occur as ν -invariants of G_2 -holonomy metrics

Questions

- ▶ How many different G_2 -metrics exist on one 7-manifold?
- ▶ Are different G_2 -metrics on a fixed 7-manifold G_2 -bordant?

Example. The example with $\cos \vartheta = \frac{1}{\sqrt{6}}$ has $\bar{\nu}(M, g) = -65$
In particular, $3 \nmid \bar{\nu}(M, g)$, so it is not G_2 -nullbordant

Conjecture

All values in $\mathbb{Z}/48$ occur as ν -invariants of G_2 -holonomy metrics

Questions

- ▶ How many different G_2 -metrics exist on one 7-manifold?
- ▶ Are different G_2 -metrics on a fixed 7-manifold G_2 -bordant?

Construct more examples

- ▶ Find more asymptotically cylindrical Calabi-Yau manifolds
- ▶ Understand their moduli space, make the K3 surfaces match
- ▶ Consider other constructions

Thanks for your attention!