

Deformed G_2 -instantons

Jason D. Lotay

Oxford

18 September 2020

(Joint work with Goncalo Oliveira)

Overview

Deformed G_2 -instantons

- special connections in 7 dimensions
- “mirror” to calibrated cycles \rightsquigarrow enumerative invariants?
- critical points of Chern–Simons-type functional \mathcal{F}

Results

- first non-trivial examples
- examples detect different G_2 -structures (including nearly parallel and isometric)
- deformation theory: obstructions and topology of moduli space
- relation to \mathcal{F}

Deformed G_2 -instantons

X^7 , φ G_2 -structure, $d * \varphi = 0$, connection A on bundle over X

Definition (J.-H. Lee–N.C. Leung)

A *deformed G_2 -instanton* (dG_2) \Leftrightarrow curvature F_A satisfies

$$F_A \wedge * \varphi + \frac{1}{6} F_A^3 = 0$$

$\mathbb{R}^7 = \mathbb{R}^3 \oplus \mathbb{R}^4$, $x = (x_1, x_2, x_3) \in \mathbb{R}^3$, $y = (y_0, y_1, y_2, y_3) \in \mathbb{R}^4$,
 $\varphi(u, v, w) = g_{\mathbb{R}^7}(u \times v, w) \rightsquigarrow$ **“mirror calibrated cycles”**

- $A = i(a_j(x)dx_j + u_k(x)dy_k)$ dG_2 -instanton \Leftrightarrow
 Graph(u) **associative** and $i(a_j dx_j)$ **flat**
- $A = i(v_j(y)dx_j + b_k(y)dy_k)$ dG_2 -instanton \Leftrightarrow
 Graph(v) **coassociative** and $i(b_k dy_k)$ **anti-self-dual (ASD)**

\rightsquigarrow primary interest in **U(1)-connections**

Lower dimensions

4 dimensions: $\pi : X^7 \rightarrow Z^4$, Z ASD Einstein, connection B on Z

Lemma

π^*B dG₂-instanton $\Leftrightarrow B$ ASD $\Leftrightarrow \pi^*B$ G₂-instanton

6 dimensions: $\pi : X^7 \rightarrow Y^6$, Y Calabi–Yau 3-fold
 ω Kähler form on Y , connection B on Y

Lemma

π^*B dG₂-instanton $\Leftrightarrow B$ *deformed Hermitian–Yang–Mills*

$$F_B^{(0,2)} = 0 \quad \text{and} \quad \text{Im}((\omega + F_B)^3) = 0.$$

- Conjecture: existence of dHYM \leftrightarrow stability condition

3-Sasakian 7-manifolds

Definition

(X^7, g^{ts}) 3-Sasakian $\Leftrightarrow (\mathbb{R}^+ \times X^7, g = dr^2 + r^2 g^{ts})$ hyperkähler

Fact: \exists infinitely many 3-Sasakian 7-manifolds

Examples: S^7 , Aloff–Wallach $(SU(3) \times SU(2))/(U(1) \times SU(2))$

- $V^3 \rightarrow X^7 \rightarrow Z^4$, $V = SU(2)/\Gamma$, Z ASD Einstein
- left-invariant coframe η_1, η_2, η_3 on V
- $\omega_1, \omega_2, \omega_3$ orthogonal self-dual 2-forms on Z with length 2
- \rightsquigarrow two 1-parameter families of 3-forms for $t > 0$:

$$\varphi_t^\pm = \pm t^3 \eta_1 \wedge \eta_2 \wedge \eta_3 - t \eta_1 \wedge \omega_1 - t \eta_2 \wedge \omega_2 \mp t \eta_3 \wedge \omega_3$$



- φ_t^\pm induces $g_t = t^2 g_V + g_Z \Rightarrow \varphi_t^+$ and φ_t^- isometric
- $\varphi^{ts} = \varphi_1^-$ nearly parallel inducing g^{ts}
- $\varphi^{np} = \varphi_{1/\sqrt{5}}^+$ nearly parallel inducing g^{np} “squashed” Einstein metric, cone has $\text{Hol} = \text{Spin}(7)$

Non-trivial examples

- (X^7, g^{ts}) 3-Sasakian, φ_t^\pm inducing g_t
- $a = (a_1, a_2, a_3) \in \mathbb{R}^3 \rightsquigarrow A = i(a_1\eta_1 + a_2\eta_2 + a_3\eta_3)$ connection on trivial line bundle, $r = |a|$ “distance to trivial connection”

Theorem (L.–Oliveira)

A dG_2 -instanton on $(X^7, \varphi_t^\pm) \Leftrightarrow$



Deformation theory

Theorem (L.–Oliveira)

- *Non-trivial* dG_2 -instantons constructed for φ^{ts} and φ^{np} are *obstructed*
- *Trivial* dG_2 -instanton *unobstructed* for φ^{ts} and φ^{np} but *obstructed* for $\varphi_{1/\sqrt{2}}^{\pm}$

Proof: (Kawai–Yamamoto) \rightsquigarrow unobstructed \Leftrightarrow rigid and isolated

At $t = \frac{1}{\sqrt{2}}$: \exists infinitesimal deformation of trivial dG_2 -instanton \Rightarrow obstructed (proof uses functional \mathcal{F})

Corollary

Moduli space of dG_2 -instantons on trivial line bundle for φ^{ts} and φ^{np} contains at least two components of different dimensions

Chern–Simons-type functional

- A_0 reference connection on line bundle L on (X^7, φ) , $d*\varphi = 0$
- A connection on L
- $\rightsquigarrow \mathbb{A} = A_0 + s(A - A_0)$ connection on L over $X \times [0, 1]$
- \rightsquigarrow curvature \mathbb{F}

Proposition (Karigiannis–N.C. Leung)

A dG₂-instanton \Leftrightarrow A critical point of functional





$$\mathcal{F}(A) = \int_{X \times [0,1]} e^{\mathbb{F} + *\varphi}$$

Recall: our examples $A = i(a_1\eta_1 + a_2\eta_2 + a_3\eta_3)$

\rightsquigarrow restriction of \mathcal{F} is function of two variables x and y :





$$x = a_3 \quad \text{and} \quad y^2 = a_1^2 + a_2^2$$

\mathcal{F} for φ_1^+ and $\varphi^{ts} = \varphi_1^-$ with level sets



- trivial connection only critical point \Rightarrow local maximum
- otherwise trivial connection saddle point
- non-trivial dG_2 -instantons are local maxima

\mathcal{F} for $\varphi^{np} = \varphi_{1/\sqrt{5}}^+$ and $\varphi_{1/\sqrt{5}}^-$ with level sets



- trivial connection is local minimum
- continuous families of dG_2 -instantons are local maxima
- two isolated examples are saddle points

Questions

- non-trivial dG_2 -instantons for holonomy G_2 -manifolds?
- (adiabatic) limits of dG_2 -instantons?
- dependence of moduli space on G_2 -structure?
- “mirror count”?
- applications of \mathcal{F} to compactness or deformation theory?
- Spin(7) version?