

## Some remarks on contact Calabi–Yau 7-manifolds

Jason D. Lotay

University of Oxford

10 September 2021

(Joint work with Henrique Sá Earp and Julieth Saavedra)

# Calabi–Yau and $G_2$ geometry: circle bundles

## 6D: Calabi–Yau geometry

circle bundles  
↔

$$(Z^6, h, \omega, \Upsilon)$$

metric  $h$

Kähler form  $\omega$

holomorphic volume form  $\Upsilon$

## 7D: $G_2$ geometry

$$\mathcal{S}_\epsilon^1 \longrightarrow (M^7, g, \eta, \varphi)$$

$$\downarrow$$

$$Z$$

metric  $g = \epsilon^2 \eta^2 + h$

connection 1-form  $\eta$

$G_2$ -structure  $\varphi = \epsilon \eta \wedge \omega + \text{Re } \Upsilon$

**Limit**  $\epsilon \rightarrow 0$

- **Geometry:** collapsing/“nearly collapsed”
- **Analysis:** series expansion in  $\epsilon$
- **Physics:** M-theory ↔ String Theory

# Contact Calabi–Yau 7-manifolds

**Example:**  $M^7 = \{(z_0, z_1, z_2, z_3, z_4) \in \mathbb{C}^5 : \sum_{k=0}^4 z_k^5 = 0\} \cap \mathcal{S}^9$

- $Z = \{[z_0, z_1, z_2, z_3, z_4] \in \mathbb{C}\mathbb{P}^4 : \sum_{k=0}^4 z_k^5 = 0\}$  Fermat quintic
- connection 1-form  $\eta$  such that  $d\eta = \omega$

## Definition

$(M^7, g, \eta, \Upsilon)$  *contact Calabi–Yau 7-manifold*:

- $(M^7, g)$  Sasakian with contact form  $\eta$
- $E = \ker \eta$  with transverse Calabi–Yau structure  $(h, \omega = d\eta, \Upsilon)$

$\rightsquigarrow$   $G_2$ -structure  $\varphi = \epsilon\eta \wedge \omega + \operatorname{Re} \Upsilon$  with  $g = \epsilon^2\eta^2 + h$

$\rightsquigarrow * \varphi = \frac{1}{2}\omega^2 - \epsilon\eta \wedge \operatorname{Im} \Upsilon \Rightarrow d * \varphi = 0$

$\rightsquigarrow \varphi$  **coclosed** with  $d\varphi = \epsilon\omega^2$

# Geometric flows

## Laplacian flow

$$\frac{\partial \varphi_t}{\partial t} = \Delta_t \varphi_t = (dd_t^* + d_t^*d)\varphi_t$$

## Laplacian coflow

$$\frac{\partial *_t \varphi_t}{\partial t} = (dd_t^* + d_t^*d) *_t \varphi_t = dd_t^* *_t \varphi_t$$

- **Critical point:** torsion-free  $G_2$ -structure  $\rightsquigarrow$  Ricci-flat metric with holonomy in  $G_2$
- Laplacian (co)flow restricted to (co)closed  $G_2$ -structures = gradient flow of **Hitchin volume functional** on cohomology class of  $\varphi_t$  ( $*_t \varphi_t$ )

### Theorem (L.–Sá Earp–Saavedra)

$(M^7, g, \eta, \Upsilon)$  contact Calabi–Yau (with  $\omega = d\eta$ )

$$\varphi_0 = \epsilon \eta \wedge \omega + \operatorname{Re} \Upsilon \quad \text{and} \quad *_0 \varphi_0 = \frac{1}{2} \omega^2 - \epsilon \eta \wedge \operatorname{Im} \Upsilon$$

- Laplacian flow has **finite-time singularity** at  $t = \frac{1}{8\epsilon^2}$
- Laplacian coflow exists for all  $t > 0$  and has an **infinite-time singularity**

$M$  compact  $\Rightarrow \operatorname{Vol}(M, g_t) \rightarrow \infty$

# Heterotic $G_2$ system (or $G_2$ -Hull–Strominger system)

**Recall:** connection  $A$  on  $(M^7, \varphi)$   $G_2$ -instanton  $\Leftrightarrow F_A \wedge \varphi = - * F_A \Leftrightarrow F_A \wedge * \varphi = 0$

## Definition

$\varphi$   $G_2$ -structure on  $M^7$ ,  $A$  connection on  $E$  over  $M$ ,  $B$  connection on  $TM$  and  $\alpha' > 0$

$\rightsquigarrow (\varphi, (A, E), B, \alpha')$  solution to **heterotic  $G_2$  system** (with vanishing dilaton):

- $\varphi$  *coclosed*  $\rightsquigarrow d\varphi = \frac{7}{3}\lambda * \varphi - *H$  ( $H$  flux,  $H \wedge \varphi = 0$  &  $H \wedge * \varphi = 0$ )
- $A$   $G_2$ -instanton and  $B$   $G_2$ -instanton (up to  $O(\alpha')^2$  corrections)
- **anomaly-free condition** (or heterotic Bianchi identity):  $dH = \frac{\alpha'}{4}(\text{tr } F_A^2 - \text{tr } F_B^2)$

## Theorem (L.–Sá Earp)

$\forall \alpha' > 0$   $(M^7, g, \eta, \Upsilon)$  contact Calabi–Yau admits solution  $(\varphi, (A, \ker \eta), B, \alpha')$  to heterotic  $G_2$  system where

$$\varphi = \epsilon \eta \wedge \omega + \text{Re } \Upsilon, \quad dH \neq 0 \quad \text{and} \quad \epsilon \rightarrow 0 \quad \text{as} \quad \alpha' \rightarrow 0$$

## Flow ansatz

## Laplacian flow

$$\frac{\partial \varphi_t}{\partial t} = \Delta_t \varphi_t$$

## Laplacian coflow

$$\frac{\partial *_t \varphi_t}{\partial t} = \Delta_t *_t \varphi_t$$

$(M^7, g, \eta, \Upsilon)$  contact Calabi–Yau,  $\omega = d\eta \rightsquigarrow$  consider

$$\varphi_t = f_t h_t^2 \eta \wedge \omega + h_t^3 \operatorname{Re} \Upsilon \quad \text{and} \quad *_t \varphi_t = \frac{1}{2} h_t^4 \omega^2 - f_t h_t^3 \eta \wedge \operatorname{Im} \Upsilon$$

$$\Delta_t \varphi_t = \frac{4f_t^3}{h_t^2} \eta \wedge \omega \quad \text{and} \quad \Delta_t *_t \varphi_t = 2f_t^2 \omega^2$$

$\rightsquigarrow$  ansatz preserved,  $d *_t \varphi_t = 0 \forall t$  and flows  $\leftrightarrow$  ODE systems

**Note:**

- for Laplacian coflow  $[_t \varphi_t] = [_0 \varphi_0] \forall t$
- but for Laplacian flow  $[_t \varphi_t] \neq [_0 \varphi_0]$  in general

# Singularity analysis

Recall:  $g_0 = \epsilon^2 \eta^2 + h$  on  $M^7$  and  $\nabla_t \varphi_t$  encoded by **torsion** 2-tensor  $T_t$

- Laplacian flow for  $t \in (-\infty, \frac{1}{8\epsilon^2})$  (ancient solution)
- Laplacian coflow for  $t \in (-\frac{1}{10\epsilon^2}, \infty)$  (immortal solution)  $\rightsquigarrow$  **does not converge**

$M$  compact  $\rightsquigarrow$  (L.-Wei, G. Chen) suggest singularity formation controlled by

$$\Lambda(t) = \sup_M (|Rm_t|_t^2 + |T_t|_t^4 + |\nabla_t T_t|_t^2)^{1/2}$$

**Laplacian flow:**

- $\Lambda(t) \rightarrow \infty$  as  $t \rightarrow \frac{1}{8\epsilon^2}$  but  $\lim_{t \rightarrow \frac{1}{8\epsilon^2}} (\frac{1}{8\epsilon^2} - t)\Lambda(t) < \infty$  (“Type I” / rapidly forming)
- volume normalized flow  $\rightsquigarrow$  converges to  $\mathbb{R}$  as  $t \rightarrow \frac{1}{8\epsilon^2}$  and to  $\mathbb{C}^3$  as  $t \rightarrow -\infty$

**Laplacian coflow** (with non-flat  $h$ ):

- $\Lambda(t) \rightarrow 0$  as  $t \rightarrow \infty$  but  $\lim_{t \rightarrow \infty} t\Lambda(t) = \infty$  (“Type IIb” / slowly forming)
- volume normalized flow  $\rightsquigarrow$  converges to  $\mathbb{C}^3$  as  $t \rightarrow \infty$  and to  $\mathbb{R}$  as  $t \rightarrow -\frac{1}{10\epsilon^2}$

# Solving the heterotic $G_2$ system: $G_2$ -structure and $G_2$ -instanton A

$(M^7, g, \eta, \Upsilon)$  contact Calabi–Yau

**Want:**  $(\varphi, (A, E), B, \alpha')$

- $\alpha' > 0$  arbitrary ✓
- $\epsilon = \epsilon(\alpha') > 0 \rightsquigarrow$

$$\varphi_\epsilon = \epsilon \eta \wedge \omega + \operatorname{Re} \Upsilon \quad \text{and} \quad * \varphi_\epsilon = \frac{1}{2} \omega^2 - \epsilon \eta \wedge \operatorname{Im} \Upsilon$$

$$\rightsquigarrow d * \varphi_\epsilon = 0 \quad \checkmark \quad \text{and flux } H_\epsilon = -\epsilon^2 \eta \wedge \omega + \epsilon \operatorname{Re} \Upsilon \rightsquigarrow dH_\epsilon = -\epsilon^2 \omega^2 \neq 0 \quad \checkmark$$

- $E = \ker \eta$ ,  $A$  transverse connection  $\rightsquigarrow$

$$F_A \wedge \omega^2 = 0, \quad F_A \wedge \Upsilon = 0 \quad \rightsquigarrow \quad F_A \wedge * \varphi_\epsilon = 0 \quad \checkmark$$

- **need**  $B$  on  $TM$  so that

$$F_B \wedge * \varphi_\epsilon = O(\alpha')^2 \quad \text{and} \quad dH_\epsilon = -\epsilon^2 \omega^2 = \frac{\alpha'}{4} (\operatorname{tr} F_A^2 - \operatorname{tr} F_B^2)$$



# Solving the heterotic $G_2$ system: approximate $G_2$ -instanton $B$ and anomaly-free condition

**Recall:** “Bismut connection”  $B_\epsilon^+$  for  $\varphi_\epsilon \rightsquigarrow$

$$\nabla_{B_\epsilon^+} g_\epsilon = 0 \qquad \nabla_{B_\epsilon^+} \varphi_\epsilon = 0 \qquad \text{totally skew torsion } H_\epsilon$$

(Note:  $B_\epsilon^-$  with torsion  $-H_\epsilon$  is “Hull connection”)

**Idea:** let  $\kappa = \kappa(\alpha') > 0$  and take  $B = B_{\kappa\epsilon}^+$ : Bismut connection for  $\varphi_{\kappa\epsilon}$

Take  $\kappa^2 = (\alpha')^{-3}$  and  $\epsilon^2 = 2(\alpha')^5 \rightsquigarrow$

- $F_B \wedge * \varphi_\epsilon = O(\alpha')^2 \checkmark$
- $dH_\epsilon = \frac{\alpha'}{4} (\text{tr } F_A^2 - \text{tr } F_B^2) \checkmark$

**Note:** we can also

- modify Hull and Levi-Civita connections for  $\varphi_{\kappa\epsilon}$
- make  $B$  approximate  $G_2$ -instanton to order  $O(\alpha')^k$  for any  $k \geq 2$

## Summary and questions

**Summary:**  $M^7$  contact Calabi–Yau  $\rightsquigarrow$  natural coclosed  $G_2$ -structures

- Laplacian flow  $\rightsquigarrow$  rapidly forming finite-time singularity, collapsing to  $\mathbb{R}$
- Laplacian coflow  $\rightsquigarrow$  slowly forming infinite-time singularity, collapsing to  $\mathbb{C}^3$
- solution to heterotic  $G_2$  system for any  $\alpha' > 0$  (“nearly collapsed”)

### Questions

- gauge theory on contact Calabi–Yau 7-manifolds? (cf. Sá Earp et. al., Y. Wang)
- other  $\mathcal{S}^1$ -bundles/ $\mathcal{S}^1$ -symmetry? (cf. Apostolov–Salamon, Foscolo–Haskins–Nordström, Fowdar)
- coclosed  $G_2$ -structures and geometric flows?
- conditions for (exact) solutions/numerical approximations to heterotic  $G_2$  system?